

Solutions to Axler's Linear Algebra

JR

July 4th, 2026

Contents

1	Vector Spaces	2
1A	$\mathbb{R}^n$ and $\mathbb{C}^n$	2
1B	Definition of Vector Space	8
1C	Subspaces	13
2	Finite-Dimensional Vector Spaces	24
2A	Span and Linear Independence	24
2B	Bases	33
2C	Dimension	40
3	Linear Maps	52
3A	Vector Space of Linear Maps	52
3B	Null Spaces and Ranges	63
3C	Matrices	81

1 Vector Spaces

1A $\mathbb{R}^n$ and $\mathbb{C}^n$

Exercise 1A1

Show that $\alpha + \beta = \beta + \alpha$ for all $\alpha, \beta \in \mathbb{C}$.

Solution. Let $\alpha = a + bi$ and $\beta = c + di$ for some $a, b, c, d \in \mathbb{R}$. Then

$$\begin{aligned}\alpha + \beta &= (a + bi) + (c + di) \\ &= (a + c) + (b + d)i \\ &= (c + a) + (d + b)i \\ &= (c + di) + (a + bi) = \beta + \alpha.\end{aligned}$$

■

Exercise 1A2

Show that $(\alpha + \beta) + \gamma = \alpha + (\beta + \gamma)$ for all $\alpha, \beta, \gamma \in \mathbb{C}$.

Solution. Let $\alpha = a + bi$, $\beta = c + di$, and $\gamma = e + fi$ for some $a, b, c, d, e, f \in \mathbb{R}$. Then

$$\begin{aligned}(\alpha + \beta) + \gamma &= ((a + bi) + (c + di)) + (e + fi) \\ &= ((a + c) + (b + d)i) + (e + fi) \\ &= (a + c + e) + (b + d + f)i \\ &= (a + (c + e)) + (b + (d + f))i \\ &= (a + bi) + ((c + di) + (e + fi)) = \alpha + (\beta + \gamma).\end{aligned}$$

■

Exercise 1A3

Show that $(\alpha\beta)\gamma = \alpha(\beta\gamma)$ for all $\alpha, \beta, \gamma \in \mathbb{C}$.

Solution. Let $\alpha = a + bi$, $\beta = c + di$, and $\gamma = e + fi$ for some $a, b, c, d, e, f \in \mathbb{R}$. Then

$$\begin{aligned}(\alpha\beta)\gamma &= ((a + bi)(c + di))(e + fi) \\ &= ((ac - bd) + (ad + bc)i)(e + fi) \\ &= ((ac - bd)e - (ad + bc)f) + ((ac - bd)f + (ad + bc)e)i \\ &= (a(ce) - b(de) - a(df) - b(cf)) + (a(cf) - b(df) + a(de) + b(ce))i \\ &= \alpha(\beta\gamma).\end{aligned}$$

■

Exercise 1A4

Show that $\gamma(\alpha + \beta) = \gamma\alpha + \gamma\beta$ for all $\alpha, \beta, \gamma \in \mathbb{C}$.

Solution. Let $\alpha = a + bi$, $\beta = c + di$, and $\gamma = e + fi$ for some $a, b, c, d, e, f \in \mathbb{R}$. Then

$$\begin{aligned}
 \gamma(\alpha + \beta) &= (e + fi)((a + bi) + (c + di)) \\
 &= (e + fi)((a + c) + (b + d)i) \\
 &= (e(a + c) - f(b + d)) + (e(b + d) + f(a + c))i \\
 &= (ea - fb + ec - fd) + (eb + fa + ed + fc)i \\
 &= (ea - fb) + (eb + fa)i + (ec - fd) + (ed + fc)i \\
 &= (e + fi)(a + bi) + (e + fi)(c + di) = \gamma\alpha + \gamma\beta.
 \end{aligned}$$

■

Exercise 1A5

Show that for every $\alpha \in \mathbb{C}$, there exists a unique $\beta \in \mathbb{C}$ such that $\alpha + \beta = 0$.

Solution. Let $\alpha = a + bi$ for some $a, b \in \mathbb{R}$. We want to find $\beta = c + di$ such that $\alpha + \beta = 0$. This gives

$$(a + bi) + (c + di) = (a + c) + (b + d)i = 0 + 0i.$$

Solving for c and d , we get $c = -a$ and $d = -b$. Hence, $\beta = c + di = -a - bi = -\alpha$.

To see that β is unique, suppose there is another $\gamma \in \mathbb{C}$ such that $\alpha + \gamma = 0$. Then

$$\beta = \beta + 0 = \beta + (\alpha + \gamma) = (\beta + \alpha) + \gamma = 0 + \gamma = \gamma.$$

Hence, β is unique. ■

Exercise 1A6

Show that for every $\alpha \in \mathbb{C}$ with $\alpha \neq 0$, there exists a unique $\beta \in \mathbb{C}$ such that $\alpha\beta = 1$.

Solution. Let $\alpha = a + bi$ for some $a, b \in \mathbb{R}$ with $\alpha \neq 0$. We want to find $\beta = c + di$ such that $\alpha\beta = 1$. This gives

$$(a + bi)(c + di) = (ac - bd) + (ad + bc)i = 1 + 0i.$$

Solving for c and d , we get the system of equations

$$\begin{cases} ac - bd = 1 \\ ad + bc = 0 \end{cases} \implies \begin{cases} a^2c - abd = a \\ abd + b^2c = 0 \end{cases}$$

Adding the two equations, we get

$$(a^2 + b^2)c = a \implies c = \frac{a}{a^2 + b^2}.$$

Substituting this into the second equation, we get

$$abd + b^2 \frac{a}{a^2 + b^2} = 0 \implies d = \frac{-b}{a^2 + b^2}.$$

Hence,

$$\beta = c + di = \frac{a}{a^2 + b^2} + \frac{-b}{a^2 + b^2}i.$$

If $\gamma \in \mathbb{C}$ is another complex number such that $\alpha\gamma = 1$, then

$$\gamma = \gamma 1 = \gamma(\alpha\beta) = (\gamma\alpha)\beta = 1\beta = \beta.$$

Hence, β is unique. ■

Exercise 1A7

Show that

$$\frac{-1 + \sqrt{3}i}{2}$$

is a cube root of 1.

Solution. Note that the complex number given can be written as

$$\frac{1}{2}(-1 + \sqrt{3}i).$$

Cubing $-1 + \sqrt{3}i$, we get

$$\begin{aligned} (-1 + \sqrt{3}i)^3 &= (-1)^3 + 3(-1)^2(\sqrt{3}i) + 3(-1)(\sqrt{3}i)^2 + (\sqrt{3}i)^3 \\ &= -1 + 3\sqrt{3}i + 3(-1)(-3) + (\sqrt{3}i)^3 \\ &= -1 + 3\sqrt{3}i + 9 + 3\sqrt{3}i(-i) \\ &= -1 + 3\sqrt{3}i + 9 - 3\sqrt{3}i \\ &= 8. \end{aligned}$$

Therefore,

$$\left(\frac{-1 + \sqrt{3}i}{2}\right)^3 = \frac{(-1 + \sqrt{3}i)^3}{8} = \frac{8}{8} = 1. \quad \blacksquare$$

Exercise 1A8

Find two distinct square roots of i .

Solution. Let $z = a + bi$ for some $a, b \in \mathbb{R}$. We want to find z such that $z^2 = i$. This gives

$$(a + bi)^2 = a^2 - b^2 + 2abi = 0 + 1i.$$

Solving for a and b , we get the system of equations

$$\begin{cases} a^2 - b^2 = 0 \\ 2ab = 1 \end{cases}$$

The first equation yields $a = b$ or $a = -b$. If $a = b$, then the second equation gives

$$2b^2 = 1 \implies b = \pm \frac{1}{\sqrt{2}} \implies a = \pm \frac{1}{\sqrt{2}}.$$

If $a = -b$, this results in $-2b^2 = 1$ which has no solution when $b \in \mathbb{R}$. Hence, the two distinct square roots of i are

$$\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i \quad \text{and} \quad -\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}}i. \quad \blacksquare$$

Exercise 1A9

Find $x \in \mathbb{R}^4$ such that

$$(4, -3, 1, 7) + 2x = (5, 9, -6, 8).$$

Solution. Let $x = (x_1, x_2, x_3, x_4) \in \mathbb{R}^4$. The equation becomes

$$(4, -3, 1, 7) + 2(x_1, x_2, x_3, x_4) = (5, 9, -6, 8),$$

which simplifies to the system of equations

$$\begin{cases} 5 = 4 + 2x_1 \\ 9 = -3 + 2x_2 \\ -6 = 1 + 2x_3 \\ 8 = 7 + 2x_4 \end{cases} \implies \begin{cases} 2x_1 = 1 \\ 2x_2 = 12 \\ 2x_3 = -7 \\ 2x_4 = 1 \end{cases}$$

Therefore,

$$x = \left(\frac{1}{2}, 6, -\frac{7}{2}, \frac{1}{2}\right). \quad \blacksquare$$

Exercise 1A10

Explain why there does not exist $\lambda \in \mathbb{C}$ such that

$$\lambda(2 - 3i, 5 + 4i, -6 + 7i) = (12 - 5i, 7 + 22i, -32 - 9i).$$

Solution. Suppose there exists $\lambda \in \mathbb{C}$ such that

$$\lambda(2 - 3i, 5 + 4i, -6 + 7i) = (12 - 5i, 7 + 22i, -32 - 9i).$$

This gives the system of equations

$$\begin{cases} \lambda(2 - 3i) = 12 - 5i \\ \lambda(5 + 4i) = 7 + 22i \\ \lambda(-6 + 7i) = -32 - 9i \end{cases}$$

Solving the first equation for λ , we get

$$\lambda = \frac{12 - 5i}{2 - 3i} = \frac{(12 - 5i)(2 + 3i)}{(2 - 3i)(2 + 3i)} = \frac{24 + 36i - 10i - 15i^2}{4 + 9} = \frac{24 + 26i + 15}{13} = 3 + 2i.$$

Solving the second equation for λ , we get

$$\lambda = \frac{7 + 22i}{5 + 4i} = \frac{(7 + 22i)(5 - 4i)}{(5 + 4i)(5 - 4i)} = \frac{35 - 28i + 110i - 88i^2}{25 + 16} = \frac{35 + 82i + 88}{41} = 3 + 2i.$$

Solving the third equation for λ , we get

$$\lambda = \frac{-32 - 9i}{-6 + 7i} = \frac{(-32 - 9i)(-6 - 7i)}{(-6 + 7i)(-6 - 7i)} = \frac{192 + 224i + 54i + 63i^2}{36 + 49} = \frac{192 + 278i - 63}{85} = \frac{129 + 278i}{85}.$$

Since the value of λ obtained from the third equation is different from the value obtained from the first two equations, no such $\lambda \in \mathbb{C}$ exists. ■

Exercise 1A11

Show that $(x + y) + z = x + (y + z)$ for all $x, y, z \in \mathbb{F}^n$.

Solution. Let $x = (x_1, x_2, \dots, x_n)$, $y = (y_1, y_2, \dots, y_n)$, and $z = (z_1, z_2, \dots, z_n)$. Then

$$\begin{aligned} (x + y) + z &= ((x_1, x_2, \dots, x_n) + (y_1, y_2, \dots, y_n)) + (z_1, z_2, \dots, z_n) \\ &= ((x_1 + y_1), (x_2 + y_2), \dots, (x_n + y_n)) + (z_1, z_2, \dots, z_n) \\ &= (x_1 + y_1 + z_1, x_2 + y_2 + z_2, \dots, x_n + y_n + z_n) \\ &= (x_1, x_2, \dots, x_n) + ((y_1 + z_1), (y_2 + z_2), \dots, (y_n + z_n)) \\ &= x + (y + z). \end{aligned}$$

■

Exercise 1A12

Show that $(ab)x = a(bx)$ for all $a, b \in \mathbb{F}$ and $x \in \mathbb{F}^n$.

Solution. Let $x = (x_1, x_2, \dots, x_n) \in \mathbb{F}^n$ and $a, b \in \mathbb{F}$. Then

$$\begin{aligned} (ab)x &= (ab)(x_1, x_2, \dots, x_n) \\ &= ((ab)x_1, (ab)x_2, \dots, (ab)x_n) \\ &= (a(bx_1), a(bx_2), \dots, a(bx_n)) \\ &= a((bx_1, bx_2, \dots, bx_n)) \\ &= a(bx). \end{aligned}$$

■

Exercise 1A13

Show that $1x = x$ for all $x \in \mathbb{F}^n$.

Solution. Let $x = (x_1, x_2, \dots, x_n) \in \mathbb{F}^n$. Then

$$\begin{aligned} 1x &= 1(x_1, x_2, \dots, x_n) \\ &= (1x_1, 1x_2, \dots, 1x_n) \\ &= (x_1, x_2, \dots, x_n) \\ &= x. \end{aligned}$$

■

Exercise 1A14

Show that $\lambda(x + y) = \lambda x + \lambda y$ for all $\lambda \in \mathbb{F}$ and $x, y \in \mathbb{F}^n$.

Solution. Let $x = (x_1, x_2, \dots, x_n)$ and $y = (y_1, y_2, \dots, y_n)$ be elements of $\mathbb{F}^n$ and $\lambda \in \mathbb{F}$. Then

$$\begin{aligned}
 \lambda(x + y) &= \lambda((x_1, x_2, \dots, x_n) + (y_1, y_2, \dots, y_n)) \\
 &= \lambda((x_1 + y_1), (x_2 + y_2), \dots, (x_n + y_n)) \\
 &= (\lambda(x_1 + y_1), \lambda(x_2 + y_2), \dots, \lambda(x_n + y_n)) \\
 &= (\lambda x_1 + \lambda y_1, \lambda x_2 + \lambda y_2, \dots, \lambda x_n + \lambda y_n) \\
 &= (\lambda x_1, \lambda x_2, \dots, \lambda x_n) + (\lambda y_1, \lambda y_2, \dots, \lambda y_n) \\
 &= \lambda(x_1, x_2, \dots, x_n) + \lambda(y_1, y_2, \dots, y_n) \\
 &= \lambda x + \lambda y.
 \end{aligned}$$

■

Exercise 1A15

Show that $(a + b)x = ax + bx$ for all $a, b \in \mathbb{F}$ and $x \in \mathbb{F}^n$.

Solution. Let $x = (x_1, x_2, \dots, x_n) \in \mathbb{F}^n$ and $a, b \in \mathbb{F}$. Then

$$\begin{aligned}
 (a + b)x &= (a + b)(x_1, x_2, \dots, x_n) \\
 &= ((a + b)x_1, (a + b)x_2, \dots, (a + b)x_n) \\
 &= (ax_1 + bx_1, ax_2 + bx_2, \dots, ax_n + bx_n) \\
 &= (ax_1, ax_2, \dots, ax_n) + (bx_1, bx_2, \dots, bx_n) \\
 &= ax + bx.
 \end{aligned}$$

■

1B Definition of Vector Space

Exercise 1B1

Prove that $-(-v) = v$ for every $v \in V$.

Solution. Let $v \in V$. By the definition of additive inverse, we have

$$v + (-v) = 0 = -(-v) + (-v).$$

Adding $-(-v)$ to both sides, we get

$$v + (-v) + (-(-v)) = -(-v) + (-v) + (-(-v)).$$

Simplifying both sides, we obtain

$$v = -(-v). \quad \blacksquare$$

Exercise 1B2

Suppose $a \in \mathbb{F}$, $v \in V$, and $av = 0$. Prove that $a = 0$ or $v = 0$.

Solution. If $a = 0$, then we are done. Suppose $a \neq 0$. Then

$$v = 1v = \frac{1}{a}(av) = \frac{1}{a} \cdot 0 = 0. \quad \blacksquare$$

Exercise 1B3

Suppose $v, w \in V$. Explain why there exists a unique $x \in V$ such that $v + 3x = w$.

Solution. Solving for x explicitly, we obtain

$$3x = w - v \implies x = \frac{1}{3}(w - v).$$

This shows that there exists $x \in V$ such that $v + 3x = w$. To see that x is unique, suppose there exists another $y \in V$ such that $v + 3y = w$. Then dividing by 3, we get

$$3x = w - v = 3y \implies x = y. \quad \blacksquare$$

Exercise 1B4

The empty set is not a vector space. The empty set fails to satisfy only one of the requirements listed in the definition of a vector space. Which one?

Solution. The empty set fails to satisfy the requirement that there exists a zero vector $0 \in V$ such that $v + 0 = v$ for all $v \in V$ since it contains no elements. ■

Exercise 1B5

Show that in the definition of a vector space, the additive inverse condition can be replaced with the condition that

$$0v = 0 \text{ for all } v \in V.$$

Here, the 0 on the left side is the number 0, and the 0 on the right side is the additive identity in V .

Solution. Suppose that $0v = 0$ for all $v \in V$. Let $v \in V$. Then

$$v + (-1)v = 1v + (-1)v = (1 + (-1))v = 0v = 0.$$

Hence, $(-1)v$ is the additive inverse of v .

Conversely, suppose that for every $v \in V$, there exists an additive inverse $-v \in V$ such that $v + (-v) = 0$. Then $0v = (0 + 0)v = 0v + 0v$. Adding $-0v$ to both sides, we get $0 = 0v$. Hence, $0v = 0$ for all $v \in V$. ■

Exercise 1B6

Let ∞ and $-\infty$ denote two distinct objects, neither of which is in $\mathbb{R}$. Define an addition and scalar multiplication on $\mathbb{R} \cup \{\infty, -\infty\}$ as you could guess from the notation. Specifically, the sum and product of two real numbers is as usual, and for $t \in \mathbb{R}$, define

$$t\infty = \begin{cases} -\infty & \text{if } t < 0, \\ 0 & \text{if } t = 0, \\ \infty & \text{if } t > 0, \end{cases} \quad t(-\infty) = \begin{cases} \infty & \text{if } t < 0, \\ 0 & \text{if } t = 0, \\ -\infty & \text{if } t > 0, \end{cases}$$

and

$$\begin{aligned} t + \infty &= \infty + t = \infty + \infty = \infty, \\ t + (-\infty) &= (-\infty) + t = (-\infty) + (-\infty) = -\infty, \\ \infty + (-\infty) &= (-\infty) + \infty = 0. \end{aligned}$$

With these operations of addition and scalar multiplication, is $\mathbb{R} \cup \{\infty, -\infty\}$ a vector space over $\mathbb{R}$? Explain.

Solution. No, $\mathbb{R} \cup \{\infty, -\infty\}$ is not a vector space over $\mathbb{R}$ as it does not satisfy the associative property of addition. Let $t \in \mathbb{R}$. Then

$$(t + \infty) + (-\infty) = \infty + (-\infty) = 0 \neq t = t + 0 = t + (\infty + (-\infty)). \quad \blacksquare$$

Exercise 1B7

Suppose S is a nonempty set. Let V^S denote the set of functions from S to V . Define a natural addition and scalar multiplication on V^S , and show that V^S is a vector space with these definitions.

Solution. Let $f, g \in V^S$ and $t \in \mathbb{F}$. Define addition and scalar multiplication as follows:

$$(f + g)(s) = f(s) + g(s), \quad (tf)(s) = t(f(s))$$

for all $s \in S$. We will show that V^S is a vector space with these operations.

1. **Commutativity:** Let $f, g \in V^S$. Then for all $s \in S$,

$$(f + g)(s) = f(s) + g(s) = g(s) + f(s) = (g + f)(s).$$

2. **Associativity:** Let $f, g, h \in V^S$ and $t, u \in \mathbb{F}$. Then for all $s \in S$,

$$\begin{aligned} ((f + g) + h)(s) &= (f + g)(s) + h(s) \\ &= (f(s) + g(s)) + h(s) \\ &= f(s) + (g(s) + h(s)) \\ &= f(s) + (g + h)(s) \\ &= (f + (g + h))(s), \end{aligned}$$

and

$$\begin{aligned} ((tu)f)(s) &= (tu)(f(s)) \\ &= t(u(f(s))) \\ &= t(uf)(s) \\ &= (t(uf))(s). \end{aligned}$$

3. **Additive identity:** Let $0 \in V$ be the additive identity in V . Define the zero function $0_S \in V^S$ by $0_S(s) = 0$ for all $s \in S$. Then for all $f \in V^S$ and $s \in S$,

$$(f + 0_S)(s) = f(s) + 0_S(s) = f(s) + 0 = f(s) = 0 + f(s) = (0_S + f)(s).$$

Hence, 0_S is the additive identity in V^S .

4. **Additive inverse:** Let $f \in V^S$. Define the function $-f \in V^S$ by $(-f)(s) = -f(s)$ for all $s \in S$. Then for all $s \in S$,

$$(f + (-f))(s) = f(s) + (-f)(s) = f(s) + (-f(s)) = 0 = 0_S(s).$$

Hence, $-f$ is the additive inverse of f in V^S .

5. **Multiplicative identity:** For all $f \in V^S$ and $s \in S$,

$$(1f)(s) = 1(f(s)) = f(s).$$

6. **Distributive properties:** Let $f, g \in V^S$ and $t, u \in \mathbb{F}$. Then for all $s \in S$,

$$\begin{aligned} ((t + u)f)(s) &= (t + u)(f(s)) \\ &= t(f(s)) + u(f(s)) \\ &= (tf)(s) + (uf)(s) \\ &= (tf + uf)(s), \end{aligned}$$

and

$$\begin{aligned}
 (t(f + g))(s) &= t((f + g)(s)) \\
 &= t(f(s) + g(s)) \\
 &= t(f(s)) + t(g(s)) \\
 &= (tf)(s) + (tg)(s) \\
 &= (tf + tg)(s).
 \end{aligned}$$

■

Exercise 1B8

Suppose V is a real vector space.

- The **complexification** of V , denoted by $V_{\mathbb{C}}$, equals $V \times V$. An element of $V_{\mathbb{C}}$ is an ordered pair (u, v) , where $u, v \in V$, but we write this as $u + iv$.
- Addition on $V_{\mathbb{C}}$ is defined by

$$(u_1 + iv_1) + (u_2 + iv_2) = (u_1 + u_2) + i(v_1 + v_2)$$

for all $u_1, u_2, v_1, v_2 \in V$.

- Complex scalar multiplication on $V_{\mathbb{C}}$ is defined by

$$(a + ib)(u + iv) = (au - bv) + i(av + bu)$$

for all $a, b \in \mathbb{R}$ and $u, v \in V$.

Prove that with the definitions of addition and scalar multiplication as above, $V_{\mathbb{C}}$ is a complex vector space.

Solution. We prove that $V_{\mathbb{C}}$ is a complex vector space by verifying the vector space axioms.

1. **Commutativity:** Let $u_1 + v_1i, u_2 + v_2i \in V_{\mathbb{C}}$. Then

$$(u_1 + v_1i) + (u_2 + v_2i) = (u_1 + u_2) + i(v_1 + v_2) = (u_2 + u_1) + i(v_2 + v_1) = (u_2 + v_2i) + (u_1 + v_1i).$$

2. **Associativity:** Let $u_1 + v_1i, u_2 + v_2i, u_3 + v_3i \in V_{\mathbb{C}}$. Then

$$\begin{aligned}
 ((u_1 + v_1i) + (u_2 + v_2i)) + (u_3 + v_3i) &= ((u_1 + u_2) + i(v_1 + v_2)) + (u_3 + v_3i) \\
 &= ((u_1 + u_2) + u_3) + i((v_1 + v_2) + v_3) \\
 &= (u_1 + (u_2 + u_3)) + i(v_1 + (v_2 + v_3)) \\
 &= (u_1 + v_1i) + ((u_2 + v_2i) + (u_3 + v_3i)).
 \end{aligned}$$

Similarly, let $a + bi, c + di \in \mathbb{C}$ and $u + vi \in V_{\mathbb{C}}$. Then

$$\begin{aligned}
 ((a + bi)(c + di))(u + vi) &= ((ac - bd) + i(ad + bc))(u + vi) \\
 &= ((ac - bd)u - (ad + bc)v) + i((ac - bd)v + (ad + bc)u) \\
 &= (a(cu - dv) - b(du + cv)) + i(a(du + cv) + b(cu - dv)) \\
 &= (a + bi)((cu - dv) + i(du + cv)) \\
 &= (a + bi)((c + di)(u + vi)).
 \end{aligned}$$

3. **Additive identity:** Let $0 \in V$ be the additive identity in V . Then for all $u + vi \in V_{\mathbb{C}}$,

$$(u + vi) + (0 + 0i) = (u + 0) + i(v + 0) = u + vi = (0 + 0i) + (u + vi).$$

Hence, $0 + 0i$ is the additive identity in $V_{\mathbb{C}}$.

4. **Additive inverse:** Let $u + vi \in V_{\mathbb{C}}$. Then

$$(u + vi) + (-u + (-v)i) = (u - u) + i(v - v) = 0 + 0i.$$

Hence, $-u + (-v)i$ is the additive inverse of $u + vi$ in $V_{\mathbb{C}}$.

5. **Multiplicative identity:** For all $u + vi \in V_{\mathbb{C}}$,

$$(1 + 0i)(u + vi) = (1u - 0v) + i(1v + 0u) = u + vi.$$

Hence, $1 + 0i$ is the multiplicative identity in $V_{\mathbb{C}}$.

6. **Distributive properties:** Let $a + bi, c + di \in \mathbb{C}$ and $u_1 + v_1i, u_2 + v_2i \in V_{\mathbb{C}}$. Then

$$\begin{aligned} (a + bi)((u_1 + v_1i) + (u_2 + v_2i)) &= (a + bi)((u_1 + u_2) + i(v_1 + v_2)) \\ &= (a(u_1 + u_2) - b(v_1 + v_2)) + i(a(v_1 + v_2) + b(u_1 + u_2)) \\ &= (au_1 - bv_1) + i(av_1 + bu_1) + (au_2 - bv_2) + i(av_2 + bu_2) \\ &= (a + bi)(u_1 + v_1i) + (a + bi)(u_2 + v_2i), \end{aligned}$$

and

$$\begin{aligned} ((a + bi) + (c + di))(u_1 + v_1i) &= ((a + c) + (b + d)i)(u_1 + v_1i) \\ &= ((a + c)u_1 - (b + d)v_1) + i((a + c)v_1 + (b + d)u_1) \\ &= (au_1 - bv_1) + i(av_1 + bu_1) + (cu_1 - dv_1) + i(cv_1 + du_1) \\ &= (a + bi)(u_1 + v_1i) + (c + di)(u_1 + v_1i). \end{aligned} \quad \blacksquare$$

1C Subspaces

Exercise 1C1

For each of the following subsets of $\mathbb{F}^3$, determine whether it is a subspace of $\mathbb{F}^3$.

- (a) $\{(x_1, x_2, x_3) \in \mathbb{F}^3 \mid x_1 + 2x_2 + 3x_3 = 0\}$
- (b) $\{(x_1, x_2, x_3) \in \mathbb{F}^3 \mid x_1 + 2x_2 + 3x_3 = 4\}$
- (c) $\{(x_1, x_2, x_3) \in \mathbb{F}^3 \mid x_1x_2x_3 = 0\}$
- (d) $\{(x_1, x_2, x_3) \in \mathbb{F}^3 \mid x_1 = 5x_3\}$

Solution. Let U be the subset of $\mathbb{F}^3$ in question.

- (a) Since the zero vector $(0, 0, 0)$ satisfies $0 + 2(0) + 3(0) = 0$, U is nonempty. Let $u = (u_1, u_2, u_3)$ and $v = (v_1, v_2, v_3)$ be elements of U . Then

$$(u_1 + v_1) + 2(u_2 + v_2) + 3(u_3 + v_3) = (u_1 + 2u_2 + 3u_3) + (v_1 + 2v_2 + 3v_3) = 0 + 0 = 0.$$

Hence, $u + v$ is in U . Let $a \in \mathbb{F}$. Then

$$au_1 + 2(au_2) + 3(au_3) = a(u_1 + 2u_2 + 3u_3) = a \cdot 0 = 0.$$

Hence, au is in U . Therefore, U is a subspace of $\mathbb{F}^3$.

- (b) The zero vector $(0, 0, 0)$ does not satisfy $0 + 2(0) + 3(0) = 4$, so U does not contain the zero vector. Hence, it is not a subspace of $\mathbb{F}^3$.
- (c) Suppose $u = (1, 0, 0)$ and $v = (0, 1, 1)$. Then u and v are in U since $1 \cdot 0 \cdot 0 = 0$ and $0 \cdot 1 \cdot 1 = 0$. However, $u + v = (1, 1, 1)$ is not in U since $1 \cdot 1 \cdot 1 = 1 \neq 0$. Hence, U is not closed under addition and is not a subspace of $\mathbb{F}^3$.
- (d) Since the zero vector $(0, 0, 0)$ satisfies $0 = 5(0)$, U is nonempty. Let $u = (u_1, u_2, u_3)$ and $v = (v_1, v_2, v_3)$ be elements of U . Then

$$u_1 + v_1 = 5u_3 + 5v_3 = 5(u_3 + v_3).$$

Hence, $u + v$ is in U . Let $a \in \mathbb{F}$. Then

$$au_1 = a(5u_3) = 5(au_3).$$

Hence, au is in U . Therefore, U is a subspace of $\mathbb{F}^3$. ■

Exercise 1C2

Verify all assertions about subspaces in Example 1.35.

- (a) If $b \in \mathbb{F}$, then

$$\{(x_1, x_2, x_3, x_4) \in \mathbb{F}^4 \mid x_3 = 5x_4 + b\}$$

is a subspace of $\mathbb{F}^4$ if and only if $b = 0$.

- (b) The set of continuous, real-valued functions on the interval $[0, 1]$ is a subspace of $\mathbb{R}^{[0,1]}$.
 (c) The set of differentiable real-valued functions on $\mathbb{R}$ is a subspace of $\mathbb{R}^{\mathbb{R}}$.
 (d) The set of differentiable real-valued functions f on the interval $(0, 3)$ such that $f'(2) = b$ is a subspace of $\mathbb{R}^{(0,3)}$ if and only if $b = 0$.
 (e) The set of all sequences of complex numbers with limit 0 is a subspace of $\mathbb{C}^{\infty}$.

Solution. Let U be the subset of $\mathbb{F}^4$ in question.

- (a) Suppose U is a subspace of $\mathbb{F}^4$. Then it must contain the zero vector $(0, 0, 0, 0)$. Then $0 = 5(0) + b$, which implies $b = 0$.

Conversely, suppose $b = 0$. Then $U = \{(x_1, x_2, x_3, x_4) \in \mathbb{F}^4 \mid x_3 = 5x_4\}$. The proof is similar to that of [Exercise 1C1](#) (d), hence U is a subspace of $\mathbb{F}^4$.

- (b) Note that the 0 function on $[0, 1]$ is continuous, so $0 \in U$. Let $f, g \in U$ be continuous functions. Then $f + g$ is continuous, so $f + g \in U$. Let $a \in \mathbb{R}$. Then af is continuous, so $af \in U$. Therefore, U is a subspace of $\mathbb{R}^{[0,1]}$.

- (c) Since the derivative of the 0 function is the 0 function, $0 \in U$. Let $f, g \in U$ be differentiable functions. Then $f + g$ is differentiable, so $f + g \in U$. Let $a \in \mathbb{R}$. Then af is differentiable, so $af \in U$. Therefore, U is a subspace of $\mathbb{R}^{\mathbb{R}}$.

- (d) Suppose U is a subspace of $\mathbb{R}^{(0,3)}$. Then it must contain the 0 function on $(0, 3)$. Then $0' = 0 = b$, which implies $b = 0$.

Conversely, suppose $b = 0$. Then $U = \{f \in \mathbb{R}^{(0,3)} \mid f'(2) = 0\}$. Let $f, g \in U$ be differentiable functions such that $f'(2) = g'(2) = 0$. Then

$$(f + g)'(2) = f'(2) + g'(2) = 0 + 0 = 0,$$

so $f + g \in U$. Let $a \in \mathbb{R}$. Then

$$(af)'(2) = af'(2) = a \cdot 0 = 0,$$

so $af \in U$. Therefore, U is a subspace of $\mathbb{R}^{(0,3)}$.

- (e) Let U be the set of all sequences of complex numbers with limit 0. Then the zero sequence is in U . Let $(z_n), (w_n) \in U$ be sequences with limit 0. Then

$$\lim_{n \rightarrow \infty} (z_n + w_n) = \lim_{n \rightarrow \infty} z_n + \lim_{n \rightarrow \infty} w_n = 0 + 0 = 0,$$

so $(z_n) + (w_n) \in U$. Let $a \in \mathbb{C}$. Then

$$\lim_{n \rightarrow \infty} (az_n) = a \lim_{n \rightarrow \infty} z_n = a \cdot 0 = 0,$$

so $a(z_n) \in U$. Therefore, U is a subspace of $\mathbb{C}^\infty$. ■

Exercise 1C3

Show that the set of differentiable real-valued functions f on the interval $(-4, 4)$ such that $f'(-1) = 3f(2)$ is a subspace of $\mathbb{R}^{(-4, 4)}$.

Solution. Let U be the set in question. Then the zero function is in U since $0'(-1) = 0 = 3 \cdot 0 = 3 \cdot 0(2)$. Let $f, g \in U$ be differentiable functions such that $f'(-1) = 3f(2)$ and $g'(-1) = 3g(2)$. Then

$$(f + g)'(-1) = f'(-1) + g'(-1) = 3f(2) + 3g(2) = 3(f(2) + g(2)) = 3(f + g)(2),$$

so $f + g \in U$. Let $a \in \mathbb{R}$. Then

$$(af)'(-1) = af'(-1) = a \cdot 3f(2) = 3(af)(2),$$

so $af \in U$. Therefore, U is a subspace of $\mathbb{R}^{(-4, 4)}$. ■

Exercise 1C4

Suppose $b \in \mathbb{R}$. Show that the set of continuous real-valued functions f on the interval $[0, 1]$ such that

$$\int_0^1 f = b$$

is a subspace of $\mathbb{R}^{[0, 1]}$ if and only if $b = 0$.

Solution. Let U be the set in question. Suppose U is a subspace of $\mathbb{R}^{[0, 1]}$. Then it must contain the zero function on $[0, 1]$. Then

$$\int_0^1 0 = 0 = b,$$

which implies $b = 0$.

Conversely, suppose $b = 0$. Then

$$U = \left\{ f \in \mathbb{R}^{[0, 1]} \mid \int_0^1 f = 0 \right\}.$$

The zero function is in U since $\int_0^1 0 = 0$. Let $f, g \in U$ be continuous functions such that $\int_0^1 f = \int_0^1 g = 0$. Then

$$\int_0^1 (f + g) = \int_0^1 f + \int_0^1 g = 0 + 0 = 0,$$

so $f + g \in U$. Let $a \in \mathbb{R}$. Then

$$\int_0^1 (af) = a \int_0^1 f = a \cdot 0 = 0,$$

so $af \in U$. Therefore, U is a subspace of $\mathbb{R}^{[0, 1]}$. ■

Exercise 1C5

Is $\mathbb{R}^2$ a subspace of the complex vector space $\mathbb{C}^2$?

Solution. No, $\mathbb{R}^2$ is not a subspace of $\mathbb{C}^2$. Let $u = (1, 0) \in \mathbb{R}^2$ and $a = i \in \mathbb{C}$. Then $au = (i, 0) \notin \mathbb{R}^2$, so $\mathbb{R}^2$ is not closed under scalar multiplication and is not a subspace of $\mathbb{C}^2$. ■

Exercise 1C6

(a) Is $\{(a, b, c) \in \mathbb{R}^3 \mid a^3 = b^3\}$ a subspace of $\mathbb{R}^3$?

(b) Is $\{(a, b, c) \in \mathbb{C}^3 \mid a^3 = b^3\}$ a subspace of $\mathbb{C}^3$?

Remark. Here, I present an elementary argument that does not require the use of De Moivre's Theorem; rather, I arrive to the conclusion directly. However, readers who are familiar with it may find it easier to note that the complex number 1 has three cubic roots of unity. Other readers may also use Vieta's formulas to arrive at the same polynomial in ω to obtain $1 + \omega = -\omega^2$.

Solution. Let U be the subset in question.

(a) Since the zero vector $(0, 0, 0)$ satisfies $0^3 = 0^3$, U is nonempty. Let $x = (x_1, x_2, x_3)$ and $y = (y_1, y_2, y_3)$ be elements of U . Then $x_1^3 = x_2^3$ and $y_1^3 = y_2^3$ both imply that $x_1 = x_2$ and $y_1 = y_2$. Hence,

$$(x_1 + y_1)^3 = (x_2 + y_2)^3,$$

so $x + y \in U$. Let $a \in \mathbb{R}$. Then

$$(ax_1)^3 = a^3 x_1^3 = a^3 x_2^3 = (ax_2)^3,$$

so $ax \in U$. Therefore, U is a subspace of $\mathbb{R}^3$.

(b) No, U is not a subspace of $\mathbb{C}^3$. To see this, consider the cubic polynomial $x^3 - 1 = 0$. This polynomial has three distinct roots in $\mathbb{C}$ as follows:

$$x^3 - 1 = (x - 1)(x^2 + x + 1) = 0$$

Applying the quadratic formula to $x^2 + x + 1 = 0$, we get

$$x = \frac{-1 \pm \sqrt{(-1)^2 - 4(1)(1)}}{2(1)} = \frac{-1 \pm i\sqrt{3}}{2}.$$

Let

$$\omega = \frac{-1 + i\sqrt{3}}{2}.$$

Then $\omega^3 = 1$, and

$$\omega^2 = \frac{-1 - i\sqrt{3}}{2}.$$

Moreover, it is not hard to see that $\omega + \omega^2 = -1$. Let $u = (1, 1, 0)$ and $v = (\omega, 1, 0)$. Then $u, v \in U$ since $1^3 = 1^3$ and $\omega^3 = 1^3$. Then $u + v = (1 + \omega, 2, 0)$. However,

$$(1 + \omega)^3 = 1 + 3\omega + 3\omega^2 + \omega^3 = 1 + 3(-1) + 1 = -1 \neq 2^3 = 8.$$

Hence, $u + v \notin U$, so U is not closed under addition and is not a subspace of $\mathbb{C}^3$. ■

Exercise 1C7

Prove or give a counterexample: If U is a nonempty subset of $\mathbb{R}^2$ such that U is closed under addition and under taking additive inverses (meaning $-u \in U$ whenever $u \in U$), then U is a subspace of $\mathbb{R}^2$.

Solution. Consider the subset $\mathbb{Q}^2$ of $\mathbb{R}^2$. Then $\mathbb{Q}^2$ is nonempty because $(0, 0) \in \mathbb{Q}^2$, closed under addition since the sum of any two rational numbers is rational, and closed under taking additive inverses since the additive inverse of a rational number is rational. However, for any irrational number $r \in \mathbb{R}$, the vector $(r, 0) \in \mathbb{R}^2$ is not in $\mathbb{Q}^2$. Hence, $\mathbb{Q}^2$ is not closed under scalar multiplication and is not a subspace of $\mathbb{R}^2$. Therefore, the statement is false. ■

Exercise 1C8

Give an example of a nonempty subset U of $\mathbb{R}^2$ such that U is closed under scalar multiplication, but U is not a subspace of $\mathbb{R}^2$.

Solution. Consider the subset

$$U = \{(x, 0) \mid x \in \mathbb{R}\} \cup \{(0, y) \mid y \in \mathbb{R}\}.$$

Then U is nonempty since $(1, 0) \in U$, and it is closed under scalar multiplication since any scalar multiple of $(1, 0)$ or $(0, 1)$ is still in U . However, U is not closed under addition since $(1, 0) + (0, 1) = (1, 1) \notin U$. Hence, U is not a subspace of $\mathbb{R}^2$. ■

Exercise 1C9

A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is called *periodic* if there exists a positive number p such that $f(x+p) = f(x)$ for all $x \in \mathbb{R}$. Is the set of periodic functions from $\mathbb{R}$ to $\mathbb{R}$ a subspace of $\mathbb{R}^{\mathbb{R}}$? Explain.

Solution. It is not a subspace of $\mathbb{R}^{\mathbb{R}}$. Consider the periodic functions $f(x) = \sin(2x)$ and $g(x) = \sin(\pi x)$. The function f has period π , and the function g has period 2. For $f(x+p) + g(x+p) = f(x) + g(x)$ to hold for all $x \in \mathbb{R}$, we would need a common period p that is a multiple of both π and 2. However, no such positive number exists since π is irrational. Therefore, the sum $f + g$ is not periodic, and the set of periodic functions is not closed under addition. Hence, it is not a subspace of $\mathbb{R}^{\mathbb{R}}$. ■

Exercise 1C10

Suppose V_1 and V_2 are subspaces of V . Prove that the intersection $V_1 \cap V_2$ is a subspace of V .

Solution. Let $U = V_1 \cap V_2$. Since both V_1 and V_2 are subspaces of V , they both contain the zero vector of V . Hence, $0 \in U$, so U is nonempty. Let $u, v \in U$. Then $u, v \in V_1$ and $u, v \in V_2$. Since both V_1 and V_2 are subspaces, they are closed under addition. Therefore, $u + v \in V_1$ and $u + v \in V_2$, which

implies that $u + v \in U$. Let $a \in \mathbb{F}$. Then since both V_1 and V_2 are closed under scalar multiplication, we have that $au \in V_1$ and $au \in V_2$, which implies that $au \in U$. Therefore, U is a subspace of V . ■

Exercise 1C11

Prove that the intersection of every collection of subspaces of V is a subspace of V .

Solution. Let C be a collection of subspaces of V , and let

$$U = \bigcap_{W \in C} W.$$

Since each subspace in C contains the zero vector, $0 \in U$, so U is nonempty. Let $u, v \in U$. Then $u, v \in W$ for all $W \in C$. Since each W is a subspace, they are closed under addition, so $u + v \in W$ for all $W \in C$, which implies $u + v \in U$. Let $a \in \mathbb{F}$. Then $au \in W$ for all $W \in C$, which implies $au \in U$. Therefore, U is a subspace of V . ■

Exercise 1C12

Prove that the union of two subspaces of V is a subspace of V if and only if one of the subspaces is contained in the other.

Solution. Let U, W be subspaces of V . Suppose $U \subseteq W$. Then $U \cup W = W$, which is a subspace of V . Similarly, if $W \subseteq U$, then $U \cup W = U$, which is a subspace of V .

Conversely, suppose $U \cup W$ is a subspace of V . If neither $U \subseteq W$ nor $W \subseteq U$, then there exist $u \in U \setminus W$ and $w \in W \setminus U$. Since $U \cup W$ is a subspace, it must be closed under addition. Then $u + w \in U \cup W$. However, $u + w \notin U$ because if it were, then $w = (u + w) - u \in U$, which is a contradiction. Similarly, $u + w \notin W$ because if it were, then $u = (u + w) - w \in W$, which is also a contradiction. Hence, $u + w \notin U \cup W$, which contradicts the assumption that $U \cup W$ is a subspace. Therefore, one of the subspaces must be contained in the other. ■

Exercise 1C13

Prove that the union of three subspaces of V is a subspace of V if and only if one of the subspaces contains the other two.

Solution. Let U_1, U_2, U_3 be subspaces of V . Without loss of generality, suppose $U_2 \subseteq U_1$ and $U_3 \subseteq U_1$. Then $U_1 \cup U_2 \cup U_3 = U_1$, which is a subspace of V .

Conversely, suppose $U_1 \cup U_2 \cup U_3$ is a subspace of V . Note that $U_1 \cup U_2 \cup U_3 = (U_1 \cup U_2) \cup U_3$. Using [Exercise 1C12](#), one of the subspaces $U_1 \cup U_2$ or U_3 must be contained in the other. If $U_1 \cup U_2 \subseteq U_3$, then $U_1 \cup U_2 \cup U_3 = U_3$, which is a subspace of V . If $U_3 \subseteq U_1 \cup U_2$, then $U_1 \cup U_2$ must be a subspace of V . By [Exercise 1C12](#) again, then either $U_1 \subseteq U_2$ or $U_2 \subseteq U_1$. If the former is true, then $U_1 \cup U_2 = U_2$, hence U_2 contains U_1 and U_3 . If the latter is true, then $U_1 \cup U_2 = U_1$, hence U_1 contains U_2 and U_3 . Therefore, one of the subspaces must contain the other two. ■

Exercise 1C14

Suppose

$$U = \{(x, -x, 2x) \in \mathbb{F}^3 \mid x \in \mathbb{F}\} \quad \text{and} \quad W = \{(x, x, 2x) \in \mathbb{F}^3 \mid x \in \mathbb{F}\}.$$

Describe $U + W$ using symbols, and also give a description of $U + W$ that uses no symbols.

Solution. Let $u = (x, -x, 2x) \in U$ and $w = (y, y, 2y) \in W$. Then

$$u + w = (x + y, -x + y, 2x + 2y) = (x + y, y - x, 2(x + y)).$$

Observe that $x + y$ can take any value in $\mathbb{F}$, and $y - x$ can also take any value in $\mathbb{F}$. Hence, we can write

$$U + W = \{(a, b, 2a) \in \mathbb{F}^3 \mid a, b \in \mathbb{F}\}.$$

In words, $U + W$ is the set of all vectors in $\mathbb{F}^3$ whose third component is twice the first component, and whose second component can be any value in $\mathbb{F}$. ■

Exercise 1C15

Suppose U is a subspace of V . What is $U + U$?

Solution. Let $u_1, u_2 \in U$. Then $u_1 + u_2 \in U$ since U is closed under addition. Hence, $U + U \subseteq U$. Conversely, let $u \in U$. Then $u = u + 0$, where 0 is the additive identity in U . Since $0 \in U$, we have that $u \in U + U$. Hence, $U \subseteq U + U$. Therefore, $U + U = U$. ■

Exercise 1C16

Is the operation of addition on subspaces of V commutative? In other words, if U and W are subspaces of V , is $U + W = W + U$?

Solution. Yes, the operation of addition on subspaces of V is commutative. Let $u \in U$ and $w \in W$. Then $u + w \in U + W$. Because addition in V is commutative, we have that $u + w = w + u$, where $w \in W$ and $u \in U$. Hence, $w + u \in W + U$. Therefore, every element of $U + W$ is also an element of $W + U$, which implies that $U + W \subseteq W + U$. By a similar argument, we can show that $W + U \subseteq U + W$. Hence, $U + W = W + U$. ■

Exercise 1C17

Is the operation of addition on the subspaces of V associative? In other words, if V_1, V_2, V_3 are subspaces of V , is

$$(V_1 + V_2) + V_3 = V_1 + (V_2 + V_3)?$$

Solution. Yes, the operation of addition on subspaces of V is associative. Let $u_1 \in V_1$, $u_2 \in V_2$, and $u_3 \in V_3$. Then $(u_1 + u_2) + u_3 \in (V_1 + V_2) + V_3$. Because addition in V is associative, we have that $(u_1 + u_2) + u_3 = u_1 + (u_2 + u_3)$, where $u_1 \in V_1$, $u_2 \in V_2$, and $u_3 \in V_3$. Hence, $u_1 + (u_2 + u_3) \in V_1 + (V_2 + V_3)$. Therefore, every element of $(V_1 + V_2) + V_3$ is also an element of $V_1 + (V_2 + V_3)$, which implies that

$(V_1 + V_2) + V_3 \subseteq V_1 + (V_2 + V_3)$. By a similar argument, we can show that $V_1 + (V_2 + V_3) \subseteq (V_1 + V_2) + V_3$. Hence, $(V_1 + V_2) + V_3 = V_1 + (V_2 + V_3)$. ■

Exercise 1C18

Does the operation of addition on the subspaces of V have an additive identity? Which subspaces have additive inverses?

Solution. Yes, the operation of addition on the subspaces of V has an additive identity. The additive identity is the zero subspace $\{0\}$, which contains only the zero vector of V . For any subspace U of V , we have that $U + \{0\} = U$.

If a subspace U of V had an additive inverse W , then we would have that $U + W = \{0\}$. However, $U \subseteq U + W$ since $u = u + 0$ for any $u \in U$. Hence, $U + W = \{0\}$ implies that $U = \{0\}$. Therefore, the only subspace of V that has an additive inverse is the zero subspace $\{0\}$. ■

Exercise 1C19

Prove or give a counterexample: If V_1, V_2, U are subspaces of V such that

$$V_1 + U = V_2 + U,$$

then $V_1 = V_2$.

Solution. The statement is false. Consider the vector space $\mathbb{R}^2$ and let

$$V_1 = \{(x, 0) \in \mathbb{R}^2 \mid x \in \mathbb{R}\}, \quad V_2 = \{(0, y) \in \mathbb{R}^2 \mid y \in \mathbb{R}\}, \quad U = \{(x, x) \in \mathbb{R}^2 \mid x \in \mathbb{R}\}.$$

Then $V_1 + U = \mathbb{R}^2$ and $V_2 + U = \mathbb{R}^2$, but $V_1 \neq V_2$. Therefore, the statement is false. ■

Exercise 1C20

Suppose

$$U = \{(x, x, y, y) \in \mathbb{F}^4 \mid x, y \in \mathbb{F}\}.$$

Find a subspace W of $\mathbb{F}^4$ such that $\mathbb{F}^4 = U \oplus W$.

Solution. Let

$$W = \{(0, z, 0, w) \in \mathbb{F}^4 \mid z, w \in \mathbb{F}\}.$$

If $(x, x, y, y) \in U$ and $(0, z, 0, w) \in W$, then $(x, x, y, y) = (0, z, 0, w)$ if and only if $x = 0$, $y = 0$, $z = 0$, and $w = 0$. Hence, $U \cap W = \{0\}$. Moreover, for any $(a, b, c, d) \in \mathbb{F}^4$, we can write

$$(a, b, c, d) = (x, x, y, y) + (0, z, 0, w),$$

where

$$x = a, \quad y = c, \quad z = b - x = b - a, \quad w = d - y = d - c.$$

Hence, every vector in $\mathbb{F}^4$ can be written as a sum of a vector in U and a vector in W . Therefore, $\mathbb{F}^4 = U \oplus W$. ■

Exercise 1C21

Suppose

$$U = \{(x, y, x + y, x - y, 2x) \in \mathbb{F}^5 \mid x, y \in \mathbb{F}\}.$$

Find a subspace W of $\mathbb{F}^5$ such that $\mathbb{F}^5 = U \oplus W$.

Solution. Let

$$W = \{(0, 0, z, w, v) \in \mathbb{F}^5 \mid z, w, v \in \mathbb{F}\}.$$

If $(x, y, x + y, x - y, 2x) \in U$ and $(0, 0, z, w, v) \in W$, then $(x, y, x + y, x - y, 2x) = (0, 0, z, w, v)$ if and only if $x = 0$, $y = 0$, $z = 0$, $w = 0$, and $v = 0$. Hence, $U \cap W = \{0\}$. Moreover, for any $(a, b, c, d, e) \in \mathbb{F}^5$, we can write

$$(a, b, c, d, e) = (x, y, x + y, x - y, 2x) + (0, 0, z, w, v),$$

which implies

$$\begin{cases} a = x + 0 \\ b = y + 0 \\ c = (x + y) + z \\ d = (x - y) + w \\ e = 2x + v \end{cases} \implies \begin{cases} x = a \\ y = b \\ z = c - (x + y) = c - (a + b) \\ w = d - (x - y) = d - (a - b) \\ v = e - 2x = e - 2a \end{cases}$$

Hence, every vector in $\mathbb{F}^5$ can be written as a sum of a vector in U and a vector in W . Therefore, $\mathbb{F}^5 = U \oplus W$. ■

Exercise 1C22

Suppose

$$U = \{(x, y, x + y, x - y, 2x) \in \mathbb{F}^5 \mid x, y \in \mathbb{F}\}.$$

Find three subspaces W_1, W_2, W_3 of $\mathbb{F}^5$, none of which equals $\{0\}$, such that $\mathbb{F}^5 = U \oplus W_1 \oplus W_2 \oplus W_3$.

Solution. Let

$$W_1 = \{(0, 0, z, 0, 0) \in \mathbb{F}^5 \mid z \in \mathbb{F}\},$$

$$W_2 = \{(0, 0, 0, w, 0) \in \mathbb{F}^5 \mid w \in \mathbb{F}\},$$

$$W_3 = \{(0, 0, 0, 0, v) \in \mathbb{F}^5 \mid v \in \mathbb{F}\}.$$

The intersection of any two of these subspaces is $\{0\}$, since they have no nonzero vectors in common. Additionally, the intersection of W_1, W_2, W_3 with U is also $\{0\}$, since any nonzero vector in U has nonzero entries in the first two coordinates, while any vector in W_1, W_2 , or W_3 has zeros in the first two coordinates. The proof of showing that $\mathbb{F}^5 = U \oplus W_1 \oplus W_2 \oplus W_3$ is similar to that of [Exercise 1C21](#), hence omitted. Therefore, $\mathbb{F}^5 = U \oplus W_1 \oplus W_2 \oplus W_3$. ■

Exercise 1C23

Prove or give a counterexample: If V_1, V_2, U are subspaces of V such that

$$V = V_1 \oplus U \quad \text{and} \quad V = V_2 \oplus U,$$

then $V_1 = V_2$.

Solution. See [Exercise 1C19](#) for a counterexample. For $V_1 \oplus U$, observe that $V_1 \cap U = \{0\}$. Moreover, for any $(a, b) \in \mathbb{R}^2$, we can write

$$(a, b) = (a - b, 0) + (b, b) \in V_1 \oplus U.$$

The proof for $V_2 \oplus U$ is similar. Hence, $V = V_1 \oplus U = V_2 \oplus U$, but $V_1 \neq V_2$. Therefore, the statement is false. ■

Exercise 1C24

A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is called *even* if

$$f(-x) = f(x)$$

for all $x \in \mathbb{R}$. A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is called *odd* if

$$f(-x) = -f(x)$$

for all $x \in \mathbb{R}$. Let V_e denote the set of real-valued even functions on $\mathbb{R}$, and let V_o denote the set of real-valued odd functions on $\mathbb{R}$. Show that $\mathbb{R}^{\mathbb{R}} = V_e \oplus V_o$.

Solution. Let $f \in \mathbb{R}^{\mathbb{R}}$. Define

$$f_e(x) = \frac{f(x) + f(-x)}{2} \quad \text{and} \quad f_o(x) = \frac{f(x) - f(-x)}{2}.$$

Then f_e is even since

$$f_e(-x) = \frac{f(-x) + f(x)}{2} = f_e(x),$$

and f_o is odd since

$$f_o(-x) = \frac{f(-x) - f(x)}{2} = -\frac{f(x) - f(-x)}{2} = -f_o(x).$$

Moreover, we have that

$$f(x) = f_e(x) + f_o(x).$$

Hence, every function in $\mathbb{R}^{\mathbb{R}}$ can be written as a sum of an even function and an odd function.

Now, suppose $g \in V_e \cap V_o$. Then g is both even and odd, which implies that

$$g(0) = g(-0) = -g(0),$$

so $g(0) = 0$. For any $x \in \mathbb{R}$, we have that

$$g(x) = g(-(-x)) = -g(-x) = -g(x),$$

which implies that $g(x) = 0$. Hence, the only function in the intersection of V_e and V_o is the zero function. Therefore, $\mathbb{R}^{\mathbb{R}} = V_e \oplus V_o$. ■

2 Finite-Dimensional Vector Spaces

2A Span and Linear Independence

Exercise 2A1

Find a list of four distinct vectors in $\mathbb{F}^3$ whose span equals

$$\{(x, y, z) \in \mathbb{F}^3 \mid x + y + z = 0\}$$

Solution. Clearly, the span of the list involves two vectors, since we may rewrite z in terms of x and y as $z = -x - y$. Thus, we may choose the following two vectors:

$$v_1 = (1, 0, -1), \quad v_2 = (0, 1, -1).$$

To find two more vectors, we may pick any two distinct linear combinations of v_1 and v_2 . For example, we may select

$$v_3 = v_1 + v_2 = (1, 1, -2), \quad v_4 = 2v_1 + 3v_2 = (2, 3, -5).$$

Then

$$\text{span}(v_1, v_2, v_3, v_4) = \text{span}(v_1, v_2) = \{(x, y, z) \in \mathbb{F}^3 \mid x + y + z = 0\}. \quad \blacksquare$$

Exercise 2A2

Prove or give a counterexample: If v_1, v_2, v_3, v_4 spans V , then the list

$$v_1 - v_2, v_2 - v_3, v_3 - v_4, v_4$$

also spans V .

Solution. Let $U = \{v_1, v_2, v_3, v_4\}$ and $W = \{v_1 - v_2, v_2 - v_3, v_3 - v_4, v_4\}$. Clearly, $v_4 \in \text{span } W$. Then we may write

$$v_3 = (v_3 - v_4) + v_4 \in \text{span } W,$$

$$v_2 = (v_2 - v_3) + v_3 \in \text{span } W,$$

$$v_1 = (v_1 - v_2) + v_2 \in \text{span } W.$$

Thus, $v_1, v_2, v_3, v_4 \in \text{span } W$, and so $V = \text{span } U \subseteq \text{span } W$. Since W is a list of vectors in V , we have $\text{span } W \subseteq V$. Therefore, $\text{span } W = V$, and so W spans V . $\blacksquare$

Exercise 2A3

Suppose $v_1, \dots, v_m$ is a list of vectors in V . For $k \in \{1, \dots, m\}$, let

$$w_k = v_1 + \dots + v_k.$$

Show that $\text{span}(v_1, \dots, v_m) = \text{span}(w_1, \dots, w_m)$.

Solution. This is a generalization of [Exercise 2A2](#) and a direct consequence of the fact that $v_k = w_k - w_{k-1}$ for $k \in \{2, \dots, m\}$ and $v_1 = w_1$. Since each v_k is a linear combination of $w_1, \dots, w_k$, we have $\text{span}(v_1, \dots, v_m) \subseteq \text{span}(w_1, \dots, w_m)$. On the other hand, each w_k is a linear combination of $v_1, \dots, v_k$, so $\text{span}(w_1, \dots, w_m) \subseteq \text{span}(v_1, \dots, v_m)$. Therefore, $\text{span}(v_1, \dots, v_m) = \text{span}(w_1, \dots, w_m)$. ■

Exercise 2A4

- (a) Show that a list of length one in a vector space is linearly independent if and only if the vector in the list is not 0.
- (b) Show that a list of length two in a vector space is linearly independent if and only if neither of the two vectors in the list is a scalar multiple of the other.

Solution.

- (a) Let v be the vector in the list. If $v = 0$, then the list is linearly dependent, since $0 \cdot v = 0$. If $v \neq 0$, then the only way to write 0 as a linear combination of v is to take the scalar multiple to be 0. Thus, the list is linearly independent.
- (b) Let v_1 and v_2 be the vectors in the list. If v_1 is a scalar multiple of v_2 , then there exists a scalar λ such that $v_1 = \lambda v_2$. Then we may write

$$0 = v_1 - \lambda v_2,$$

which shows that the list is linearly dependent. Conversely, if the list is linearly dependent, then there exist scalars α and β , not both zero, such that

$$\alpha v_1 + \beta v_2 = 0.$$

If $\alpha \neq 0$, then

$$v_1 = -\frac{\beta}{\alpha} v_2.$$

Similarly, if $\beta \neq 0$, then

$$v_2 = -\frac{\alpha}{\beta} v_1.$$

Thus, the list is linearly independent if and only if neither vector is a scalar multiple of the other. ■

Exercise 2A5

Find a number t such that

$$(3, 1, 4), (2, -3, 5), (5, 9, t)$$

is not linearly independent in $\mathbb{R}^3$.

Solution. To say that the list is not linearly independent is to say that there exist scalars α, β, γ , not all zero, such that

$$\alpha(3, 1, 4) + \beta(2, -3, 5) + \gamma(5, 9, t) = (0, 0, 0).$$

This is equivalent to the following system of equations:

$$\begin{aligned} 0 &= 3\alpha + 2\beta + 5\gamma, \\ 0 &= \alpha - 3\beta + 9\gamma, \\ 0 &= 4\alpha + 5\beta + t\gamma. \end{aligned}$$

From the second equation, we obtain $\alpha = 3\beta - 9\gamma$. Substituting this into the first equation, we have

$$\begin{aligned} 0 &= 3(3\beta - 9\gamma) + 2\beta + 5\gamma \\ &= 11\beta - 22\gamma. \end{aligned}$$

Thus, $\beta = 2\gamma$. Substituting this into the expression for α , we have $\alpha = 3(2\gamma) - 9\gamma = -3\gamma$. Substituting these values into the third equation, we have

$$\begin{aligned} 0 &= 4(-3\gamma) + 5(2\gamma) + t\gamma \\ &= -12\gamma + 10\gamma + t\gamma \\ &= (t - 2)\gamma. \end{aligned}$$

Since $\gamma \neq 0$, we must have $t - 2 = 0$, or $t = 2$. Then the list is not linearly independent when $t = 2$. ■

Exercise 2A6

Show that the list $(2, 3, 1), (1, -1, 2), (7, 3, c)$ is linearly dependent in $\mathbb{F}^3$ if and only if $c = 8$.

Solution. We may proceed similarly to [Exercise 2A5](#) to obtain the following system of equations:

$$\begin{aligned} 0 &= 2\alpha + \beta + 7\gamma, \\ 0 &= 3\alpha - \beta + 3\gamma, \\ 0 &= \alpha + 2\beta + c\gamma. \end{aligned}$$

The second equation gives us $\beta = 3\alpha + 3\gamma$. Substituting this into the first equation, we have

$$\begin{aligned} 0 &= 2\alpha + (3\alpha + 3\gamma) + 7\gamma \\ &= 5\alpha + 10\gamma. \end{aligned}$$

Thus, $\alpha = -2\gamma$. Substituting this into the expression for β , we have $\beta = 3(-2\gamma) + 3\gamma = -3\gamma$. Substituting these values into the third equation, we have

$$\begin{aligned} 0 &= (-2\gamma) + 2(-3\gamma) + c\gamma \\ &= -2\gamma - 6\gamma + c\gamma \\ &= (c - 8)\gamma. \end{aligned}$$

Since $\gamma \neq 0$, we must have $c - 8 = 0$, or $c = 8$. Then the list is linearly dependent when $c = 8$.

Conversely, if $c \neq 8$, then $(c - 8)\gamma = 0$ implies $\gamma = 0$. Then $\alpha = -2\gamma = 0$ and $\beta = -3\gamma = 0$, so the only solution to the system of equations is $\alpha = \beta = \gamma = 0$. Thus, the list is linearly independent when $c \neq 8$. ■

Exercise 2A7

- (a) Show that if we think of $\mathbb{C}$ as a vector space over $\mathbb{R}$, then the list $1 + i, 1 - i$ is linearly independent.
- (b) Show that if we think of $\mathbb{C}$ as a vector space over $\mathbb{C}$, then the list $1 + i, 1 - i$ is linearly dependent.

Solution.

- (a) Suppose $\alpha, \beta \in \mathbb{R}$ such that

$$\alpha(1 + i) + \beta(1 - i) = 0.$$

Then we have

$$(\alpha + \beta) + (\alpha - \beta)i = 0.$$

Since $0 = 0 + 0i$, we must have $\alpha + \beta = 0$ and $\alpha - \beta = 0$. Solving this system of equations, we find that $\alpha = \beta = 0$. Thus, the list is linearly independent.

- (b) Suppose $a + bi, c + di \in \mathbb{C}$ such that

$$(a + bi)(1 + i) + (c + di)(1 - i) = 0.$$

Then we have

$$(a - b + c + d) + (a + b - c + d)i = 0.$$

Then we have the system of equations

$$0 = a - b + c + d,$$

$$0 = a + b - c + d.$$

Adding the two equations results in $0 = 2a + 2d$, or $a = -d$. Substituting this into the first equation, we obtain $b = c$. Then we have infinitely many nontrivial solutions to the system of equations, such as $a = 1, b = 0, c = 0, d = -1$. Thus, the list is linearly dependent. ■

Exercise 2A8

Suppose v_1, v_2, v_3, v_4 is linearly independent in V . Prove that the list $v_1 - v_2, v_2 - v_3, v_3 - v_4, v_4$ is also linearly independent.

Solution. Let $\alpha, \beta, \gamma, \delta \in \mathbb{F}$ such that

$$\alpha(v_1 - v_2) + \beta(v_2 - v_3) + \gamma(v_3 - v_4) + \delta v_4 = 0.$$

Then we have

$$\alpha v_1 + (-\alpha + \beta)v_2 + (-\beta + \gamma)v_3 + (-\gamma + \delta)v_4 = 0.$$

Since v_1, v_2, v_3, v_4 is linearly independent, we must have

$$0 = \alpha,$$

$$0 = -\alpha + \beta,$$

$$0 = -\beta + \gamma,$$

$$0 = -\gamma + \delta.$$

From the first equation, we have $\alpha = 0$. Substituting this into the second equation gives us $\beta = 0$. Substituting this into the third equation gives us $\gamma = 0$. Finally, substituting this into the fourth equation gives us $\delta = 0$. Thus, the only solution to the original equation is $\alpha = \beta = \gamma = \delta = 0$, and so the list is linearly independent. ■

Exercise 2A9

Prove or give a counterexample: If $v_1, v_2, \dots, v_m$ is a linearly independent list of vectors in V , then $5v_1 - 4v_2, v_2, v_3, \dots, v_m$ is linearly independent.

Solution. Let $\alpha_1, \dots, \alpha_m \in \mathbb{F}$ such that

$$\alpha_1(5v_1 - 4v_2) + \alpha_2v_2 + \dots + \alpha_mv_m = 0.$$

Then we have

$$5\alpha_1v_1 + (-4\alpha_1 + \alpha_2)v_2 + \alpha_3v_3 + \dots + \alpha_mv_m = 0.$$

Since $v_1, v_2, \dots, v_m$ is linearly independent, we must have

$$\begin{aligned} 0 &= 5\alpha_1, \\ 0 &= -4\alpha_1 + \alpha_2, \\ 0 &= \alpha_3, \\ &\vdots \\ 0 &= \alpha_m. \end{aligned}$$

From the first equation, we have $\alpha_1 = 0$. Substituting this into the second equation gives us $\alpha_2 = 0$. The remaining equations give us $\alpha_k = 0$ for $k = 3, \dots, m$. Thus, the only solution to the original equation is $\alpha_k = 0$ for $k = 1, \dots, m$, and so the list is linearly independent. ■

Exercise 2A10

Prove or give a counterexample: If $v_1, v_2, \dots, v_m$ is a linearly independent list of vectors in V and $\lambda \in \mathbb{F}$ with $\lambda \neq 0$, then $\lambda v_1, \lambda v_2, \dots, \lambda v_m$ is linearly independent.

Solution. Let $\alpha_1, \dots, \alpha_m \in \mathbb{F}$ such that

$$\alpha_1(\lambda v_1) + \dots + \alpha_m(\lambda v_m) = 0.$$

Then we have

$$\lambda(\alpha_1v_1 + \dots + \alpha_mv_m) = 0.$$

Since $\lambda \neq 0$, we must have

$$\alpha_1v_1 + \dots + \alpha_mv_m = 0.$$

Since $v_1, v_2, \dots, v_m$ is linearly independent, we must have $\alpha_k = 0$ for $k = 1, \dots, m$. Thus, the only solution to the original equation is $\alpha_k = 0$ for $k = 1, \dots, m$, and so the list is linearly independent. ■

Exercise 2A11

Prove or give a counterexample: If $v_1, \dots, v_m$ and $w_1, \dots, w_m$ are linearly independent lists of vectors in V , then the list $v_1 + w_1, \dots, v_m + w_m$ is linearly independent.

Solution. Consider the following counterexample in $\mathbb{R}^2$:

$$v_1 = (1, 0), \quad v_2 = (0, 1), \quad w_1 = (0, 1), \quad w_2 = (1, 0).$$

Then v_1, v_2 and w_1, w_2 are linearly independent lists of vectors in $\mathbb{R}^2$. However,

$$v_1 + w_1 = (1, 1), \quad v_2 + w_2 = (1, 1),$$

and so the list $v_1 + w_1, v_2 + w_2$ is linearly dependent. Thus, the statement is false. ■

Exercise 2A12

Suppose $v_1, \dots, v_m$ is linearly independent in V and $w \in V$. Prove that if $v_1 + w, \dots, v_m + w$ is linearly dependent, then $w \in \text{span}(v_1, \dots, v_m)$.

Solution. Let $\alpha_1, \dots, \alpha_m \in \mathbb{F}$, not all zero, such that

$$\alpha_1(v_1 + w) + \dots + \alpha_m(v_m + w) = 0.$$

Then we have

$$(\alpha_1 v_1 + \dots + \alpha_m v_m) + (\alpha_1 + \dots + \alpha_m)w = 0.$$

If $\alpha_1 + \dots + \alpha_m = 0$, then we have

$$\alpha_1 v_1 + \dots + \alpha_m v_m = 0,$$

which contradicts the linear independence of $v_1, \dots, v_m$. Thus, we must have $\alpha_1 + \dots + \alpha_m \neq 0$. Let $\beta = \alpha_1 + \dots + \alpha_m$. Then we may write

$$w = -\frac{\alpha_1}{\beta}v_1 - \dots - \frac{\alpha_m}{\beta}v_m.$$

Thus, w is a linear combination of $v_1, \dots, v_m$, and so $w \in \text{span}(v_1, \dots, v_m)$. ■

Exercise 2A13

Suppose $v_1, \dots, v_m$ is linearly independent in V and $w \in V$. Show that

$$v_1, \dots, v_m, w \text{ is linearly independent} \iff w \notin \text{span}(v_1, \dots, v_m).$$

Solution. Suppose $v_1, \dots, v_m, w$ is linearly independent. If there were scalars $\alpha_1, \dots, \alpha_m, \beta$ such that

$$\alpha_1 v_1 + \dots + \alpha_m v_m + \beta w = 0,$$

then linear independence would imply that $\alpha_1 = \dots = \alpha_m = \beta = 0$. Suppose $w \in \text{span}(v_1, \dots, v_m)$. Then there exist scalars $\gamma_1, \dots, \gamma_m$ not all zero such that

$$\gamma_1 v_1 + \dots + \gamma_m v_m = w.$$

Then we may write

$$-\gamma_1 v_1 - \cdots - \gamma_m v_m + 1 \cdot w = 0,$$

which contradicts the linear independence of $v_1, \dots, v_m, w$. Thus, $w \notin \text{span}(v_1, \dots, v_m)$.

Conversely, suppose $w \notin \text{span}(v_1, \dots, v_m)$. Then there do not exist scalars $\alpha_1, \dots, \alpha_m$ such that

$$\alpha_1 v_1 + \cdots + \alpha_m v_m = w.$$

If $v_1, \dots, v_m, w$ is linearly dependent, then there exist scalars $\beta_1, \dots, \beta_m, \gamma$, not all zero, such that

$$\beta_1 v_1 + \cdots + \beta_m v_m + \gamma w = 0.$$

If $\gamma = 0$, then we have

$$\beta_1 v_1 + \cdots + \beta_m v_m = 0,$$

which contradicts the linear independence of $v_1, \dots, v_m$. Thus, we must have $\gamma \neq 0$. Then we may write

$$w = -\frac{\beta_1}{\gamma} v_1 - \cdots - \frac{\beta_m}{\gamma} v_m,$$

which contradicts the assumption that w is not in the span of $v_1, \dots, v_m$. Thus, $v_1, \dots, v_m, w$ is linearly independent. ■

Exercise 2A14

Suppose $v_1, \dots, v_m$ is a list of vectors in V . For $k \in \{1, \dots, m\}$, let

$$w_k = v_1 + \cdots + v_k.$$

Show that the list $v_1, \dots, v_m$ is linearly independent if and only if the list $w_1, \dots, w_m$ is linearly independent.

Solution. Suppose $v_1, \dots, v_m$ is linearly independent. Let $\alpha_1, \dots, \alpha_m \in \mathbb{F}$ such that

$$\alpha_1 w_1 + \cdots + \alpha_m w_m = 0.$$

Noting that $v_k = w_k - w_{k-1}$ for $k \in \{2, \dots, m\}$ and $v_1 = w_1$, we may rewrite the equation as

$$\alpha_1 v_1 + (\alpha_2 - \alpha_1) v_2 + \cdots + (\alpha_m - \alpha_{m-1}) v_m = 0.$$

Since $v_1, \dots, v_m$ is linearly independent, we must have $\alpha_1 = 0$. Then this implies that $\alpha_2 - \alpha_1 = 0$, and so on, until we have $\alpha_m - \alpha_{m-1} = 0$. Thus, the only solution to the original equation is $\alpha_1 = \cdots = \alpha_m = 0$, and so $w_1, \dots, w_m$ is linearly independent.

Conversely, suppose $w_1, \dots, w_m$ is linearly independent. Let $\beta_1, \dots, \beta_m \in \mathbb{F}$ such that

$$\beta_1 v_1 + \cdots + \beta_m v_m = 0.$$

Then we have

$$\beta_1 w_1 + (\beta_2 + \beta_1) w_2 + \cdots + (\beta_m + \cdots + \beta_1) w_m = 0.$$

Since $w_1, \dots, w_m$ is linearly independent, we must have $\beta_1 = 0$. Then this implies that $\beta_2 + \beta_1 = 0$, and so on, until we have $\beta_m + \cdots + \beta_1 = 0$. Thus, the only solution to the original equation is $\beta_1 = \cdots = \beta_m = 0$, and so $v_1, \dots, v_m$ is linearly independent. ■

Exercise 2A15

Explain why there does not exist a list of six polynomials that is linearly independent in $\mathcal{P}_4(\mathbb{F})$.

Solution. Since $\mathcal{P}_4(\mathbb{F})$ can be spanned by the list $1, x, x^2, x^3, x^4$, and the length of a linearly independent list cannot exceed the length of a spanning list, there does not exist a list of six polynomials that is linearly independent in $\mathcal{P}_4(\mathbb{F})$. ■

Exercise 2A16

Explain why no list of four polynomials spans $\mathcal{P}_4(\mathbb{F})$.

Solution. Since $\mathcal{P}_4(\mathbb{F})$ can be spanned by the list $1, x, x^2, x^3, x^4$, and the length of a spanning list cannot be less than the length of a linearly independent list, no list of four polynomials spans $\mathcal{P}_4(\mathbb{F})$. ■

Exercise 2A17

Prove that V is infinite-dimensional if and only if there is a sequence $v_1, v_2, \dots$ of vectors in V such that $v_1, \dots, v_m$ is linearly independent for every positive integer m .

Solution. Suppose V is infinite-dimensional. We proceed by induction to construct a sequence of vectors $v_1, v_2, \dots$ in V such that $v_1, \dots, v_m$ is linearly independent for every positive integer m . Since V is infinite-dimensional, there exists a nonzero vector $v_1 \in V$. Suppose we have constructed a linearly independent list of vectors $v_1, \dots, v_m$ in V . Since V is infinite-dimensional, there exists a vector $v_{m+1} \in V$ such that $v_{m+1} \notin \text{span}(v_1, \dots, v_m)$. Then $v_1, \dots, v_m, v_{m+1}$ is linearly independent. By induction, we have constructed a sequence of vectors $v_1, v_2, \dots$ in V such that $v_1, \dots, v_m$ is linearly independent for every positive integer m .

Conversely, suppose there exists a sequence of vectors $v_1, v_2, \dots$ in V such that $v_1, \dots, v_m$ is linearly independent for every positive integer m . Then for every positive integer m , there exists a linearly independent list of vectors in V of length m . Since there is no maximum length for a linearly independent list of vectors in V , this implies that V is infinite-dimensional. ■

Exercise 2A18

Prove that $\mathbb{F}^\infty$ is infinite-dimensional.

Solution. By [Exercise 2A17](#), it suffices to construct a sequence of vectors $v_1, v_2, \dots$ in $\mathbb{F}^\infty$ such that $v_1, \dots, v_m$ is linearly independent for every positive integer m . Let v_k be the vector in $\mathbb{F}^\infty$ whose k -th coordinate is 1 and all other coordinates are 0. Explicitly, we have

$$v_1 = (1, 0, 0, \dots), \quad v_2 = (0, 1, 0, \dots), \quad v_3 = (0, 0, 1, \dots), \quad \dots$$

Then for any positive integer m , the list $v_1, \dots, v_m$ is linearly independent, since the only way to write the zero vector as a linear combination of $v_1, \dots, v_m$ is to take all coefficients to be zero. Thus, by [Exercise 2A17](#), $\mathbb{F}^\infty$ is infinite-dimensional. ■

Exercise 2A19

Prove that the real vector space of all continuous real-valued functions on the interval $[0, 1]$ is infinite-dimensional.

Solution. By [Exercise 2A17](#), it suffices to construct a sequence of continuous real-valued functions $f_1, f_2, \dots$ on the interval $[0, 1]$ such that $f_1, \dots, f_m$ is linearly independent for every positive integer m . Let $f_k(x) = x^{k-1}$ for $k = 1, 2, \dots$. Then for any positive integer m , the list $f_1, \dots, f_m$ is linearly independent, since the only way to write the zero function as a linear combination of $f_1, \dots, f_m$ is to take all coefficients to be zero. Thus, by [Exercise 2A17](#), $\mathbb{R}^{[0,1]}$ is infinite-dimensional. ■

Exercise 2A20

Suppose $p_0, p_1, \dots, p_m$ are polynomials in $\mathcal{P}_m(\mathbb{F})$ such that $p_k(2) = 0$ for each $k \in \{0, \dots, m\}$. Prove that $p_0, p_1, \dots, p_m$ is not linearly independent in $\mathcal{P}_m(\mathbb{F})$.

Solution. Since $p_k(2) = 0$ for each $k \in \{0, \dots, m\}$, we have

$$p_k(x) = (x - 2)q_k(x)$$

for some polynomial $q_k(x) \in \mathcal{P}_{m-1}(\mathbb{F})$. Then we may write

$$p_0(x), p_1(x), \dots, p_m(x) = (x - 2)q_0(x), (x - 2)q_1(x), \dots, (x - 2)q_m(x).$$

Since $\mathcal{P}_{m-1}(\mathbb{F})$ can be spanned by the m polynomials $1, x, \dots, x^{m-1}$, the list of $m + 1$ polynomials $q_0, q_1, \dots, q_m$ must be linearly dependent. Thus, there exist scalars $\alpha_0, \dots, \alpha_m$, not all zero, such that

$$\alpha_0 q_0(x) + \dots + \alpha_m q_m(x) = 0.$$

Then we may write

$$\alpha_0 p_0(x) + \dots + \alpha_m p_m(x) = (x - 2)(\alpha_0 q_0(x) + \dots + \alpha_m q_m(x)) = 0.$$

Since not all the scalars $\alpha_0, \dots, \alpha_m$ are zero, this shows that $p_0, p_1, \dots, p_m$ is linearly dependent in $\mathcal{P}_m(\mathbb{F})$. ■

2B Bases

Exercise 2B1

Find all vector spaces that have exactly one basis.

Solution. Let V be a vector space. If V has exactly one basis, then V must be finite-dimensional, since infinite-dimensional vector spaces have infinitely many bases. Let $v_1, \dots, v_n$ be a basis of V . Then any other basis of V must also have length n . Since there is only one basis of V , it must be that $n = 0$, and so the only vector space that has exactly one basis is the zero vector space. ■

Exercise 2B2

Verify all assertions in Example 2.27.

- (a) The list $(1, 0, \dots, 0), (0, 1, 0, \dots, 0), \dots, (0, \dots, 0, 1)$ is a basis of $\mathbb{F}^n$, called the *standard basis* of $\mathbb{F}^n$.
- (b) The list $(1, 2), (3, 5)$ is a basis of $\mathbb{F}^2$. Note that this list has length two, which is the same as the length of the standard basis of $\mathbb{F}^2$. In the next section, we will see that this is not a coincidence.
- (c) The list $(1, 2, -4), (7, -5, 6)$ is linearly independent in $\mathbb{F}^3$ but is not a basis of $\mathbb{F}^3$ because it does not span $\mathbb{F}^3$.
- (d) The list $(1, 2), (3, 5), (4, 13)$ spans $\mathbb{F}^2$ but is not a basis of $\mathbb{F}^2$ because it is not linearly independent.
- (e) The list $(1, 1, 0), (0, 0, 1)$ is a basis of $\{(x, x, y) \in \mathbb{F}^3 \mid x, y \in \mathbb{F}\}$.
- (f) The list $(1, -1, 0), (1, 0, -1)$ is a basis of

$$\{(x, y, z) \in \mathbb{F}^3 \mid x + y + z = 0\}.$$
- (g) The list $1, z, \dots, z^m$ is a basis of $\mathcal{P}_m(\mathbb{F})$, called the *standard basis* of $\mathcal{P}_m(\mathbb{F})$.

Solution.

- (a) Suppose $\alpha_1, \dots, \alpha_n \in \mathbb{F}$ such that

$$\alpha_1(1, 0, \dots, 0) + \alpha_2(0, 1, 0, \dots, 0) + \dots + \alpha_n(0, \dots, 0, 1) = 0.$$

Then each $\alpha_i = 0$, so the list is linearly independent. Moreover, any vector $(x_1, \dots, x_n) \in \mathbb{F}^n$ can be written as

$$(x_1, \dots, x_n) = x_1(1, 0, \dots, 0) + x_2(0, 1, 0, \dots, 0) + \dots + x_n(0, \dots, 0, 1),$$

so the list spans $\mathbb{F}^n$. Thus, the list is a basis of $\mathbb{F}^n$.

- (b) If $(3, 5) = a(1, 2)$ for some $a \in \mathbb{F}$, then we must have $3 = a$ and $5 = 2a$, which is impossible. Thus, the list is linearly independent. Moreover, for any element $(x, y) \in \mathbb{F}^2$, we can solve the system

of equations

$$\begin{aligned}x &= 3\alpha + \beta, \\y &= 5\alpha + 2\beta\end{aligned}$$

for $\alpha, \beta \in \mathbb{F}$. Solving this system of equations, we find that

$$\alpha = 2x - y, \quad \beta = 5x + 3y.$$

Since we can find $\alpha, \beta \in \mathbb{F}$ for any $(x, y) \in \mathbb{F}^2$, the list spans $\mathbb{F}^2$. Thus, the list is a basis of $\mathbb{F}^2$.

- (c) If $(7, -5, 6) = a(1, 2, -4)$ for some $a \in \mathbb{F}$, then we must have $7 = a$, $-5 = 2a$, and $6 = -4a$, which is impossible. Thus, the list is linearly independent. Moreover, the list does not span $\mathbb{F}^3$. If it did, then we could write $(0, 0, 1) = \alpha(1, 2, -4) + \beta(7, -5, 6)$ for some $\alpha, \beta \in \mathbb{F}$. This would give us the system of equations

$$\begin{aligned}0 &= \alpha + 7\beta, \\0 &= 2\alpha - 5\beta, \\1 &= -4\alpha + 6\beta.\end{aligned}$$

From the first equation, we see that $\alpha = -7\beta$. Substituting this into the second equation gives us $0 = -14\beta - 5\beta = -19\beta$, so $\beta = 0$ and $\alpha = 0$. However, substituting these values into the third equation gives us $1 = 0$, which is impossible. Thus, the list does not span $\mathbb{F}^3$, and so it is not a basis of $\mathbb{F}^3$.

- (d) From part (b), we saw that $(1, 2), (3, 5)$ is a basis of $\mathbb{F}^2$. Since $(4, 13) = -5(3, 5) + 19(1, 2)$, the list $(1, 2), (3, 5), (4, 13)$ is linearly dependent. Moreover, since $(1, 2), (3, 5)$ is a basis of $\mathbb{F}^2$, the list $(1, 2), (3, 5), (4, 13)$ spans $\mathbb{F}^2$. Thus, the list is not a basis of $\mathbb{F}^2$.

- (e) Let U be the subspace in question. If $\alpha, \beta \in \mathbb{F}$ such that

$$\alpha(1, 1, 0) + \beta(0, 0, 1) = (0, 0, 0),$$

then we must have $\alpha = \beta = 0$, so the list is linearly independent. Moreover, for any $(x, x, y) \in U$, we can write

$$(x, x, y) = x(1, 1, 0) + y(0, 0, 1),$$

so the list spans U . Thus, the list is a basis of U .

- (f) Let U be the subspace in question. If $\alpha, \beta \in \mathbb{F}$ such that

$$\alpha(1, -1, 0) + \beta(1, 0, -1) = (0, 0, 0),$$

then we must have $\alpha = \beta = 0$, so the list is linearly independent. Moreover, for any $(x, y, z) \in U$, we can write

$$(x, y, z) = (x - z)(1, -1, 0) + (y - z)(1, 0, -1),$$

so the list spans U . Thus, the list is a basis of U .

- (g) If $\alpha_0, \dots, \alpha_m \in \mathbb{F}$ such that

$$\alpha_0 + \alpha_1 z + \dots + \alpha_m z^m = 0,$$

then we must have $\alpha_0 = \dots = \alpha_m = 0$, so the list is linearly independent. Moreover, any polynomial $p(z) \in \mathcal{P}_m(\mathbb{F})$ can be written as

$$p(z) = \alpha_0 + \alpha_1 z + \dots + \alpha_m z^m,$$

so the list spans $\mathcal{P}_m(\mathbb{F})$. Thus, the list is a basis of $\mathcal{P}_m(\mathbb{F})$. ■

Exercise 2B3

- (a) Let
- U
- be the subspace of
- $\mathbb{R}^5$
- defined by

$$U = \{(x_1, x_2, x_3, x_4, x_5) \in \mathbb{R}^5 \mid x_1 = 3x_2 \text{ and } x_3 = 7x_4\}.$$

Find a basis of U .

- (b) Extend the basis in (a) to a basis of $\mathbb{R}^5$.
- (c) Find a subspace W of $\mathbb{R}^5$ such that $\mathbb{R}^5 = U \oplus W$.

Solution.

- (a) Note that we can rewrite the subspace
- U
- as

$$U = \{(3x_2, x_2, 7x_4, x_4, x_5) \in \mathbb{R}^5 \mid x_2, x_4, x_5 \in \mathbb{R}\}.$$

Then we can write any vector in U as a linear combination of the following three vectors:

$$u_1 = (3, 1, 0, 0, 0), \quad u_2 = (0, 0, 7, 1, 0), \quad u_3 = (0, 0, 0, 0, 1).$$

Moreover, these three vectors are linearly independent, since the only way to write the zero vector as a linear combination of u_1, u_2, u_3 is to take all coefficients to be zero. Thus, $\{u_1, u_2, u_3\}$ is a basis of U .

- (b) We can extend the basis
- $\{u_1, u_2, u_3\}$
- of
- U
- to a basis of
- $\mathbb{R}^5$
- by adding two more vectors that are linearly independent of
- u_1, u_2, u_3
- . For example, we can add the vectors

$$u_4 = (1, 0, 0, 0, 0), \quad u_5 = (0, 0, 1, 0, 0),$$

which are linearly independent of u_1, u_2, u_3 since they are not in U . Thus, $\{u_1, u_2, u_3, u_4, u_5\}$ is a basis of $\mathbb{R}^5$.

- (c) Let
- W
- be the subspace of
- $\mathbb{R}^5$
- spanned by the vectors
- u_4
- and
- u_5
- . Then we have

$$\mathbb{R}^5 = U \oplus W,$$

since $U \cap W = \{0\}$ and every vector in $\mathbb{R}^5$ can be written as a sum of a vector in U and a vector in W . ■

Exercise 2B4

- (a) Let
- U
- be the subspace of
- $\mathbb{C}^5$
- defined by

$$U = \{(z_1, z_2, z_3, z_4, z_5) \in \mathbb{C}^5 \mid 6z_1 = z_2 \text{ and } z_3 + 2z_4 + 3z_5 = 0\}.$$

Find a basis of U .

- (b) Extend the basis in (a) to a basis of $\mathbb{C}^5$.
- (c) Find a subspace W of $\mathbb{C}^5$ such that $\mathbb{C}^5 = U \oplus W$.

Solution.

- (a) Note that we can rewrite the subspace
- U
- as

$$U = \{(z_1, 6z_1, -2z_4 - 3z_5, z_4, z_5) \in \mathbb{C}^5 \mid z_1, z_4, z_5 \in \mathbb{C}\}.$$

Then we can write any vector in U as a linear combination of the following three vectors:

$$u_1 = (1, 6, 0, 0, 0), \quad u_2 = (0, 0, -2, 1, 0), \quad u_3 = (0, 0, -3, 0, 1).$$

Moreover, these three vectors are linearly independent, since the only way to write the zero vector as a linear combination of u_1, u_2, u_3 is to take all coefficients to be zero. Thus, $\{u_1, u_2, u_3\}$ is a basis of U .

- (b) We can extend the basis
- $\{u_1, u_2, u_3\}$
- of
- U
- to a basis of
- $\mathbb{C}^5$
- by adding two more vectors that are linearly independent of
- u_1, u_2, u_3
- . For example, we can add the vectors

$$u_4 = (1, 0, 0, 0, 0), \quad u_5 = (0, 0, 1, 0, 0),$$

which are linearly independent of u_1, u_2, u_3 since they are not in U . Thus, $\{u_1, u_2, u_3, u_4, u_5\}$ is a basis of $\mathbb{C}^5$.

- (c) Let
- W
- be the subspace of
- $\mathbb{C}^5$
- spanned by the vectors
- u_4
- and
- u_5
- . Then we have

$$\mathbb{C}^5 = U \oplus W,$$

since $U \cap W = \{0\}$ and every vector in $\mathbb{C}^5$ can be written as a sum of a vector in U and a vector in W . ■

Exercise 2B5

Suppose V is finite-dimensional and U, W are subspaces of V such that $V = U + W$. Prove that there exists a basis of V consisting of vectors in $U \cup W$.

Solution. Let $u_1, \dots, u_m$ be a basis of U and $w_1, \dots, w_n$ be a basis of W . Then the list $u_1, \dots, u_m, w_1, \dots, w_n$ spans V , since any vector in V can be written as a sum of a vector in U and a vector in W . We can extract a basis of V from this list by removing any linearly dependent vectors. Thus, there exists a basis of V consisting of vectors in $U \cup W$. ■

Exercise 2B6

Prove or give a counterexample: If p_0, p_1, p_2, p_3 is a list in $\mathcal{P}_3(\mathbb{F})$ such that none of the polynomials p_0, p_1, p_2, p_3 has degree 2, then the list is not a basis of $\mathcal{P}_3(\mathbb{F})$.

Solution. This is not true. Let $p_0 = 1, p_1 = x, p_2 = x^3, p_3 = x^2 + x^3$. Then none of the polynomials p_0, p_1, p_2, p_3 has degree 2. Moreover, the list p_0, p_1, p_2, p_3 is linearly independent, since the only way to write the zero polynomial as a linear combination of p_0, p_1, p_2, p_3 is to take all coefficients to be zero. Finally, noting that $x^2 = p_3 - p_2$, then for any polynomial $p(x) = a_0 + a_1x + a_2x^2 + a_3x^3 \in \mathcal{P}_3(\mathbb{F})$, we can write

$$p(x) = a_0p_0 + a_1p_1 + a_2(p_3 - p_2) + a_3p_2 = a_0p_0 + a_1p_1 + (a_3 - a_2)p_2 + a_2p_3,$$

which shows that the list spans $\mathcal{P}_3(\mathbb{F})$. Thus, the list p_0, p_1, p_2, p_3 is a basis of $\mathcal{P}_3(\mathbb{F})$. ■

Exercise 2B7

Suppose v_1, v_2, v_3, v_4 is a basis of V . Prove that

$$v_1 + v_2, v_2 + v_3, v_3 + v_4, v_4$$

is also a basis of V .

Solution. Let $\alpha_1, \alpha_2, \alpha_3, \alpha_4 \in \mathbb{F}$ such that

$$\alpha_1(v_1 + v_2) + \alpha_2(v_2 + v_3) + \alpha_3(v_3 + v_4) + \alpha_4 v_4 = 0.$$

Then we may rewrite this equation as

$$\alpha_1 v_1 + (\alpha_1 + \alpha_2)v_2 + (\alpha_2 + \alpha_3)v_3 + (\alpha_3 + \alpha_4)v_4 = 0.$$

Since v_1, v_2, v_3, v_4 is a basis of V , it is linearly independent, and so we must have

$$\alpha_1 = 0, \quad \alpha_1 + \alpha_2 = 0, \quad \alpha_2 + \alpha_3 = 0, \quad \alpha_3 + \alpha_4 = 0.$$

From the first equation, we have $\alpha_1 = 0$. Then the second equation gives us $\alpha_2 = 0$. The third equation then gives us $\alpha_3 = 0$, and finally the fourth equation gives us $\alpha_4 = 0$. Thus, the list is linearly independent. Moreover, since v_1, v_2, v_3, v_4 is a basis of V , it spans V , and so the list also spans V . Therefore, the list is a basis of V . ■

Exercise 2B8

Prove or give a counterexample: If v_1, v_2, v_3, v_4 is a basis of V and U is a subspace of V such that $v_1, v_2 \in U$ and $v_3 \notin U$ and $v_4 \notin U$, then v_1, v_2 is a basis of U .

Solution. This is not true. Let $V = \mathbb{R}^4$ and v_1, v_2, v_3, v_4 be the standard basis of $\mathbb{R}^4$. Let U be the subspace of $\mathbb{R}^4$ defined by

$$U = \text{span}(v_1, v_2, v_3 + v_4) = \{(x, y, z, z) \in \mathbb{R}^4 \mid x, y, z \in \mathbb{R}\}.$$

Then $v_1, v_2 \in U$ and $v_3, v_4 \notin U$. However, v_1, v_2 does not span U , since $(0, 0, 1, 1) \in U$ but cannot be written as a linear combination of v_1 and v_2 . Thus, v_1, v_2 is not a basis of U . ■

Exercise 2B9

Suppose $v_1, \dots, v_m$ is a list of vectors in V . For $k \in \{1, \dots, m\}$, let

$$w_k = v_1 + \dots + v_k.$$

Show that $v_1, \dots, v_m$ is a basis of V if and only if $w_1, \dots, w_m$ is a basis of V .

Solution. By [Exercise 2A14](#), we know that $v_1, \dots, v_m$ is linearly independent if and only if $w_1, \dots, w_m$ is linearly independent. By [Exercise 2A3](#), we know that $v_1, \dots, v_m$ spans V if and only if $w_1, \dots, w_m$ spans V . Thus, $v_1, \dots, v_m$ is a basis of V if and only if $w_1, \dots, w_m$ is a basis of V . ■

Exercise 2B10

Suppose U and W are subspaces of V such that $V = U \oplus W$. Suppose also that $u_1, \dots, u_m$ is a basis of U and $w_1, \dots, w_n$ is a basis of W . Prove that

$$u_1, \dots, u_m, w_1, \dots, w_n$$

is a basis of V .

Solution. Let $u_1, \dots, u_m$ be a basis of U and $w_1, \dots, w_n$ be a basis of W . We claim that the list

$$u_1, \dots, u_m, w_1, \dots, w_n$$

is a basis of V . Let $\alpha_1, \dots, \alpha_m, \beta_1, \dots, \beta_n \in \mathbb{F}$ such that

$$\alpha_1 u_1 + \dots + \alpha_m u_m + \beta_1 w_1 + \dots + \beta_n w_n = 0.$$

Then we may write

$$\alpha_1 u_1 + \dots + \alpha_m u_m = -\beta_1 w_1 - \dots - \beta_n w_n.$$

Since the left-hand side is in U and the right-hand side is in W , and $U \cap W = \{0\}$, we must have

$$\alpha_1 u_1 + \dots + \alpha_m u_m = 0 \quad \text{and} \quad \beta_1 w_1 + \dots + \beta_n w_n = 0.$$

Since $u_1, \dots, u_m$ is a basis of U , we have $\alpha_1 = \dots = \alpha_m = 0$. Similarly, since $w_1, \dots, w_n$ is a basis of W , we have $\beta_1 = \dots = \beta_n = 0$. Thus, the list is linearly independent. Moreover, since $V = U \oplus W$, then $v = u + w$ for some $u \in U$ and $w \in W$. Then we may write

$$u = \gamma_1 u_1 + \dots + \gamma_m u_m \quad \text{and} \quad w = \delta_1 w_1 + \dots + \delta_n w_n$$

for some $\gamma_1, \dots, \gamma_m, \delta_1, \dots, \delta_n \in \mathbb{F}$. Hence,

$$v = u + w = \gamma_1 u_1 + \dots + \gamma_m u_m + \delta_1 w_1 + \dots + \delta_n w_n,$$

which shows that the list spans V . Therefore, the list is a basis of V . ■

Exercise 2B11

Suppose V is a real vector space. Show that if $v_1, \dots, v_n$ is a basis of V (as a real vector space), then $v_1, \dots, v_n$ is also a basis of the complexification $V_{\mathbb{C}}$ (as a complex vector space).

Note. Please see [Exercise 1B8](#) for the definition of the complexification $V_{\mathbb{C}}$.

Solution. Let $v_1, \dots, v_n$ be a basis of V . We claim that $v_1, \dots, v_n$ is also a basis of $V_{\mathbb{C}}$. Let $\alpha_1, \dots, \alpha_n \in \mathbb{C}$ such that

$$\alpha_1 v_1 + \dots + \alpha_n v_n = 0.$$

Then we may write each $\alpha_k = a_k + b_k i$ for some $a_k, b_k \in \mathbb{R}$. Then we have

$$(a_1 + b_1 i)v_1 + \dots + (a_n + b_n i)v_n = 0.$$

This implies that

$$(a_1 v_1 + \dots + a_n v_n) + i(b_1 v_1 + \dots + b_n v_n) = 0.$$

Since the left-hand side is a sum of two vectors in V , and V is a real vector space, we must have

$$a_1v_1 + \cdots + a_nv_n = 0 \quad \text{and} \quad b_1v_1 + \cdots + b_nv_n = 0.$$

Since $v_1, \dots, v_n$ is a basis of V , we have $a_1 = \cdots = a_n = 0$ and $b_1 = \cdots = b_n = 0$. Thus, $\alpha_1 = \cdots = \alpha_n = 0$, which shows that the list is linearly independent. Moreover, let $v \in V_{\mathbb{C}}$. Then we may write $v = u + wi$ for some $u, w \in V$. Since $v_1, \dots, v_n$ is a basis of V , we may write

$$u = \gamma_1v_1 + \cdots + \gamma_nv_n \quad \text{and} \quad w = \delta_1v_1 + \cdots + \delta_nv_n$$

for some $\gamma_1, \dots, \gamma_n, \delta_1, \dots, \delta_n \in \mathbb{R}$. Hence,

$$v = u + wi = (\gamma_1 + \delta_1i)v_1 + \cdots + (\gamma_n + \delta_ni)v_n,$$

which shows that the list spans $V_{\mathbb{C}}$. Therefore, the list is a basis of $V_{\mathbb{C}}$. ■

2C Dimension

Exercise 2C1

Show that the subspaces of $\mathbb{R}^2$ are precisely $\{0\}$, all lines in $\mathbb{R}^2$ containing the origin, and $\mathbb{R}^2$.

Solution. Since $\dim(\mathbb{R}^2) = 2$, any subspace of $\mathbb{R}^2$ must have dimension 0, 1, or 2. The only subspace of dimension 0 is $\{0\}$, and the only subspace of dimension 2 is $\mathbb{R}^2$. Any subspace with dimension 1 consists of all scalar multiples of some nonzero $(x, y) \in \mathbb{R}^2$, which is precisely the line with direction (x, y) containing the origin. Thus, the subspaces of $\mathbb{R}^2$ are precisely $\{0\}$, all lines in $\mathbb{R}^2$ containing the origin, and $\mathbb{R}^2$. ■

Exercise 2C2

Show that the subspaces of $\mathbb{R}^3$ are precisely $\{0\}$, all lines in $\mathbb{R}^3$ containing the origin, all planes in $\mathbb{R}^3$ containing the origin, and $\mathbb{R}^3$.

Solution. Since $\dim(\mathbb{R}^3) = 3$, any subspace of $\mathbb{R}^3$ must have dimension 0, 1, 2, or 3. The only subspace of dimension 0 is $\{0\}$, and the only subspace of dimension 3 is $\mathbb{R}^3$. Any subspace with dimension 1 consists of all scalar multiples of some nonzero $(x, y, z) \in \mathbb{R}^3$, which is precisely the line containing the origin in the direction of (x, y, z) . Any subspace with dimension 2 consists of all linear combinations of two linearly independent vectors in $\mathbb{R}^3$, which is precisely a plane containing the origin. Thus, the subspaces of $\mathbb{R}^3$ are precisely $\{0\}$, all lines in $\mathbb{R}^3$ containing the origin, all planes in $\mathbb{R}^3$ containing the origin, and $\mathbb{R}^3$. ■

Exercise 2C3

- (a) Let $U = \{p \in \mathcal{P}_4(\mathbb{F}) \mid p(6) = 0\}$. Find a basis of U .
- (b) Extend the basis in (a) to a basis of $\mathcal{P}_4(\mathbb{F})$.
- (c) Find a subspace W of $\mathcal{P}_4(\mathbb{F})$ such that $\mathcal{P}_4(\mathbb{F}) = U \oplus W$.

Solution.

- (a) Let $p = a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 \in U$. Then we have $p(6) = 0$, which gives us the equation

$$a_0 + 6a_1 + 36a_2 + 216a_3 + 1296a_4 = 0.$$

We can solve this equation for a_0 in terms of a_1, a_2, a_3, a_4 :

$$a_0 = -6a_1 - 36a_2 - 216a_3 - 1296a_4.$$

Then we can write any polynomial in U as a linear combination of the following four polynomials:

$$u_1 = -6 + x, \quad u_2 = -36 + x^2, \quad u_3 = -216 + x^3, \quad u_4 = -1296 + x^4.$$

Moreover, these four polynomials are linearly independent, since the only way to write the zero polynomial as a linear combination of u_1, u_2, u_3, u_4 is to take all coefficients to be zero. Thus, $\{u_1, u_2, u_3, u_4\}$ is a basis of U .

- (b) We can extend the basis $\{u_1, u_2, u_3, u_4\}$ of U to a basis of $\mathcal{P}_4(\mathbb{F})$ by adding one more polynomial that is linearly independent of u_1, u_2, u_3, u_4 . For example, we can add the polynomial

$$u_5 = 1,$$

which is linearly independent of u_1, u_2, u_3, u_4 since it is not in U . Thus, $\{u_1, u_2, u_3, u_4, u_5\}$ is a basis of $\mathcal{P}_4(\mathbb{F})$.

- (c) Let $W = \text{span } u_5 = \text{span } 1$. Then we have

$$\mathcal{P}_4(\mathbb{F}) = U + W,$$

since every polynomial in $\mathcal{P}_4(\mathbb{F})$ can be uniquely written as a sum of a polynomial in U and a polynomial in W . Moreover, if $p \in U \cap W$, then $p = \alpha$ for some $\alpha \in \mathbb{F}$. Since $p(6) = 0$, we have $\alpha = 0$. Thus, $U \cap W = \{0\}$, and so we have

$$\mathcal{P}_4(\mathbb{F}) = U \oplus W. \quad \blacksquare$$

Exercise 2C4

- (a) Let $U = \{p \in \mathcal{P}_4(\mathbb{R}) \mid p''(6) = 0\}$. Find a basis of U .
 (b) Extend the basis in (a) to a basis of $\mathcal{P}_4(\mathbb{R})$.
 (c) Find a subspace W of $\mathcal{P}_4(\mathbb{R})$ such that $\mathcal{P}_4(\mathbb{R}) = U \oplus W$.

Solution.

- (a) Let $p = a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 \in U$. Then we have

$$p''(x) = 2a_2 + 6a_3x + 12a_4x^2,$$

and so $p''(6) = 0$ gives us the equation

$$2a_2 + 36a_3 + 432a_4 = 0.$$

We can solve this equation for a_2 in terms of a_3, a_4 :

$$a_2 = -18a_3 - 216a_4.$$

Then we can write any polynomial in U as a linear combination of the following four polynomials:

$$u_1 = 1, \quad u_2 = x, \quad u_3 = -18x^2 + x^3, \quad u_4 = -216x^2 + x^4.$$

Moreover, these four polynomials are linearly independent, since the only way to write the zero polynomial as a linear combination of u_1, u_2, u_3, u_4 is to take all coefficients to be zero. Thus, $\{u_1, u_2, u_3, u_4\}$ is a basis of U .

- (b) We can extend the basis $\{u_1, u_2, u_3, u_4\}$ of U to a basis of $\mathcal{P}_4(\mathbb{R})$ by adding one more polynomial that is linearly independent of u_1, u_2, u_3, u_4 . For example, we can add the polynomial

$$u_5 = x^2,$$

which is linearly independent of u_1, u_2, u_3, u_4 since it is not in U . Thus, $\{u_1, u_2, u_3, u_4, u_5\}$ is a basis of $\mathcal{P}_4(\mathbb{R})$.

(c) Let $W = \text{span } u_5 = \text{span } x^2$. Then we have

$$\mathcal{P}_4(\mathbb{R}) = U + W,$$

since every polynomial in $\mathcal{P}_4(\mathbb{R})$ can be written as a sum of a polynomial in U and a polynomial in W . Moreover, if $p \in U \cap W$, then $p = \alpha x^2$ for some $\alpha \in \mathbb{R}$. Since $p''(6) = 0$, we have $2\alpha = 0$, which implies that $\alpha = 0$. Thus, $U \cap W = \{0\}$, and so we have

$$\mathcal{P}_4(\mathbb{R}) = U \oplus W. \quad \blacksquare$$

Exercise 2C5

- (a) Let $U = \{p \in \mathcal{P}_4(\mathbb{F}) \mid p(2) = p(5)\}$. Find a basis of U .
- (b) Extend the basis in (a) to a basis of $\mathcal{P}_4(\mathbb{F})$.
- (c) Find a subspace W of $\mathcal{P}_4(\mathbb{F})$ such that $\mathcal{P}_4(\mathbb{F}) = U \oplus W$.

Solution.

(a) Let $p = a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 \in U$. Then we have $p(2) = p(5)$, which gives us the equation

$$a_0 + 2a_1 + 4a_2 + 8a_3 + 16a_4 = a_0 + 5a_1 + 25a_2 + 125a_3 + 625a_4.$$

We can simplify this equation to get

$$-3a_1 - 21a_2 - 117a_3 - 609a_4 = 0.$$

We can solve this equation for a_1 in terms of a_2, a_3, a_4 :

$$a_1 = -7a_2 - 39a_3 - 203a_4.$$

Then we can write any polynomial in U as a linear combination of the following four polynomials:

$$u_1 = 1, \quad u_2 = -7x + x^2, \quad u_3 = -39x + x^3, \quad u_4 = -203x + x^4.$$

Moreover, these four polynomials are linearly independent, since the only way to write the zero polynomial as a linear combination of u_1, u_2, u_3, u_4 is to take all coefficients to be zero. Thus, $\{u_1, u_2, u_3, u_4\}$ is a basis of U .

(b) We can extend the basis $\{u_1, u_2, u_3, u_4\}$ of U to a basis of $\mathcal{P}_4(\mathbb{F})$ by adding one more polynomial that is linearly independent of u_1, u_2, u_3, u_4 . For example, we can add the polynomial

$$u_5 = x,$$

which is linearly independent of u_1, u_2, u_3, u_4 since it is not in U . Thus, $\{u_1, u_2, u_3, u_4, u_5\}$ is a basis of $\mathcal{P}_4(\mathbb{F})$.

(c) Let $W = \text{span } u_5 = \text{span } x$. Then we have

$$\mathcal{P}_4(\mathbb{F}) = U + W,$$

since every polynomial in $\mathcal{P}_4(\mathbb{F})$ can be written as a sum of a polynomial in U and a polynomial in W . Moreover, if $p \in U \cap W$, then $p = \alpha x$ for some $\alpha \in \mathbb{F}$. Since $p(2) = p(5)$, we have $\alpha 2 = \alpha 5$, which implies that $\alpha = 0$. Thus, $U \cap W = \{0\}$, and so we have

$$\mathcal{P}_4(\mathbb{F}) = U \oplus W. \quad \blacksquare$$

Exercise 2C6

- (a) Let $U = \{p \in \mathcal{P}_4(\mathbb{F}) \mid p(2) = p(5) = p(6)\}$. Find a basis of U .
- (b) Extend the basis in (a) to a basis of $\mathcal{P}_4(\mathbb{F})$.
- (c) Find a subspace W of $\mathcal{P}_4(\mathbb{F})$ such that $\mathcal{P}_4(\mathbb{F}) = U \oplus W$.

Solution.

- (a) Let $p = a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 \in U$. Then we have $p(2) = p(5) = p(6)$. We know from [Exercise 2C5](#) that the condition $p(2) = p(5)$ gives us the equation

$$a_1 = -7a_2 - 39a_3 - 203a_4.$$

The condition $p(5) = p(6)$ gives us the equation

$$5a_1 + 25a_2 + 125a_3 + 625a_4 = 6a_1 + 36a_2 + 216a_3 + 1296a_4,$$

which simplifies to

$$a_1 + 11a_2 + 91a_3 + 671a_4 = 0.$$

Since we know that $a_1 = -7a_2 - 39a_3 - 203a_4$, we can substitute this into the second equation to get

$$4a_2 + 52a_3 + 468a_4 = 0.$$

We can solve this equation for a_2 in terms of a_3, a_4 :

$$a_2 = -13a_3 - 117a_4.$$

Then $a_1 = 52a_3 + 616a_4$. Thus, we can write any polynomial in U as a linear combination of the following three polynomials:

$$u_1 = 1, \quad u_2 = 52x - 13x^2 + x^3, \quad u_3 = 616x - 117x^2 + x^4.$$

Moreover, these three polynomials are linearly independent, since the only way to write the zero polynomial as a linear combination of u_1, u_2, u_3 is to take all coefficients to be zero. Thus, $\{u_1, u_2, u_3\}$ is a basis of U .

- (b) We can extend the basis $\{u_1, u_2, u_3\}$ of U to a basis of $\mathcal{P}_4(\mathbb{F})$ by adding two more polynomials that are linearly independent of u_1, u_2, u_3 . For example, we can add the polynomials

$$u_4 = x^2, \quad u_5 = x^3,$$

which are linearly independent of u_1, u_2, u_3 since they are not in U . Thus, $\{u_1, u_2, u_3, u_4, u_5\}$ is a basis of $\mathcal{P}_4(\mathbb{F})$.

- (c) Let $W = \text{span}(u_4, u_5) = \text{span}(x^2, x^3)$. Then we have

$$\mathcal{P}_4(\mathbb{F}) = U + W,$$

since every polynomial in $\mathcal{P}_4(\mathbb{F})$ can be written as a sum of a polynomial in U and a polynomial in W . Moreover, if $p \in U \cap W$, then $p = \alpha x^2 + \beta x^3$ for some $\alpha, \beta \in \mathbb{F}$. Since $p(2) = p(5) = p(6)$, we have

$$4\alpha + 8\beta = 25\alpha + 125\beta = 36\alpha + 216\beta.$$

Solving this system of equations, we find that $\alpha = \beta = 0$. Thus, $U \cap W = \{0\}$, and so we have

$$\mathcal{P}_4(\mathbb{F}) = U \oplus W. \quad \blacksquare$$

Exercise 2C7

(a) Let

$$U = \left\{ p \in \mathcal{P}_4(\mathbb{R}) \mid \int_{-1}^1 p = 0 \right\}.$$

Find a basis of U .(b) Extend the basis in (a) to a basis of $\mathcal{P}_4(\mathbb{R})$.(c) Find a subspace W of $\mathcal{P}_4(\mathbb{R})$ such that $\mathcal{P}_4(\mathbb{R}) = U \oplus W$.**Solution.**(a) Let $p = a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 \in U$. Then we have

$$\begin{aligned} \int_{-1}^1 p &= \int_{-1}^1 (a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4) dx \\ &= \int_{-1}^1 (a_0 + a_2x^2 + a_4x^4) dx \\ &= 2 \int_0^1 (a_0 + a_2x^2 + a_4x^4) dx \\ &= 2 \left(a_0 + \frac{a_2}{3} + \frac{a_4}{5} \right) = 0. \end{aligned}$$

We can solve this equation for a_4 in terms of a_0, a_2 :

$$a_4 = -5a_2 - 15a_0.$$

Then we can write any polynomial in U as a linear combination of the following four polynomials:

$$u_1 = 1 - 15x^4, \quad u_2 = x, \quad u_3 = x^2 - 5x^4, \quad u_4 = x^3.$$

Moreover, these four polynomials are linearly independent, since the only way to write the zero polynomial as a linear combination of u_1, u_2, u_3, u_4 is to take all coefficients to be zero. Thus, $\{u_1, u_2, u_3, u_4\}$ is a basis of U .(b) We can extend the basis $\{u_1, u_2, u_3, u_4\}$ of U to a basis of $\mathcal{P}_4(\mathbb{R})$ by adding one more polynomial that is linearly independent of u_1, u_2, u_3, u_4 . For example, we can add the polynomial

$$u_5 = x^4,$$

which is linearly independent of u_1, u_2, u_3, u_4 since it is not in U . Thus, $\{u_1, u_2, u_3, u_4, u_5\}$ is a basis of $\mathcal{P}_4(\mathbb{R})$.(c) Let $W = \text{span } u_5 = \text{span } x^4$. Then we have

$$\mathcal{P}_4(\mathbb{R}) = U + W,$$

since every polynomial in $\mathcal{P}_4(\mathbb{R})$ can be written as a sum of a polynomial in U and a polynomial in W . Moreover, if $p \in U \cap W$, then $p = \alpha x^4$ for some $\alpha \in \mathbb{R}$. Since the integral of an even function over a symmetric interval is nonzero unless the function is identically zero, we have $\alpha = 0$. Thus, $U \cap W = \{0\}$, and so we have

$$\mathcal{P}_4(\mathbb{R}) = U \oplus W. \quad \blacksquare$$

Exercise 2C8

Suppose $v_1, \dots, v_m$ is linearly independent in V and $w \in V$. Prove that

$$\dim \text{span}(v_1 + w, \dots, v_m + w) \geq m - 1.$$

Solution. Observe that $(v_i + w) - (v_{i+1} + w) = v_i - v_{i+1}$ for $1 \leq i \leq m - 1$. Then the list of vectors

$$v_1 - v_2, v_2 - v_3, \dots, v_{m-1} - v_m$$

is contained in $\text{span}(v_1 + w, \dots, v_m + w)$. We claim that this list is linearly independent. Suppose that

$$\alpha_1(v_1 - v_2) + \alpha_2(v_2 - v_3) + \dots + \alpha_{m-1}(v_{m-1} - v_m) = 0$$

for some $\alpha_1, \dots, \alpha_{m-1} \in \mathbb{F}$. Then we can rewrite this equation as

$$\alpha_1 v_1 + (\alpha_2 - \alpha_1)v_2 + \dots + (\alpha_{m-1} - \alpha_{m-2})v_{m-1} - \alpha_{m-1}v_m = 0.$$

Since $v_1, \dots, v_m$ is linearly independent, we must have $\alpha_1 = 0$, $\alpha_2 - \alpha_1 = 0$, $\dots$, $\alpha_{m-1} - \alpha_{m-2} = 0$, and $-\alpha_{m-1} = 0$. This implies that $\alpha_1 = \dots = \alpha_{m-1} = 0$. Thus, the list is linearly independent, and so we have

$$\dim \text{span}(v_1 + w, \dots, v_m + w) \geq m - 1. \quad \blacksquare$$

Exercise 2C9

Suppose m is a positive integer and $p_0, p_1, \dots, p_m \in \mathcal{P}(\mathbb{F})$ are such that each p_k has degree k . Prove that $p_0, p_1, \dots, p_m$ is a basis of $\mathcal{P}_m(\mathbb{F})$.

Solution. Since each p_k has degree k , we can write each polynomial as

$$p_k = a_{k0} + a_{k1}x + \dots + a_{kk}x^k$$

where $a_{ij} \in \mathbb{F}$ for all $0 \leq i \leq j \leq k$ and $a_{kk} \neq 0$. We claim that $p_0, p_1, \dots, p_m$ is linearly independent. Suppose that

$$\alpha_0 p_0 + \alpha_1 p_1 + \dots + \alpha_m p_m = 0$$

for some $\alpha_0, \dots, \alpha_m \in \mathbb{F}$. Then we can write this equation as

$$(\alpha_0 a_{00} + \alpha_1 a_{10} + \dots + \alpha_m a_{m0}) + (\alpha_1 a_{11} + \dots + \alpha_m a_{m1})x + \dots + (\alpha_m a_{mm})x^m = 0.$$

Since the zero polynomial has all coefficients equal to zero, we must have

$$\alpha_m a_{mm} = 0.$$

Since $a_{mm} \neq 0$, we must have $\alpha_m = 0$. Next, the coefficient of the x^{m-1} term gives us

$$\alpha_{m-1} a_{m-1, m-1} + \alpha_m a_{m, m-1} = 0.$$

Since $\alpha_m = 0$ and $a_{m-1, m-1} \neq 0$, we must have $\alpha_{m-1} = 0$. Continuing in this way, we find that $\alpha_{m-2} = \dots = \alpha_0 = 0$. Thus, the list is linearly independent. Moreover, since $\dim \mathcal{P}_m(\mathbb{F}) = m + 1$, any linearly independent list of $m + 1$ vectors in $\mathcal{P}_m(\mathbb{F})$ is a basis. Therefore, $p_0, p_1, \dots, p_m$ is a basis of $\mathcal{P}_m(\mathbb{F})$. $\blacksquare$

Exercise 2C10

Suppose m is a positive integer. For $0 \leq k \leq m$, let

$$p_k(x) = x^k(1-x)^{m-k}.$$

Show that $p_0, \dots, p_m$ is a basis of $\mathcal{P}_m(\mathbb{F})$.

Solution. We first determine what exponents appear in the polynomial p_k . Using the Binomial Theorem, we can expand $(1-x)^{m-k}$ and obtain the expression for $p_k(x)$ as a polynomial in x :

$$p_k(x) = x^k \sum_{i=0}^{m-k} \binom{m-k}{i} (-1)^i x^i = \sum_{i=0}^{m-k} \binom{m-k}{i} (-1)^i x^{k+i}.$$

Thus, the exponents that appear in p_k are $k, k+1, \dots, m$, and we may write p_k as

$$p_k(x) = a_{kk}x^k + a_{k,k+1}x^{k+1} + \dots + a_{km}x^m$$

where $a_{kk} = 1$ and $a_{kj} \in \mathbb{F}$ for all $k \leq j \leq m$. We claim that $p_0, p_1, \dots, p_m$ is linearly independent. Suppose that

$$\alpha_0 p_0 + \alpha_1 p_1 + \dots + \alpha_m p_m = 0$$

for some $\alpha_0, \dots, \alpha_m \in \mathbb{F}$. Then we can write this equation as

$$(\alpha_0 a_{00}) + (\alpha_0 a_{01} + \alpha_1 a_{11})x + \dots + (\alpha_0 a_{0m} + \dots + \alpha_m a_{mm})x^m = 0.$$

Since the zero polynomial has all coefficients equal to zero, and $a_{00} \neq 0$ by definition of p_0 , we must have $\alpha_0 = 0$. Next, the coefficient of the x is $\alpha_0 a_{01} + \alpha_1 a_{11} = 0$. Since $\alpha_0 = 0$ and $a_{11} \neq 0$, we must have $\alpha_1 = 0$. Continuing in this way, we find that $\alpha_2 = \dots = \alpha_m = 0$. Thus, the list is linearly independent. Moreover, since $\dim \mathcal{P}_m(\mathbb{F}) = m+1$, any linearly independent list of $m+1$ vectors in $\mathcal{P}_m(\mathbb{F})$ is a basis. Therefore, $p_0, p_1, \dots, p_m$ is a basis of $\mathcal{P}_m(\mathbb{F})$. ■

Exercise 2C11

Suppose U and W are both four-dimensional subspaces of $\mathbb{C}^6$. Prove that there exist two vectors in $U \cap W$ such that neither of these vectors is a scalar multiple of the other.

Solution. Let U and W be four-dimensional subspaces of $\mathbb{C}^6$. By the dimension formula for the sum of two subspaces, we have

$$\dim(U+W) = \dim U + \dim W - \dim(U \cap W) = 4 + 4 - \dim(U \cap W) = 8 - \dim(U \cap W).$$

Since $U+W$ is a subspace of $\mathbb{C}^6$, we have $\dim(U+W) \leq 6$. Thus, we have

$$8 - \dim(U \cap W) \leq 6,$$

which implies that $\dim(U \cap W) \geq 2$. Therefore, there exist at least two linearly independent vectors in $U \cap W$, and neither of these vectors is a scalar multiple of the other. ■

Exercise 2C12

Suppose that U and W are subspaces of $\mathbb{R}^8$ such that $\dim U = 3$, $\dim W = 5$, and $U + W = \mathbb{R}^8$. Prove that $\mathbb{R}^8 = U \oplus W$.

Solution. By the dimension formula for the sum of two subspaces, we have

$$\dim(U + W) = \dim U + \dim W - \dim(U \cap W) = 3 + 5 - \dim(U \cap W) = 8 - \dim(U \cap W).$$

Since $U + W = \mathbb{R}^8$, we have $\dim(U + W) = 8$. Thus, we have

$$8 - \dim(U \cap W) = 8,$$

which implies that $\dim(U \cap W) = 0$. Therefore, $U \cap W = \{0\}$, and so we have

$$\mathbb{R}^8 = U \oplus W. \quad \blacksquare$$

Exercise 2C13

Suppose U and W are both five-dimensional subspaces of $\mathbb{R}^8$. Prove that $U \cap W \neq \{0\}$.

Solution. By the dimension formula for the sum of two subspaces, we have

$$\dim(U + W) = \dim U + \dim W - \dim(U \cap W) = 5 + 5 - \dim(U \cap W) = 10 - \dim(U \cap W).$$

Since $U + W$ is a subspace of $\mathbb{R}^8$, we have $\dim(U + W) \leq 8$. Thus, we have

$$10 - \dim(U \cap W) \leq 8,$$

which implies that $\dim(U \cap W) \geq 2$. Therefore, $U \cap W$ is nontrivial, and so we have $U \cap W \neq \{0\}$. $\blacksquare$

Exercise 2C14

Suppose V is a ten-dimensional vector space and V_1, V_2, V_3 are subspaces of V with $\dim V_1 = \dim V_2 = \dim V_3 = 7$. Prove that $V_1 \cap V_2 \cap V_3 \neq \{0\}$.

Solution. Let $W = V_1 \cap V_2$. By the dimension formula for the sum of two subspaces, we have

$$\dim(V_1 + V_2) = \dim V_1 + \dim V_2 - \dim W = 7 + 7 - \dim W = 14 - \dim W.$$

Since $V_1 + V_2$ is a subspace of V , we have $\dim(V_1 + V_2) \leq 10$. Thus, we have $14 - \dim W \leq 10$, which implies that $\dim W \geq 4$. Therefore, $V_1 \cap V_2$ is at least four-dimensional. Now, consider the subspace $W \cap V_3 = V_1 \cap V_2 \cap V_3$. By the dimension formula for the sum of two subspaces, we have

$$\dim(W + V_3) = \dim W + \dim V_3 - \dim(W \cap V_3) = \dim W + 7 - \dim(W \cap V_3).$$

Since $W + V_3$ is a subspace of V , we have $\dim(W + V_3) \leq 10$. Thus, we have

$$\dim W + 7 - \dim(W \cap V_3) \leq 10,$$

which implies that $\dim(W \cap V_3) \geq \dim W + 7 - 10 = \dim W - 3$. Since $\dim W \geq 4$, we have $\dim(W \cap V_3) \geq 1$. Therefore, $V_1 \cap V_2 \cap V_3$ is nontrivial, and so we have $V_1 \cap V_2 \cap V_3 \neq \{0\}$. $\blacksquare$

Exercise 2C15

Suppose V is finite-dimensional and V_1, V_2, V_3 are subspaces of V with $\dim V_1 + \dim V_2 + \dim V_3 > 2 \dim V$. Prove that $V_1 \cap V_2 \cap V_3 \neq \{0\}$.

Solution. Let $W = V_1 \cap V_2$. By the dimension formula for the sum of two subspaces, we have

$$\dim(V_1 + V_2) = \dim V_1 + \dim V_2 - \dim W.$$

Since $V_1 + V_2$ is a subspace of V , we have $\dim(V_1 + V_2) \leq \dim V$. Thus, we have

$$\dim V_1 + \dim V_2 - \dim W \leq \dim V,$$

which implies that $\dim W \geq \dim V_1 + \dim V_2 - \dim V$. Now, consider the subspace $W \cap V_3 = V_1 \cap V_2 \cap V_3$. By the dimension formula for the sum of two subspaces, we have

$$\dim(W + V_3) = \dim W + \dim V_3 - \dim(W \cap V_3).$$

Since $W + V_3$ is a subspace of V , we have $\dim(W + V_3) \leq \dim V$. Thus, we have

$$\dim W + \dim V_3 - \dim(W \cap V_3) \leq \dim V,$$

which implies that $\dim(W \cap V_3) \geq \dim W + \dim V_3 - \dim V$. Since $\dim W \geq \dim V_1 + \dim V_2 - \dim V$, we have

$$\dim(W \cap V_3) \geq (\dim V_1 + \dim V_2 - \dim V) + \dim V_3 - \dim V = \dim V_1 + \dim V_2 + \dim V_3 - 2 \dim V.$$

Since $\dim V_1 + \dim V_2 + \dim V_3 > 2 \dim V$, we have $\dim(W \cap V_3) > 0$. Therefore, $V_1 \cap V_2 \cap V_3$ is nontrivial, and so we have $V_1 \cap V_2 \cap V_3 \neq \{0\}$. ■

Exercise 2C16

Suppose V is finite-dimensional and U is a subspace of V with $U \neq V$. Let $n = \dim V$ and $m = \dim U$. Prove that there exist $n - m$ subspaces of V , each of dimension $n - 1$, whose intersection equals U .

Solution. Let U be a subspace of V with $\dim U = m < n = \dim V$. We can extend a basis of U to a basis of V . Let $\{u_1, \dots, u_m\}$ be a basis of U , and extend it to a basis of V by adding vectors $v_1, \dots, v_{n-m}$. Then $\{u_1, \dots, u_m, v_1, \dots, v_{n-m}\}$ is a basis of V . For each $i = 1, \dots, n - m$, let

$$W_i = \text{span}(u_1, \dots, u_m, v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_{n-m}).$$

Then each W_i is a subspace of V with dimension $n - 1$, since it is spanned by $n - 1$ linearly independent vectors. Moreover, we have

$$U = W_1 \cap W_2 \cap \dots \cap W_{n-m},$$

since any vector in the intersection must be a linear combination of the basis vectors of U , and any vector in U is contained in all the subspaces W_i . Therefore, there exist $n - m$ subspaces of V , each of dimension $n - 1$, whose intersection equals U . ■

Exercise 2C17

Suppose that $V_1, \dots, V_m$ are finite-dimensional subspaces of V . Prove that $V_1 + \dots + V_m$ is finite-dimensional and

$$\dim(V_1 + \dots + V_m) \leq \dim V_1 + \dots + \dim V_m.$$

Solution. We will prove the statement by induction on m . For the base case, let $m = 2$. Then we have

$$V_1 + V_2 = V_1 + V_2,$$

which is finite-dimensional and

$$\dim(V_1 + V_2) = \dim V_1 + \dim V_2 - \dim(V_1 \cap V_2) \leq \dim V_1 + \dim V_2.$$

Hence, the statement holds for $m = 2$. Now, suppose that the statement holds for some positive integer m . That is, suppose that if $V_1, \dots, V_m$ are finite-dimensional subspaces of V , then $V_1 + \dots + V_m$ is finite-dimensional and

$$\dim(V_1 + \dots + V_m) \leq \dim V_1 + \dots + \dim V_m.$$

We want to show that the statement holds for $m + 1$. Let $V_1, \dots, V_{m+1}$ be finite-dimensional subspaces of V . By the induction hypothesis, we know that $W = V_1 + \dots + V_m$ is finite-dimensional and

$$\dim W \leq \dim V_1 + \dots + \dim V_m.$$

Then we have

$$\dim(W + V_{m+1}) = \dim W + \dim V_{m+1} - \dim(W \cap V_{m+1}).$$

Since $W \cap V_{m+1}$ is a subspace of V , we have $\dim(W \cap V_{m+1}) \geq 0$. Thus, we have

$$\dim(W + V_{m+1}) \leq \dim W + \dim V_{m+1} \leq \dim V_1 + \dots + \dim V_m + \dim V_{m+1}.$$

Therefore, $V_1 + \dots + V_{m+1}$ is finite-dimensional and

$$\dim(V_1 + \dots + V_{m+1}) \leq \dim V_1 + \dots + \dim V_m + \dim V_{m+1}. \quad \blacksquare$$

Exercise 2C18

Suppose V is finite-dimensional, with $\dim V = n \geq 1$. Prove that there exist one-dimensional subspaces $V_1, \dots, V_n$ of V such that

$$V = V_1 \oplus \dots \oplus V_n.$$

Solution. Let $v_1, \dots, v_n$ be a basis of V . For each $i = 1, \dots, n$, let $V_i = \text{span } v_i$. Then each V_i is a one-dimensional subspace of V , since it is spanned by a single nonzero vector. Moreover, we have

$$V = V_1 + \dots + V_n,$$

and the sum is direct because the basis vectors are linearly independent. Therefore, we have

$$V = V_1 \oplus \dots \oplus V_n. \quad \blacksquare$$

Exercise 2C19

Explain why you might guess, motivated by analogy with the formula for the number of elements in the union of three finite sets, that if V_1, V_2, V_3 are subspaces of a finite-dimensional vector space, then

$$\begin{aligned} \dim(V_1 + V_2 + V_3) = & \dim V_1 + \dim V_2 + \dim V_3 \\ & - \dim(V_1 \cap V_2) - \dim(V_1 \cap V_3) - \dim(V_2 \cap V_3) \\ & + \dim(V_1 \cap V_2 \cap V_3). \end{aligned}$$

Then either prove the formula above or give a counterexample.

Solution. One may guess that the given formula above is true by analogy with the formula for the number of elements in the union of three finite sets. However, this erroneously conflates that the idea of subspace sums is equivalent to the idea of set unions. In fact, the formula is not true in general. For example, let $V = \mathbb{R}^2$, and let

$$V_1 = \text{span}((1, 0)), \quad V_2 = \text{span}((0, 1)), \quad V_3 = \text{span}((1, 1)).$$

It is easy to see that $\dim V_1 = \dim V_2 = \dim V_3 = 1$, but $\dim(V_1 \cap V_2) = \dim(V_1 \cap V_3) = \dim(V_2 \cap V_3) = 0$ and $\dim(V_1 \cap V_2 \cap V_3) = 0$. Thus, the right-hand side of the formula is

$$1 + 1 + 1 - 0 - 0 - 0 + 0 = 3,$$

but the left-hand side is $\dim(V_1 + V_2 + V_3) = \dim \mathbb{R}^2 = 2$. Therefore, the formula is not true in general. ■

Exercise 2C20

Prove that if V_1, V_2 , and V_3 are subspaces of a finite-dimensional vector space, then

$$\begin{aligned} \dim(V_1 + V_2 + V_3) = & \dim V_1 + \dim V_2 + \dim V_3 \\ & - \frac{\dim(V_1 \cap V_2) + \dim(V_1 \cap V_3) + \dim(V_2 \cap V_3)}{3} \\ & - \frac{\dim((V_1 + V_2) \cap V_3) + \dim((V_1 + V_3) \cap V_2) + \dim((V_2 + V_3) \cap V_1)}{3}. \end{aligned}$$

Solution. Note that we need to account for all pairs of subspaces all pairs of sums of subspaces, since the sum of subspaces is not equivalent to the union of sets. We can use the dimension formula repeatedly to obtain the answer. To that end, define the following subspaces:

$$W_1 = V_1 + V_2, \quad W_2 = V_1 + V_3, \quad W_3 = V_2 + V_3.$$

Applying the dimension formula to each of W_1, W_2, W_3 , we have

$$\begin{aligned} \dim W_1 &= \dim V_1 + \dim V_2 - \dim(V_1 \cap V_2), \\ \dim W_2 &= \dim V_1 + \dim V_3 - \dim(V_1 \cap V_3), \\ \dim W_3 &= \dim V_2 + \dim V_3 - \dim(V_2 \cap V_3). \end{aligned}$$

Next, we can apply the dimension formula to each of the sums $W_1 + V_3$, $W_2 + V_2$, $W_3 + V_1$:

$$\begin{aligned}
 \dim(W_1 + V_3) &= \dim W_1 + \dim V_3 - \dim(W_1 \cap V_3), \\
 &= \dim V_1 + \dim V_2 - \dim(V_1 \cap V_2) + \dim V_3 - \dim(W_1 \cap V_3) \\
 \dim(W_2 + V_2) &= \dim W_2 + \dim V_2 - \dim(W_2 \cap V_2), \\
 &= \dim V_1 + \dim V_3 - \dim(V_1 \cap V_3) + \dim V_2 - \dim(W_2 \cap V_2) \\
 \dim(W_3 + V_1) &= \dim W_3 + \dim V_1 - \dim(W_3 \cap V_1), \\
 &= \dim V_2 + \dim V_3 - \dim(V_2 \cap V_3) + \dim V_1 - \dim(W_3 \cap V_1).
 \end{aligned}$$

Note that $W_1 + V_3 = W_2 + V_2 = W_3 + V_1 = V_1 + V_2 + V_3$. Thus, we can add the three equations above and divide by 3 to obtain the desired formula:

$$\begin{aligned}
 \dim(V_1 + V_2 + V_3) &= \dim V_1 + \dim V_2 + \dim V_3 \\
 &\quad - \frac{\dim(V_1 \cap V_2) + \dim(V_1 \cap V_3) + \dim(V_2 \cap V_3)}{3} \\
 &\quad - \frac{\dim((V_1 + V_2) \cap V_3) + \dim((V_1 + V_3) \cap V_2) + \dim((V_2 + V_3) \cap V_1)}{3}. \quad \blacksquare
 \end{aligned}$$

3 Linear Maps

3A Vector Space of Linear Maps

Exercise 3A1

Suppose $b, c \in \mathbb{R}$. Define $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ by

$$T(x, y, z) = (2x - 4y + 3z + b, 6x + cxyz).$$

Show that T is linear if and only if $b = c = 0$.

Solution. Suppose T is linear. Then

$$T(1, 1, 1) = T(1, 0, 0) + T(0, 1, 0) + T(0, 0, 1) = (2 + b, 6) + (-4 + b, 0) + (3 + b, 0) = (1 + 3b, 6).$$

On the other hand,

$$T(1, 1, 1) = (2 - 4 + 3 + b, 6 + c) = (1 + b, 6 + c).$$

Comparing the two expressions, we get $1 + 3b = 1 + b$ and $6 = 6 + c$, which implies $b = 0$ and $c = 0$.

Conversely, if $b = c = 0$, then $T(x, y, z) = (2x - 4y + 3z, 6x)$. Let $(x_1, y_1, z_1), (x_2, y_2, z_2) \in \mathbb{R}^3$ and $\alpha \in \mathbb{R}$. Then

$$\begin{aligned} T(x_1 + x_2, y_1 + y_2, z_1 + z_2) &= (2(x_1 + x_2) - 4(y_1 + y_2) + 3(z_1 + z_2), 6(x_1 + x_2)) \\ &= (2x_1 + 2x_2 - 4y_1 - 4y_2 + 3z_1 + 3z_2, 6x_1 + 6x_2) \\ &= (2x_1 - 4y_1 + 3z_1, 6x_1) + (2x_2 - 4y_2 + 3z_2, 6x_2) \\ &= T(x_1, y_1, z_1) + T(x_2, y_2, z_2), \end{aligned}$$

and

$$\begin{aligned} T(\alpha x_1, \alpha y_1, \alpha z_1) &= (2(\alpha x_1) - 4(\alpha y_1) + 3(\alpha z_1), 6(\alpha x_1)) \\ &= (\alpha(2x_1 - 4y_1 + 3z_1), \alpha(6x_1)) \\ &= \alpha(2x_1 - 4y_1 + 3z_1, 6x_1) \\ &= \alpha T(x_1, y_1, z_1). \end{aligned}$$

Therefore, T is linear if and only if $b = c = 0$. ■

Exercise 3A2

Suppose $b, c \in \mathbb{R}$. Define $T : \mathcal{P}(\mathbb{R}) \rightarrow \mathbb{R}^2$ by

$$Tp = \left(3p(4) + 5p'(6) + bp(1)p(2), \int_{-1}^2 x^3 p(x) dx + c \sin p(0) \right).$$

Show that T is linear if and only if $b = c = 0$.

Solution. Suppose T is linear. Then for any $p \in \mathcal{P}(\mathbb{R})$, we have $T(2p) = 2Tp$. Looking at the first component of $T(2p)$, we have

$$3(2p)(4) + 5(2p)'(6) + b(2p)(1)(2p)(2) = 6p(4) + 10p'(6) + 4bp(1)p(2).$$

On the other hand, the first component of $2Tp$ is

$$2(3p(4) + 5p'(6) + bp(1)p(2)) = 6p(4) + 10p'(6) + 2bp(1)p(2).$$

Comparing the two expressions, we get $4bp(1)p(2) = 2bp(1)p(2)$ for all $p \in \mathcal{P}(\mathbb{R})$. Since there exist polynomials p such that $p(1)p(2) \neq 0$, we must have $b = 0$.

The second component of $T(2p)$ must be equal to the second component of $2Tp$. We have

$$\int_{-1}^2 x^3(2p)(x) dx + c \sin(2p(0)) = 2 \int_{-1}^2 x^3 p(x) dx + 2c \sin(p(0)).$$

The integral terms may be subtracted, so we get $c \sin(2p(0)) = 2c \sin(p(0))$ for all $p \in \mathcal{P}(\mathbb{R})$. If $c \neq 0$, then we can divide both sides by c to get $\sin(2p(0)) = 2 \sin(p(0))$, which implies $\sin(2x) = 2 \sin(x)$ for all $x \in \mathbb{R}$. Then $\cos(x) = 1$ for all $x \in \mathbb{R}$, which is a contradiction. Therefore, $c = 0$.

Conversely, if $b = c = 0$, let $p, q \in \mathcal{P}(\mathbb{R})$ and $\alpha \in \mathbb{R}$. Then

$$\begin{aligned} T(p+q) &= \left(3(p+q)(4) + 5(p+q)'(6), \int_{-1}^2 x^3(p+q)(x) dx \right) \\ &= \left(3p(4) + 3q(4) + 5p'(6) + 5q'(6), \int_{-1}^2 x^3 p(x) dx + \int_{-1}^2 x^3 q(x) dx \right) \\ &= \left(3p(4) + 5p'(6), \int_{-1}^2 x^3 p(x) dx \right) + \left(3q(4) + 5q'(6), \int_{-1}^2 x^3 q(x) dx \right) \\ &= Tp + Tq, \end{aligned}$$

and

$$\begin{aligned} T(\alpha p) &= \left(3(\alpha p)(4) + 5(\alpha p)'(6), \int_{-1}^2 x^3(\alpha p)(x) dx \right) \\ &= \left(\alpha(3p(4) + 5p'(6)), \alpha \int_{-1}^2 x^3 p(x) dx \right) \\ &= \alpha \left(3p(4) + 5p'(6), \int_{-1}^2 x^3 p(x) dx \right) \\ &= \alpha Tp. \end{aligned}$$

Therefore, T is linear if and only if $b = c = 0$. ■

Exercise 3A3

Suppose that $T \in \mathcal{L}(\mathbb{F}^n, \mathbb{F}^m)$. Show that there exist scalars $A_{j,k} \in \mathbb{F}$ for $j = 1, \dots, m$ and $k = 1, \dots, n$ such that

$$T(x_1, \dots, x_n) = (A_{1,1}x_1 + \dots + A_{1,n}x_n, \dots, A_{m,1}x_1 + \dots + A_{m,n}x_n)$$

for every $(x_1, \dots, x_n) \in \mathbb{F}^n$.

Solution. Let $e_1, \dots, e_n$ be the standard basis of $\mathbb{F}^n$. Then for each $j = 1, \dots, m$, we have

$$T(e_k) = (A_{1,k}, A_{2,k}, \dots, A_{m,k})$$

for some scalars $A_{jk} \in \mathbb{F}$. Now let $(x_1, \dots, x_n) \in \mathbb{F}^n$. Then

$$\begin{aligned} T(x_1, \dots, x_n) &= T(x_1 e_1 + \dots + x_n e_n) \\ &= x_1 T(e_1) + \dots + x_n T(e_n) \\ &= x_1 (A_{1,1}, A_{2,1}, \dots, A_{m,1}) + \dots + x_n (A_{1,n}, A_{2,n}, \dots, A_{m,n}) \\ &= (A_{1,1}x_1 + \dots + A_{1,n}x_n, A_{2,1}x_1 + \dots + A_{2,n}x_n, \dots, A_{m,1}x_1 + \dots + A_{m,n}x_n). \end{aligned}$$

Therefore,

$$T(x_1, \dots, x_n) = (A_{1,1}x_1 + \dots + A_{1,n}x_n, A_{2,1}x_1 + \dots + A_{2,n}x_n, \dots, A_{m,1}x_1 + \dots + A_{m,n}x_n). \quad \blacksquare$$

Exercise 3A4

Suppose $T \in \mathcal{L}(V, W)$ and $v_1, \dots, v_m$ is a list of vectors in V such that $Tv_1, \dots, Tv_m$ is a linearly independent list in W . Prove that $v_1, \dots, v_m$ is linearly independent.

Solution. Suppose that $a_1 v_1 + \dots + a_m v_m = 0$ for some scalars $a_1, \dots, a_m \in \mathbb{F}$. Applying T to both sides, we get

$$a_1 T v_1 + \dots + a_m T v_m = T(0) = 0.$$

Since $Tv_1, \dots, Tv_m$ is linearly independent, it follows that $a_1 = \dots = a_m = 0$. Therefore, $v_1, \dots, v_m$ is linearly independent. $\blacksquare$

Exercise 3A5

Prove that $\mathcal{L}(V, W)$ is a vector space, as was asserted in 3.6.

Solution. We must first ensure that the operations of addition and scalar multiplication are well-defined, i.e., if $S, T \in \mathcal{L}(V, W)$ and $\alpha \in \mathbb{F}$, then $S + T \in \mathcal{L}(V, W)$ and $\alpha T \in \mathcal{L}(V, W)$. To that end, let $v, u \in V$. Then

$$\begin{aligned} (S + T)(v + u) &= S(v + u) + T(v + u) \\ &= (Sv + Su) + (Tv + Tu) \\ &= (Sv + Tv) + (Su + Tu) \\ &= (S + T)v + (S + T)u, \end{aligned}$$

and

$$\begin{aligned} (\alpha T)(v + u) &= \alpha T(v + u) \\ &= \alpha(Tv + Tu) \\ &= \alpha Tv + \alpha Tu \\ &= (\alpha T)v + (\alpha T)u. \end{aligned}$$

Also, for any $v \in V$ and $\alpha \in \mathbb{F}$,

$$\begin{aligned} (S + T)(\alpha v) &= S(\alpha v) + T(\alpha v) \\ &= \alpha Sv + \alpha Tv \\ &= \alpha(Sv + Tv) \\ &= \alpha(S + T)v, \end{aligned}$$

and

$$\begin{aligned}
 (\alpha T)(\beta v) &= \alpha T(\beta v) \\
 &= \alpha(\beta Tv) \\
 &= (\alpha\beta)Tv \\
 &= (\alpha\beta)Tv.
 \end{aligned}$$

Therefore, $S + T \in \mathcal{L}(V, W)$ and $\alpha T \in \mathcal{L}(V, W)$. We now check the vector space axioms:

1. **Commutativity:** Let $S, T \in \mathcal{L}(V, W)$. Then for any $v \in V$,

$$\begin{aligned}
 (S + T)v &= Sv + Tv \\
 &= Tv + Sv \\
 &= (T + S)v.
 \end{aligned}$$

2. **Associativity:** Let $R, S, T \in \mathcal{L}(V, W)$. Then for any $v \in V$,

$$\begin{aligned}
 ((R + S) + T)v &= (R + S)v + Tv \\
 &= (Rv + Sv) + Tv \\
 &= Rv + (Sv + Tv) \\
 &= Rv + (S + T)v \\
 &= (R + (S + T))v.
 \end{aligned}$$

Similarly, for any $\alpha, \beta \in \mathbb{F}$ and $T \in \mathcal{L}(V, W)$,

$$\begin{aligned}
 ((\alpha\beta)T)v &= (\alpha\beta)(Tv) \\
 &= \alpha(\beta Tv) \\
 &= \alpha(\beta T)v \\
 &= (\alpha(\beta T))v.
 \end{aligned}$$

3. **Additive identity:** Let $0 \in \mathcal{L}(V, W)$ be the zero map defined by $0v = 0$ for all $v \in V$. Then for any $T \in \mathcal{L}(V, W)$ and $v \in V$,

$$(T + 0)v = Tv + 0v = Tv + 0 = Tv.$$

4. **Inverse:** Let $T \in \mathcal{L}(V, W)$. Define $-T \in \mathcal{L}(V, W)$ by $(-T)v = -Tv$ for all $v \in V$. Then for any $v \in V$,

$$(T + (-T))v = Tv + (-T)v = Tv - Tv = 0 = 0v.$$

5. **Multiplicative identity:** Let $1 \in \mathbb{F}$ be the multiplicative identity. Then for any $T \in \mathcal{L}(V, W)$ and $v \in V$,

$$(1T)v = 1(Tv) = Tv.$$

6. **Distributive properties:** Let $T, S \in \mathcal{L}(V, W)$ and $\alpha, \beta \in \mathbb{F}$. Then for any $v \in V$,

$$\begin{aligned}
 (\alpha(T + S))v &= (\alpha T + \alpha S)v \\
 &= (\alpha T)v + (\alpha S)v \\
 &= \alpha(Tv) + \alpha(Sv) \\
 &= \alpha(Tv + Sv) \\
 &= \alpha((T + S)v),
 \end{aligned}$$

and

$$\begin{aligned}
 ((\alpha + \beta)T)v &= (\alpha + \beta)(Tv) \\
 &= \alpha(Tv) + \beta(Tv) \\
 &= (\alpha T)v + (\beta T)v \\
 &= (\alpha T + \beta T)v.
 \end{aligned}$$

■

Exercise 3A6

Prove that multiplication of linear maps has the associative, identity, and distributive properties asserted in 3.8.

- (a) **Associativity:** $(T_1 T_2) T_3 = T_1 (T_2 T_3)$ whenever T_1, T_2 , and T_3 are linear maps such that the products make sense (meaning T_3 maps into the domain of T_2 , and T_2 maps into the domain of T_1).
- (b) **Identity:** $TI = IT = T$ whenever $T \in \mathcal{L}(V, W)$; here the first I is the identity operator on V and the second I is the identity operator on W .
- (c) **Distributive properties:** $(S_1 + S_2)T = S_1 T + S_2 T$ and $S(T_1 + T_2) = ST_1 + ST_2$ whenever $T, T_1, T_2 \in \mathcal{L}(U, V)$ and $S, S_1, S_2 \in \mathcal{L}(V, W)$.

Solution.

- (a) Let $T_1 \in \mathcal{L}(V, W)$, $T_2 \in \mathcal{L}(U, V)$, and $T_3 \in \mathcal{L}(X, U)$. Then for any $x \in X$,

$$\begin{aligned}
 ((T_1 T_2) T_3)x &= (T_1 T_2)(T_3 x) \\
 &= T_1(T_2(T_3 x)) \\
 &= T_1((T_2 T_3)x) \\
 &= (T_1(T_2 T_3))x.
 \end{aligned}$$

- (b) Let $T \in \mathcal{L}(V, W)$. Then for any $v \in V$,

$$(TI)v = T(Iv) = Tv \quad \text{and} \quad (IT)v = I(Tv) = Tv.$$

- (c) Let $T, T_1, T_2 \in \mathcal{L}(U, V)$ and $S, S_1, S_2 \in \mathcal{L}(V, W)$. Then for any $u \in U$,

$$\begin{aligned}
 ((S_1 + S_2)T)u &= (S_1 + S_2)(Tu) \\
 &= S_1(Tu) + S_2(Tu) \\
 &= (S_1 T)u + (S_2 T)u \\
 &= (S_1 T + S_2 T)u,
 \end{aligned}$$

and

$$\begin{aligned}
 (S(T_1 + T_2))u &= S((T_1 + T_2)u) \\
 &= S(T_1 u + T_2 u) \\
 &= S(T_1 u) + S(T_2 u) \\
 &= (ST_1)u + (ST_2)u \\
 &= (ST_1 + ST_2)u.
 \end{aligned}$$

■

Exercise 3A7

Show that every linear map from a one-dimensional vector space to itself is multiplication by some scalar. More precisely, prove that if $\dim V = 1$ and $T \in \mathcal{L}(V)$, then there exists $\lambda \in \mathbb{F}$ such that $T\varphi = \lambda\varphi$ for all $v \in V$.

Solution. Suppose $\dim V = 1$ and let $v \in V$ be a nonzero vector. Then $\{v\}$ is a basis of V . Let $T \in \mathcal{L}(V)$. Since $Tv \in V$, there exists $\lambda \in \mathbb{F}$ such that $Tv = \lambda v$. Now let $u \in V$. Then there exists $\alpha \in \mathbb{F}$ such that $u = \alpha v$. Therefore,

$$Tu = T(\alpha v) = \alpha Tv = \alpha(\lambda v) = \lambda(\alpha v) = \lambda u. \quad \blacksquare$$

Exercise 3A8

Give an example of a function $\varphi : \mathbb{R}^2 \rightarrow \mathbb{R}$ such that

$$\varphi(av) = a\varphi(v)$$

for all $a \in \mathbb{R}$ and all $v \in \mathbb{R}^2$ but φ is not linear.

Remark. This was a surprisingly difficult exercise for me, namely because there were no obvious examples of homogeneous functions that were not linear. However, the idea is that while x^n functions aren't linear, they don't maintain homogeneity due to an extra a^{n-1} factor. It follows that we must divide x^n by some $n - 1$ power to maintain homogeneity. However, dividing by just y risks division by zero, so we must divide by $x + y$ to avoid this.

Solution. Let $\varphi : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by

$$\varphi(x, y) = \frac{x^2}{x + y}.$$

Then for any $a \in \mathbb{R}$ and $(x, y) \in \mathbb{R}^2$,

$$\varphi(a(x, y)) = \varphi(ax, ay) = \frac{(ax)^2}{ax + ay} = \frac{a^2x^2}{a(x + y)} = a \frac{x^2}{x + y} = a\varphi(x, y).$$

However, φ is not additive. For example, let $v = (1, 0)$ and $w = (0, 1)$. Then

$$\varphi(v + w) = \varphi(1, 1) = \frac{1^2}{1 + 1} = \frac{1}{2},$$

but

$$\varphi(v) + \varphi(w) = \varphi(1, 0) + \varphi(0, 1) = \frac{1^2}{1 + 0} + \frac{0^2}{0 + 1} = 1 + 0 = 1.$$

Therefore, φ is not linear. ■

Exercise 3A9

Give an example of a function $\varphi : \mathbb{C} \rightarrow \mathbb{C}$ such that

$$\varphi(w + z) = \varphi(w) + \varphi(z)$$

for all $w, z \in \mathbb{C}$ but φ is not linear. (Here $\mathbb{C}$ is thought of as a complex vector space.)

Solution. Let $\varphi : \mathbb{C} \rightarrow \mathbb{C}$ be defined by

$$\varphi(x + iy) = x.$$

Then for any $w = x_1 + iy_1$ and $z = x_2 + iy_2$ in $\mathbb{C}$,

$$\varphi(w + z) = \varphi((x_1 + x_2) + i(y_1 + y_2)) = x_1 + x_2 = \varphi(w) + \varphi(z).$$

However, φ is not homogeneous. For example, let $w = i$ and $z = 1$. Then

$$\varphi(iz) = \varphi(i) = 0,$$

but

$$i\varphi(z) = i\varphi(1) = i(1) = i.$$

Therefore, φ is not linear. ■

Exercise 3A10

Prove or give a counterexample: If $q \in \mathcal{P}(\mathbb{R})$ and $T : \mathcal{P}(\mathbb{R}) \rightarrow \mathcal{P}(\mathbb{R})$ is defined by $Tp = q \circ p$, then T is a linear map.

Solution. This is false. Let $q(x) = x^2$ and define $T : \mathcal{P}(\mathbb{R}) \rightarrow \mathcal{P}(\mathbb{R})$ by $Tp = q \circ p$. Then for any $p_1, p_2 \in \mathcal{P}(\mathbb{R})$,

$$T(p_1 + p_2) = q \circ (p_1 + p_2) = (p_1 + p_2)^2 = p_1^2 + 2p_1p_2 + p_2^2 \neq p_1^2 + p_2^2 = Tp_1 + Tp_2. \quad \blacksquare$$

Exercise 3A11

Suppose V is finite-dimensional and $T \in \mathcal{L}(V)$. Prove that T is a scalar multiple of the identity if and only if $ST = TS$ for every $S \in \mathcal{L}(V)$.

Remark. *The backwards direction is difficult. The idea is to construct many such S that will force T to be a scalar multiple of the identity.*

Solution. Suppose T is a scalar multiple of the identity, i.e., $T = \lambda I$ for some $\lambda \in \mathbb{F}$. Then for any $S \in \mathcal{L}(V)$,

$$ST = S(\lambda I) = \lambda S = (\lambda I)S = TS.$$

Conversely, suppose $ST = TS$ for every $S \in \mathcal{L}(V)$. Let $v_1, \dots, v_n$ be a basis of V . Define $S_{ij} \in \mathcal{L}(V)$ by

$$S_{ij}v_k = \begin{cases} v_j & \text{if } k = i, \\ 0 & \text{if } k \neq i. \end{cases}$$

In other words, S_{ij} sends v_i to v_j and sends all other basis vectors to 0. Fix some i , and let $Tv_i = \alpha_1 v_1 + \cdots + \alpha_n v_n$. Then

$$S_{ij}Tv_i = S_{ij}(\alpha_1 v_1 + \cdots + \alpha_n v_n) = \alpha_i v_j,$$

and

$$TS_{ij}v_i = T(v_j) = \beta_1 v_1 + \cdots + \beta_n v_n$$

for some $\beta_1, \dots, \beta_n \in \mathbb{F}$. Since $S_{ij}T = TS_{ij}$, we have

$$\alpha_i v_j = \beta_1 v_1 + \cdots + \beta_n v_n.$$

Since $v_1, \dots, v_n$ is a basis, it follows that $\beta_k = 0$ for all $k \neq j$ and $\beta_j = \alpha_i$. Therefore, $Tv_j = \alpha_i v_j$. Since i was arbitrary, it follows that $Tv_j = \alpha_j v_j$ for some $\alpha_j \in \mathbb{F}$, hence T sends each basis vector to a scalar multiple of itself.

Now consider any $\ell \neq j$. Then

$$S_{\ell j}Tv_\ell = S_{\ell j}(\alpha_\ell v_\ell) = \alpha_\ell v_j,$$

and

$$TS_{\ell j}v_\ell = T(v_j) = \alpha_j v_j.$$

Since $S_{\ell j}T = TS_{\ell j}$, we have $\alpha_\ell v_j = \alpha_j v_j$, which implies $\alpha_\ell = \alpha_j$. Since ℓ and j were arbitrary, it follows that all α_k are equal, say $\alpha_k = \lambda$ for all k . Therefore, $Tv_k = \lambda v_k$ for all k , and hence $T = \lambda I$. ■

Exercise 3A12

Suppose U is a subspace of V with $U \neq V$. Suppose $S \in \mathcal{L}(U, W)$ and $S \neq 0$ (which means that $Su \neq 0$ for some $u \in U$). Define $T : V \rightarrow W$ by

$$Tv = \begin{cases} Sv & \text{if } v \in U, \\ 0 & \text{if } v \in V \text{ and } v \notin U. \end{cases}$$

Prove that T is not a linear map on V .

Solution. Since $U \neq V$, there exists $v \in V$ such that $v \notin U$. Let $u \in U$ be such that $Su \neq 0$. Then

$$T(u + v) = 0,$$

but

$$Tu + Tv = Su + 0 = Su \neq 0.$$

Therefore, T is not linear. ■

Exercise 3A13

Suppose V is finite-dimensional. Prove that every linear map on a subspace of V can be extended to a linear map on V . In other words, show that if U is a subspace of V and $S \in \mathcal{L}(U, W)$, then there exists $T \in \mathcal{L}(V, W)$ such that $Tu = Su$ for all $u \in U$.

Solution. Let $u_1, \dots, u_m$ be a basis of U . Since U is a subspace of V , we can extend this basis to a basis of V , say $u_1, \dots, u_m, v_1, \dots, v_n$. Define $T : V \rightarrow W$ by

$$Tu = Su \quad \text{for all } u \in U,$$

and

$$Tv_k = 0 \quad \text{for all } k = 1, \dots, n.$$

Then T is well-defined on the basis of V , and we can extend it linearly to all of V . For any $u \in U$, we may write

$$u = \alpha_1 u_1 + \dots + \alpha_m u_m$$

for some scalars $\alpha_1, \dots, \alpha_m \in \mathbb{F}$, and 0 as the coefficients for $v_1, \dots, v_n$. Then

$$Tu = \alpha_1 Tu_1 + \dots + \alpha_m Tu_m + 0 \cdot Tv_1 + \dots + 0 \cdot Tv_n = \alpha_1 Su_1 + \dots + \alpha_m Su_m = Su.$$

Therefore, T extends S to all of V . ■

Exercise 3A14

Suppose V is finite-dimensional with $\dim V > 0$, and suppose W is infinite-dimensional. Prove that $\mathcal{L}(V, W)$ is infinite-dimensional.

Solution. Let $v_1, \dots, v_n$ be a basis of V . Since W is infinite-dimensional, there exists an infinite linearly independent list of vectors $w_1, w_2, \dots$ in W . For each $k \in \mathbb{N}$, define $T_k \in \mathcal{L}(V, W)$ by

$$T_k v_j = \begin{cases} w_k & \text{if } j = 1, \\ 0 & \text{if } j \neq 1. \end{cases}$$

In other words, T_k sends the first basis vector v_1 to w_k and all other basis vectors to 0. We claim that the list $T_1, T_2, \dots$ is linearly independent in $\mathcal{L}(V, W)$. Suppose that

$$a_1 T_1 + a_2 T_2 + \dots + a_m T_m = 0$$

for some scalars $a_1, \dots, a_m \in \mathbb{F}$. Evaluating both sides at v_1 , we get

$$a_1 T_1 v_1 + a_2 T_2 v_1 + \dots + a_m T_m v_1 = a_1 w_1 + a_2 w_2 + \dots + a_m w_m = 0.$$

Since $w_1, \dots, w_m$ is linearly independent in W , it follows that $a_1 = \dots = a_m = 0$. Therefore, $T_1, T_2, \dots$ is linearly independent in $\mathcal{L}(V, W)$, and hence $\mathcal{L}(V, W)$ is infinite-dimensional. ■

Exercise 3A15

Suppose $v_1, \dots, v_m$ is a linearly dependent list of vectors in V . Suppose also that $W \neq \{0\}$. Prove that there exist $w_1, \dots, w_m \in W$ such that no $T \in \mathcal{L}(V, W)$ satisfies $Tv_k = w_k$ for each $k = 1, \dots, m$.

Solution. Since $v_1, \dots, v_m$ is linearly dependent, there exist scalars $a_1, \dots, a_m \in \mathbb{F}$, not all zero, such that

$$a_1 v_1 + \dots + a_m v_m = 0.$$

Let $w_1, \dots, w_m \in W$ be defined by

$$w_k = \begin{cases} 0 & \text{if } a_k = 0, \\ w & \text{if } a_k \neq 0, \end{cases}$$

where w is any nonzero vector in W . Suppose for contradiction that there exists $T \in \mathcal{L}(V, W)$ such that $Tv_k = w_k$ for each $k = 1, \dots, m$. Then

$$T(a_1v_1 + \dots + a_mv_m) = a_1Tv_1 + \dots + a_mTv_m = a_1w_1 + \dots + a_mw_m.$$

Since $a_1v_1 + \dots + a_mv_m = 0$, we have $T(0) = 0$, so

$$0 = a_1w_1 + \dots + a_mw_m.$$

However, by the definition of w_k , we have $a_kw_k = a_kw$ for all k such that $a_k \neq 0$, and $a_kw_k = 0$ for all k such that $a_k = 0$. Therefore, the sum $a_1w_1 + \dots + a_mw_m$ is a nonzero scalar multiple of w , which is nonzero since $w \neq 0$. This is a contradiction. Therefore, no such $T \in \mathcal{L}(V, W)$ exists. ■

Exercise 3A16

Suppose V is finite-dimensional with $\dim V > 1$. Prove that there exist $S, T \in \mathcal{L}(V)$ such that $ST \neq TS$.

Solution. Let v_1, v_2 be linearly independent vectors in V . We may extend this list to a basis of V , say $v_1, v_2, \dots, v_n$. Define $S, T \in \mathcal{L}(V)$ by

$$Sv_1 = v_2, \quad Sv_2 = v_1, \quad Sv_k = 0 \text{ for } k > 2,$$

and

$$Tv_1 = v_1, \quad Tv_2 = 0, \quad Tv_k = 0 \text{ for } k > 2.$$

Then

$$STv_1 = S(Tv_1) = Sv_1 = v_2,$$

but

$$TSv_1 = T(Sv_1) = Tv_2 = 0.$$

Therefore, $ST \neq TS$. ■

Exercise 3A17

Suppose V is finite-dimensional. Show that the only two-sided ideals of $\mathcal{L}(V)$ are $\{0\}$ and $\mathcal{L}(V)$.

Note. A subspace $\mathcal{E}$ of $\mathcal{L}(V)$ is called a *two-sided ideal* of $\mathcal{L}(V)$ if $TE \in \mathcal{E}$ and $ET \in \mathcal{E}$ for all $E \in \mathcal{E}$ and all $T \in \mathcal{L}(V)$.

Solution. Let $\mathcal{E}$ be a two-sided ideal of $\mathcal{L}(V)$. If $\mathcal{E} = \{0\}$, then we are done. Otherwise, there exists $E \in \mathcal{E}$ such that $E \neq 0$. Then there exists $v_1 \in V$ such that $Ev_1 \neq 0$. Since V is finite-dimensional, we may extend the list $\{v_1\}$ to a basis of V , say $v_1, v_2, \dots, v_n$. Let $w_1 = Ev_1 \neq 0$. We may also extend the list $\{w_1\}$ to a basis of V , say $w_1, w_2, \dots, w_n$. Define $T_1 \in \mathcal{L}(V)$ by

$$T_1w_1 = v_1, \quad T_1w_k = 0 \text{ for } k > 1.$$

Then $T_1E \in \mathcal{E}$ since $\mathcal{E}$ is a two-sided ideal. Observe that $T_1Ev_1 = T_1(Ev_1) = T_1w_1 = v_1 \neq 0$. Therefore, $T_1E \neq 0$. Consider $S_1 \in \mathcal{L}(V)$ defined by

$$S_1v_1 = v_1, \quad S_1v_k = 0 \text{ for } k > 1.$$

Then $T_1ES_1 \in \mathcal{E}$ since $\mathcal{E}$ is a two-sided ideal. Observe that $T_1ES_1v_1 = T_1Ev_1 = T_1w_1 = v_1 \neq 0$. Therefore, $T_1ES_1 \neq 0$. We now discuss the construction of the identity map $I \in \mathcal{L}(V)$.

Construct $T_i, S_i \in \mathcal{L}(V)$ for each $i = 1, \dots, n$ as follows:

$$T_iw_1 = v_i, \quad T_iw_k = 0 \text{ for } k \neq 1,$$

and

$$S_iv_i = v_1, \quad S_iv_k = 0 \text{ for } k \neq i.$$

Then $T_iES_i \in \mathcal{E}$ and $T_iES_iv_i = v_i \neq 0$. Since $\mathcal{E}$ is a subspace, it contains all linear combinations of these maps. In particular, for any $v \in V$, we can write $v = \alpha_1v_1 + \alpha_2v_2 + \dots + \alpha_nv_n$ for some scalars $\alpha_1, \alpha_2, \dots, \alpha_n \in \mathbb{F}$. Observe that

$$T_iES_iv = T_iES_i(\alpha_1v_1) + \dots + T_iES_i(\alpha_nv_n) = \alpha_iv_i.$$

Consider the map

$$E^* = T_1ES_1 + T_2ES_2 + \dots + T_nES_n \in \mathcal{E},$$

where $E^* \in \mathcal{E}$ since $\mathcal{E}$ is a subspace. Then

$$E^*v = T_1ES_1v + T_2ES_2v + \dots + T_nES_nv = \alpha_1v_1 + \alpha_2v_2 + \dots + \alpha_nv_n = v.$$

Therefore, for any $v \in V$, there exists $E^* \in \mathcal{E}$ such that $E^*v = v$. Now let $T \in \mathcal{L}(V)$ be arbitrary. Then

$$T = TE^* \in \mathcal{E}$$

since $TE^* \in \mathcal{E}$ and $\mathcal{E}$ is a two-sided ideal. Therefore, $\mathcal{E} = \mathcal{L}(V)$. We conclude that the only two-sided ideals of $\mathcal{L}(V)$ are $\{0\}$ and $\mathcal{L}(V)$. ■

3B Null Spaces and Ranges

Exercise 3B1

Give an example of a linear map T with $\dim \text{null } T = 3$ and $\dim \text{range } T = 2$.

Solution. Let $T \in \mathcal{L}(\mathbb{R}^5, \mathbb{R}^2)$ be defined by

$$T(x_1, x_2, x_3, x_4, x_5) = (x_1, x_2).$$

Then $\text{null } T = \{(0, 0, x_3, x_4, x_5) \mid x_3, x_4, x_5 \in \mathbb{R}\}$, which has dimension 3, and $\text{range } T = \mathbb{R}^2$, which has dimension 2. ■

Exercise 3B2

Suppose $S, T \in \mathcal{L}(V)$ are such that $\text{range } S \subseteq \text{null } T$. Prove that $(ST)^2 = 0$.

Solution. Let $v \in V$. Then

$$(ST)^2 v = ST(STv) = S(T(STv)).$$

Since $STv \in \text{range } S$ and $\text{range } S \subseteq \text{null } T$, we have $T(STv) = 0$. Therefore,

$$(ST)^2 v = S(0) = 0.$$

Since v was arbitrary, it follows that $(ST)^2 = 0$. ■

Exercise 3B3

Suppose $v_1, \dots, v_m$ is a list of vectors in V . Define $T \in \mathcal{L}(\mathbb{F}^m, V)$ by

$$T(z_1, \dots, z_m) = z_1 v_1 + \dots + z_m v_m.$$

- (a) What property of T corresponds to $v_1, \dots, v_m$ spanning V ?
- (b) What property of T corresponds to the list $v_1, \dots, v_m$ being linearly independent?

Solution.

- (a) The list $v_1, \dots, v_m$ spans V if and only if $\text{range } T = V$, i.e., T is surjective.
- (b) The list $v_1, \dots, v_m$ is linearly independent if and only if $\text{null } T = \{0\}$, i.e., T is injective. ■

Exercise 3B4

Show that $\{T \in \mathcal{L}(\mathbb{R}^5, \mathbb{R}^4) \mid \dim \text{null } T > 2\}$ is not a subspace of $\mathcal{L}(\mathbb{R}^5, \mathbb{R}^4)$.

Solution. Let $T_1, T_2 \in \mathcal{L}(\mathbb{R}^5, \mathbb{R}^4)$ be defined by

$$T_1(x_1, x_2, x_3, x_4, x_5) = (x_1, x_2, 0, 0)$$

and

$$T_2(x_1, x_2, x_3, x_4, x_5) = (0, 0, x_3, x_4).$$

Then $\text{null } T_1 = \{(0, 0, x_3, x_4, x_5) \mid x_3, x_4, x_5 \in \mathbb{R}\}$ and $\text{null } T_2 = \{(x_1, x_2, 0, 0, x_5) \mid x_1, x_2, x_5 \in \mathbb{R}\}$ both have dimension 3, so T_1 and T_2 are in the set $\{T \in \mathcal{L}(\mathbb{R}^5, \mathbb{R}^4) \mid \dim \text{null } T > 2\}$. However,

$$(T_1 + T_2)(x_1, x_2, x_3, x_4, x_5) = (x_1, x_2, x_3, x_4),$$

which has null space $\text{null}(T_1 + T_2) = \{(0, 0, 0, 0, x_5) \mid x_5 \in \mathbb{R}\}$ of dimension 1. Therefore,

$$\dim \text{null}(T_1 + T_2) = 1 < 2,$$

so $T_1 + T_2$ is not in the set. Hence, the set is not closed under addition and is not a subspace of $\mathcal{L}(\mathbb{R}^5, \mathbb{R}^4)$. ■

Exercise 3B5

Give an example of $T \in \mathcal{L}(\mathbb{R}^4)$ such that $\text{range } T = \text{null } T$.

Solution. Let v_1, v_2, v_3, v_4 be the standard basis of $\mathbb{R}^4$. Define $T \in \mathcal{L}(\mathbb{R}^4)$ by

$$Tv_1 = v_3, \quad Tv_2 = v_4, \quad Tv_3 = 0, \quad Tv_4 = 0.$$

Then for any $x = (x_1, x_2, x_3, x_4) \in \mathbb{R}^4$,

$$Tx = x_1Tv_1 + x_2Tv_2 + x_3Tv_3 + x_4Tv_4 = x_1v_3 + x_2v_4.$$

Then $\text{range } T = \text{span } v_3, v_4$. If $Tx = 0$, then $x_1v_3 + x_2v_4 = 0$, which implies $x_1 = x_2 = 0$. Therefore, $\text{null } T = \{(0, 0, x_3, x_4) \mid x_3, x_4 \in \mathbb{R}\} = \text{span } v_3, v_4$. Hence, $\text{range } T = \text{null } T$. ■

Exercise 3B6

Prove that there does not exist $T \in \mathcal{L}(\mathbb{R}^5)$ such that $\text{range } T = \text{null } T$.

Solution. Suppose for contradiction that there exists $T \in \mathcal{L}(\mathbb{R}^5)$ such that $\text{range } T = \text{null } T$. Then by the rank-nullity theorem,

$$\dim \mathbb{R}^5 = \dim \text{null } T + \dim \text{range } T = 2 \dim \text{range } T.$$

Therefore, $5 = 2 \dim \text{range } T$, which is impossible since 5 is odd and $2 \dim \text{range } T$ is even. Hence, no such T exists. ■

Exercise 3B7

Suppose V and W are finite-dimensional with $2 \leq \dim V \leq \dim W$. Show that $\{T \in \mathcal{L}(V, W) \mid T \text{ is not injective}\}$ is not a subspace of $\mathcal{L}(V, W)$.

Solution. Let $\dim V = n$ and $\dim W = m$ with $2 \leq n \leq m$. Let $v_1, v_2, \dots, v_n$ be a basis of V , and let $w_1, w_2, \dots, w_m$ be a basis of W . Define $T_1, T_2 \in \mathcal{L}(V, W)$ by

$$T_1 v_1 = w_1, \quad T_1 v_k = 0 \text{ for } k \geq 2,$$

and

$$T_2 v_1 = 0, \quad T_2 v_k = w_k \text{ for } k \geq 2.$$

Then $v_2 \in \text{null } T_1$ and $v_1 \in \text{null } T_2$, so both T_1 and T_2 are not injective. However,

$$(T_1 + T_2)v_1 = T_1 v_1 + T_2 v_1 = w_1 + 0 = w_1 \neq 0,$$

and for $k \geq 2$,

$$(T_1 + T_2)v_k = T_1 v_k + T_2 v_k = 0 + w_k = w_k \neq 0.$$

Suppose $v \in V$ such that $(T_1 + T_2)v = 0$. Then v can be expressed as a linear combination of the basis vectors:

$$v = \alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_n v_n.$$

Then

$$(T_1 + T_2)v = \alpha_1 w_1 + \alpha_2 w_2 + \dots + \alpha_n w_n = 0.$$

Since $w_1, w_2, \dots, w_m$ are linearly independent, it follows that $\alpha_1 = \alpha_2 = \dots = \alpha_n = 0$, hence $v = 0$. Therefore, $T_1 + T_2$ is injective. Since T_1 and T_2 are not injective, but their sum is injective, the set $\{T \in \mathcal{L}(V, W) \mid T \text{ is not injective}\}$ is not closed under addition and thus is not a subspace of $\mathcal{L}(V, W)$. ■

Exercise 3B8

Suppose V and W are finite-dimensional with $\dim V \geq \dim W \geq 2$. Show that $\{T \in \mathcal{L}(V, W) \mid T \text{ is not surjective}\}$ is not a subspace of $\mathcal{L}(V, W)$.

Solution. We proceed similarly to the previous exercise. Let $\dim V = n$ and $\dim W = m$ with $n \geq m \geq 2$. Let $v_1, v_2, \dots, v_n$ be a basis of V , and let $w_1, w_2, \dots, w_m$ be a basis of W . Define $T_1, T_2 \in \mathcal{L}(V, W)$ by

$$T_1 v_1 = w_1, \quad T_1 v_k = 0 \text{ for } k \geq 2,$$

and

$$T_2 v_1 = 0, \quad T_2 v_k = w_k \text{ for } k = 2, \dots, m, \quad T_2 v_k = 0 \text{ for } k > m.$$

Then $\text{range } T_1 = \text{span } w_1$ and $\text{range } T_2 = \text{span } w_2, \dots, w_m$, both of which are proper subspaces of W , so T_1 and T_2 are not surjective. However,

$$(T_1 + T_2)v_1 = T_1 v_1 + T_2 v_1 = w_1 + 0 = w_1,$$

and for $k = 2, \dots, m$,

$$(T_1 + T_2)v_k = T_1 v_k + T_2 v_k = 0 + w_k = w_k.$$

Therefore, $\text{range}(T_1 + T_2) = \text{span } w_1, w_2, \dots, w_m = W$, so $T_1 + T_2$ is surjective. Since T_1 and T_2 are not surjective, but their sum is surjective, the set $\{T \in \mathcal{L}(V, W) \mid T \text{ is not surjective}\}$ is not closed under addition and thus is not a subspace of $\mathcal{L}(V, W)$. ■

Exercise 3B9

Suppose $T \in \mathcal{L}(V, W)$ is injective and $v_1, \dots, v_n$ is linearly independent in V . Prove that $Tv_1, \dots, Tv_n$ is linearly independent in W .

Solution. Suppose $a_1, \dots, a_n \in \mathbb{F}$ are such that

$$a_1Tv_1 + \dots + a_nTv_n = 0.$$

By linearity of T , we have

$$T(a_1v_1 + \dots + a_nv_n) = 0.$$

Injectivity of T implies that $\text{null } T = \{0\}$, so

$$a_1v_1 + \dots + a_nv_n = 0.$$

Since $v_1, \dots, v_n$ is linearly independent, it follows that $a_1 = \dots = a_n = 0$. Therefore, $Tv_1, \dots, Tv_n$ is linearly independent in W . ■

Exercise 3B10

Suppose $v_1, \dots, v_n$ spans V and $T \in \mathcal{L}(V, W)$. Show that $Tv_1, \dots, Tv_n$ spans $\text{range } T$.

Solution. Let $w \in \text{range } T$. Then there exists $v \in V$ such that $Tv = w$. Since $v_1, \dots, v_n$ spans V , we can write

$$v = a_1v_1 + \dots + a_nv_n$$

for some scalars $a_1, \dots, a_n \in \mathbb{F}$. By linearity of T , we have

$$w = Tv = T(a_1v_1 + \dots + a_nv_n) = a_1Tv_1 + \dots + a_nTv_n.$$

Therefore, w is a linear combination of $Tv_1, \dots, Tv_n$. Since w was arbitrary in $\text{range } T$, it follows that $Tv_1, \dots, Tv_n$ spans $\text{range } T$. ■

Exercise 3B11

Suppose that V is finite-dimensional and that $T \in \mathcal{L}(V, W)$. Prove that there exists a subspace U of V such that

$$U \cap \text{null } T = \{0\} \quad \text{and} \quad \text{range } T = \{Tu \mid u \in U\}.$$

Remark. Note that this exercise proves the existence of a subspace U of V such that $V = U \oplus \text{null } T$.

Solution. Since $\text{null } T$ is a subspace of V , we can extend a basis of $\text{null } T$ to a basis of V . Let $n_1, \dots, n_k$ be a basis of $\text{null } T$, and extend this to a basis of V by adding vectors $u_1, \dots, u_m$. Hence, a basis for V is given by

$$n_1, \dots, n_k, u_1, \dots, u_m.$$

Define $U = \text{span}(u_1, \dots, u_m)$. Suppose $v \in U \cap \text{null } T$. Then v can be expressed as a linear combination of the basis vectors of U :

$$v = \alpha_1u_1 + \dots + \alpha_mu_m.$$

Since $v \in \text{null } T$ as well, we can also express v as a linear combination of the basis vectors of $\text{null } T$:

$$v = \beta_1 n_1 + \cdots + \beta_k n_k.$$

Equating the two expressions for v , we have

$$\alpha_1 u_1 + \cdots + \alpha_m u_m = \beta_1 n_1 + \cdots + \beta_k n_k.$$

Since $n_1, \dots, n_k, u_1, \dots, u_m$ is a basis for V , the only way this equality can hold is if all coefficients are zero, i.e., $\alpha_1 = \cdots = \alpha_m = 0$ and $\beta_1 = \cdots = \beta_k = 0$. Therefore, $v = 0$, and we conclude that $U \cap \text{null } T = \{0\}$.

Now, let $w \in \text{range } T$. Then there exists $v \in V$ such that $Tv = w$. We can express v as a linear combination of the basis vectors of V :

$$v = \gamma_1 n_1 + \cdots + \gamma_k n_k + \delta_1 u_1 + \cdots + \delta_m u_m.$$

Applying T to both sides, we have

$$Tv = T(\gamma_1 n_1 + \cdots + \gamma_k n_k + \delta_1 u_1 + \cdots + \delta_m u_m) = \gamma_1 Tn_1 + \cdots + \gamma_k Tn_k + \delta_1 Tu_1 + \cdots + \delta_m Tu_m.$$

Since $n_1, \dots, n_k \in \text{null } T$, we have $Tn_i = 0$ for all $i = 1, \dots, k$. Therefore,

$$Tv = \delta_1 Tu_1 + \cdots + \delta_m Tu_m.$$

Let $u = \delta_1 u_1 + \cdots + \delta_m u_m \in U$. Then $Tu = w$. Since w was arbitrary in $\text{range } T$, it follows that $\text{range } T = \{Tu \mid u \in U\}$. Thus, we have found a subspace U of V such that $U \cap \text{null } T = \{0\}$ and $\text{range } T = \{Tu \mid u \in U\}$. ■

Exercise 3B12

Suppose T is a linear map from $\mathbb{F}^4$ to $\mathbb{F}^2$ such that

$$\text{null } T = \{(x_1, x_2, x_3, x_4) \in \mathbb{F}^4 \mid x_1 = 5x_2 \text{ and } x_3 = 7x_4\}.$$

Prove that T is surjective.

Solution. We may write the null space of T as

$$\text{null } T = \{(5x_2, x_2, 7x_4, x_4) \mid x_2, x_4 \in \mathbb{F}\}.$$

From this, we can see that $\text{null } T$ is spanned by the vectors $(5, 1, 0, 0)$ and $(0, 0, 7, 1)$. Therefore, $\dim \text{null } T = 2$. By the rank-nullity theorem, we have

$$\dim \mathbb{F}^4 = \dim \text{null } T + \dim \text{range } T \implies 4 = 2 + \dim \text{range } T.$$

Solving for $\dim \text{range } T$, we find that $\dim \text{range } T = 2$. Since $\dim \mathbb{F}^2 = 2$, it follows that $\text{range } T = \mathbb{F}^2$. Therefore, T is surjective. ■

Exercise 3B13

Suppose U is a three-dimensional subspace of $\mathbb{R}^8$ and that T is a linear map from $\mathbb{R}^8$ to $\mathbb{R}^5$ such that $\text{null } T = U$. Prove that T is surjective.

Solution. By the rank-nullity theorem, we have

$$\dim \mathbb{R}^8 = \dim \text{null } T + \dim \text{range } T \implies 8 = 3 + \dim \text{range } T.$$

Solving for $\dim \text{range } T$, we find that $\dim \text{range } T = 5$. Since $\dim \mathbb{R}^5 = 5$, it follows that $\text{range } T = \mathbb{R}^5$. Therefore, T is surjective. ■

Exercise 3B14

Prove that there does not exist a linear map from $\mathbb{F}^5$ to $\mathbb{F}^2$ whose null space equals $\{(x_1, x_2, x_3, x_4, x_5) \in \mathbb{F}^5 \mid x_1 = 3x_2 \text{ and } x_3 = x_4 = x_5\}$.

Solution. We can express the null space as

$$\text{null } T = \{(3x_2, x_2, x_4, x_4, x_4) \mid x_2, x_4 \in \mathbb{F}\}.$$

From this, we can see that $\text{null } T$ is spanned by the vectors $(3, 1, 0, 0, 0)$ and $(0, 0, 1, 1, 1)$. Therefore, $\dim \text{null } T = 2$. By the rank-nullity theorem, we have

$$\dim \mathbb{F}^5 = \dim \text{null } T + \dim \text{range } T \implies 5 = 2 + \dim \text{range } T.$$

Solving for $\dim \text{range } T$, we find that $\dim \text{range } T = 3$. However, $\dim \mathbb{F}^2 = 2$, which is less than 3. This is a contradiction, as the dimension of the range cannot exceed the dimension of the target space. Therefore, no such linear map exists. ■

Exercise 3B15

Suppose there exists a linear map on V whose null space and range are both finite-dimensional. Prove that V is finite-dimensional.

Solution. Let $T \in \mathcal{L}(V)$ be such that $\text{null } T$ and $\text{range } T$ are both finite-dimensional. By the rank-nullity theorem, we have

$$\dim V = \dim \text{null } T + \dim \text{range } T.$$

Since both $\dim \text{null } T$ and $\dim \text{range } T$ are finite, their sum is also finite. Therefore, $\dim V$ is finite, which implies that V is finite-dimensional. ■

Exercise 3B16

Suppose V and W are both finite-dimensional. Prove that there exists an injective linear map from V to W if and only if $\dim V \leq \dim W$.

Solution. Suppose there exists an injective linear map $T \in \mathcal{L}(V, W)$. By the rank-nullity theorem, we have

$$\dim V = \dim \text{null } T + \dim \text{range } T.$$

Since T is injective, $\text{null } T = \{0\}$, so $\dim \text{null } T = 0$. Therefore,

$$\dim V = 0 + \dim \text{range } T = \dim \text{range } T.$$

Since $\text{range } T$ is a subspace of W , we have $\dim \text{range } T \leq \dim W$. Thus,

$$\dim V = \dim \text{range } T \leq \dim W.$$

Conversely, suppose $\dim V \leq \dim W$. Let $n = \dim V$ and $m = \dim W$. Choose a basis $\{v_1, \dots, v_n\}$ of V . Let $\{w_1, \dots, w_m\}$ be a basis of W . Define a linear map $T \in \mathcal{L}(V, W)$ by

$$Tv_i = w_i \quad \text{for } i = 1, \dots, n.$$

Since $n \leq m$, we can assign each basis vector of V to a distinct basis vector of W . To show that T is injective, let $v = \alpha_1 v_1 + \dots + \alpha_n v_n \in V$ such that $Tv = 0$. Then

$$Tv = \alpha_1 Tv_1 + \dots + \alpha_n Tv_n = \alpha_1 w_1 + \dots + \alpha_n w_n = 0.$$

Since $w_1, \dots, w_n$ are linearly independent, it follows that $\alpha_1 = \dots = \alpha_n = 0$. Therefore, $v = 0$, and T is injective. Hence, there exists an injective linear map from V to W if and only if $\dim V \leq \dim W$. ■

Exercise 3B17

Suppose V and W are both finite-dimensional. Prove that there exists a surjective linear map from V onto W if and only if $\dim V \geq \dim W$.

Solution. Suppose there exists a surjective linear map $T \in \mathcal{L}(V, W)$. By the rank-nullity theorem, we have

$$\dim V = \dim \text{null } T + \dim \text{range } T.$$

Since T is surjective, $\text{range } T = W$, so $\dim \text{range } T = \dim W$. Therefore,

$$\dim V = \dim \text{null } T + \dim W.$$

Since $\dim \text{null } T \geq 0$, it follows that

$$\dim V \geq \dim W.$$

Conversely, suppose $\dim V \geq \dim W$. Let $n = \dim V$ and $m = \dim W$. Choose a basis $\{v_1, \dots, v_n\}$ of V . Let $\{w_1, \dots, w_m\}$ be a basis of W . Define a linear map $T \in \mathcal{L}(V, W)$ by

$$Tv_i = w_i \quad \text{for } i = 1, \dots, m, \quad Tv_i = 0 \quad \text{for } i = m+1, \dots, n.$$

To show that T is surjective, let $w \in W$. Then w can be expressed as a linear combination of the basis vectors of W :

$$w = \beta_1 w_1 + \dots + \beta_m w_m.$$

Consider the vector $v = \beta_1 v_1 + \dots + \beta_m v_m \in V$. Then

$$Tv = \beta_1 Tv_1 + \dots + \beta_m Tv_m = \beta_1 w_1 + \dots + \beta_m w_m = w.$$

Since w was arbitrary in W , it follows that $\text{range } T = W$. Therefore, T is surjective. Hence, there exists a surjective linear map from V onto W if and only if $\dim V \geq \dim W$. ■

Exercise 3B18

Suppose V and W are finite-dimensional and that U is a subspace of V . Prove that there exists $T \in \mathcal{L}(V, W)$ such that $\text{null } T = U$ if and only if $\dim U \geq \dim V - \dim W$.

Solution. Suppose there exists $T \in \mathcal{L}(V, W)$ such that $\text{null } T = U$. By the rank-nullity theorem, we have

$$\dim V = \dim \text{null } T + \dim \text{range } T = \dim U + \dim \text{range } T.$$

Since $\text{range } T$ is a subspace of W , we have $\dim \text{range } T \leq \dim W$. Therefore,

$$\dim V = \dim U + \dim \text{range } T \leq \dim U + \dim W.$$

Rearranging this inequality gives

$$\dim U \geq \dim V - \dim W.$$

Conversely, suppose $\dim U \geq \dim V - \dim W$. Let $n = \dim V$, $m = \dim W$, and $k = \dim U$. Let $u_1, \dots, u_k$ be a basis of U . Extend this basis to a basis of V by adding vectors $v_1, \dots, v_{n-k}$. Define a linear map $T \in \mathcal{L}(V, W)$ by

$$Tu_i = 0 \quad \text{for } i = 1, \dots, k, \quad Tv_j = w_j \quad \text{for } j = 1, \dots, n-k,$$

where $w_1, \dots, w_{n-k}$ are linearly independent vectors in W . Since $\dim U = k$ and $\dim V - \dim W = n - m$, we have $k \geq n - m$, which implies that $n - k \leq m$. Therefore, we can choose $w_1, \dots, w_{n-k}$ to be linearly independent in W . To show that $\text{null } T = U$, let $v \in \text{null } T$. Then $Tv = 0$. Express v as a linear combination of the basis vectors of V :

$$v = \alpha_1 u_1 + \dots + \alpha_k u_k + \beta_1 v_1 + \dots + \beta_{n-k} v_{n-k}.$$

Applying T to both sides, we have

$$Tv = \alpha_1 Tu_1 + \dots + \alpha_k Tu_k + \beta_1 Tv_1 + \dots + \beta_{n-k} Tv_{n-k} = 0 + \beta_1 w_1 + \dots + \beta_{n-k} w_{n-k} = 0.$$

Since $w_1, \dots, w_{n-k}$ are linearly independent, it follows that $\beta_1 = \dots = \beta_{n-k} = 0$. Therefore,

$$v = \alpha_1 u_1 + \dots + \alpha_k u_k \in U.$$

Hence, $\text{null } T \subseteq U$. Conversely, if $v \in U$, then $Tv = 0$ by the definition of T . Therefore, $U \subseteq \text{null } T$. Combining these two inclusions, we have $\text{null } T = U$. Thus, there exists $T \in \mathcal{L}(V, W)$ such that $\text{null } T = U$ if and only if $\dim U \geq \dim V - \dim W$. ■

Exercise 3B19

Suppose W is finite-dimensional and $T \in \mathcal{L}(V, W)$. Prove that T is injective if and only if there exists $S \in \mathcal{L}(W, V)$ such that ST is the identity operator on V .

Solution. Suppose T is injective. Then $\text{null } T = \{0\}$. Let $n = \dim V$ and $m = \dim W$. Since T is injective, we have $\dim \text{range } T = n$. Choose a basis $\{v_1, \dots, v_n\}$ of V . Then $\{Tv_1, \dots, Tv_n\}$ is linearly independent in W . Extend this set to a basis of W by adding vectors $w_1, \dots, w_{m-n}$. Define a linear map $S \in \mathcal{L}(W, V)$ by

$$S(Tv_i) = v_i \quad \text{for } i = 1, \dots, n, \quad S(w_j) = 0 \quad \text{for } j = 1, \dots, m-n.$$

Then for any $v \in V$, we can express v as a linear combination of the basis vectors:

$$v = \alpha_1 v_1 + \dots + \alpha_n v_n.$$

Applying T to both sides, we have

$$Tv = \alpha_1 Tv_1 + \cdots + \alpha_n Tv_n.$$

Now, applying S to Tv , we get

$$STv = S(\alpha_1 Tv_1 + \cdots + \alpha_n Tv_n) = \alpha_1 S(Tv_1) + \cdots + \alpha_n S(Tv_n) = \alpha_1 v_1 + \cdots + \alpha_n v_n = v.$$

Therefore, ST is the identity operator on V .

Conversely, suppose there exists $S \in \mathcal{L}(W, V)$ such that ST is the identity operator on V . Let $v \in \text{null } T$. Then $Tv = 0$, and applying S gives

$$STv = S(0) = 0.$$

Since ST is the identity operator on V , we have $v = STv = 0$. Therefore, $\text{null } T = \{0\}$, which means T is injective. ■

Exercise 3B20

Suppose W is finite-dimensional and $T \in \mathcal{L}(V, W)$. Prove that T is surjective if and only if there exists $S \in \mathcal{L}(W, V)$ such that TS is the identity operator on W .

Solution. Suppose T is surjective. Then $\text{range } T = W$. Let $m = \dim W$ and choose a basis $\{w_1, \dots, w_m\}$ of W . Since T is surjective, for each w_i , there exists $v_i \in V$ such that $Tv_i = w_i$. Define a linear map $S \in \mathcal{L}(W, V)$ by

$$S(w_i) = v_i \quad \text{for } i = 1, \dots, m.$$

Then for any $w \in W$, we can express w as a linear combination of the basis vectors:

$$w = \alpha_1 w_1 + \cdots + \alpha_m w_m.$$

Applying S to both sides, we have

$$Sw = S(\alpha_1 w_1 + \cdots + \alpha_m w_m) = \alpha_1 S(w_1) + \cdots + \alpha_m S(w_m) = \alpha_1 v_1 + \cdots + \alpha_m v_m.$$

Now, applying T to Sw , we get

$$TSw = T(\alpha_1 v_1 + \cdots + \alpha_m v_m) = \alpha_1 Tv_1 + \cdots + \alpha_m Tv_m = \alpha_1 w_1 + \cdots + \alpha_m w_m = w.$$

Therefore, TS is the identity operator on W .

Conversely, suppose there exists $S \in \mathcal{L}(W, V)$ such that TS is the identity operator on W . Let $w \in W$. Then

$$TSw = w.$$

This means that for every $w \in W$, there exists a vector $v = Sw \in V$ such that $Tv = T(Sw) = w$. Therefore, $\text{range } T = W$, which means that T is surjective. ■

Exercise 3B21

Suppose V is finite-dimensional, $T \in \mathcal{L}(V, W)$, and U is a subspace of W . Prove that $\{v \in V \mid Tv \in U\}$ is a subspace of V and

$$\dim\{v \in V \mid Tv \in U\} = \dim \text{null } T + \dim(U \cap \text{range } T).$$

Solution. Let $X = \{v \in V \mid Tv \in U\}$. We first show that X is a subspace of V . Let $v_1, v_2 \in X$ and $\alpha, \beta \in \mathbb{F}$. Then

$$T(\alpha v_1 + \beta v_2) = \alpha T v_1 + \beta T v_2.$$

Since $T v_1, T v_2 \in U$ and U is a subspace of W , it follows that $\alpha T v_1 + \beta T v_2 \in U$. Therefore, $\alpha v_1 + \beta v_2 \in X$, and thus X is a subspace of V .

Now, we will compute the dimension of X . Let $Y = U \cap \text{range } T$. Let $\dim T = k$ and $\dim Y = m$. Let $n_1, \dots, n_k$ be a basis of $\text{null } T$, and let $y_1, \dots, y_m$ be a basis of Y . Since $y_i \in \text{range } T$, there exist vectors $v_i \in V$ such that $T v_i = y_i$ for $i = 1, \dots, m$. We claim that the set $\{n_1, \dots, n_k, v_1, \dots, v_m\}$ is a basis of X .

First, we show that this set spans X . Let $v \in X$. Then $T v \in U$, and since $T v \in \text{range } T$, we have $T v \in Y$. Therefore, there exist scalars $\alpha_1, \dots, \alpha_m \in \mathbb{F}$ such that

$$T v = \alpha_1 y_1 + \dots + \alpha_m y_m = \alpha_1 T v_1 + \dots + \alpha_m T v_m.$$

Let $v' = v - (\alpha_1 v_1 + \dots + \alpha_m v_m)$. Then

$$T v' = T v - T(\alpha_1 v_1 + \dots + \alpha_m v_m) = T v - (\alpha_1 T v_1 + \dots + \alpha_m T v_m) = 0.$$

Therefore, $v' \in \text{null } T$, and we can express v' as a linear combination of the basis vectors of $\text{null } T$:

$$v' = \beta_1 n_1 + \dots + \beta_k n_k$$

for some scalars $\beta_1, \dots, \beta_k \in \mathbb{F}$. Thus,

$$v = v' + (\alpha_1 v_1 + \dots + \alpha_m v_m) = \beta_1 n_1 + \dots + \beta_k n_k + \alpha_1 v_1 + \dots + \alpha_m v_m.$$

This shows that v is a linear combination of the vectors in $\{n_1, \dots, n_k, v_1, \dots, v_m\}$, and hence this set spans X .

Next, we show that this set is linearly independent. Suppose

$$\gamma_1 n_1 + \dots + \gamma_k n_k + \delta_1 v_1 + \dots + \delta_m v_m = 0$$

for some scalars $\gamma_1, \dots, \gamma_k, \delta_1, \dots, \delta_m \in \mathbb{F}$. Observe that $T n_i = 0$ for all $i = 1, \dots, k$, and $T v_j = y_j$ for all $j = 1, \dots, m$. Therefore,

$$0 = T(\gamma_1 n_1 + \dots + \gamma_k n_k + \delta_1 v_1 + \dots + \delta_m v_m) = \delta_1 y_1 + \dots + \delta_m y_m.$$

Since $y_1, \dots, y_m$ are linearly independent, it follows that $\delta_1 = \dots = \delta_m = 0$. Therefore,

$$\gamma_1 n_1 + \dots + \gamma_k n_k = 0.$$

Since $n_1, \dots, n_k$ are linearly independent, then $\gamma_1 = \dots = \gamma_k = 0$. Thus, the set $\{n_1, \dots, n_k, v_1, \dots, v_m\}$ is linearly independent and a basis of X . Therefore,

$$\dim X = k + m = \dim \text{null } T + \dim(U \cap \text{range } T).$$

■

Exercise 3B22

Suppose U and V are finite-dimensional vector spaces and $S \in \mathcal{L}(V, W)$ and $T \in \mathcal{L}(U, V)$. Prove that

$$\dim \text{null } ST \leq \dim \text{null } S + \dim \text{null } T.$$

Solution. Let $u \in U$ such that $u \in \text{null } ST$. Then $STu = 0$, which implies that $Tu \in \text{null } S$. Therefore, we can express $\text{null } ST$ as

$$\text{null } ST = \{u \in U \mid Tu \in \text{null } S\}.$$

Observe that this looks similar to the set in Exercise 3b21, where T is a linear map from U to V and $\text{null } S$ is a subspace of V . By applying the result from Exercise 3b21, we have

$$\dim \text{null } ST = \dim \text{null } T + \dim(\text{null } S \cap \text{range } T).$$

Since $\text{null } S \cap \text{range } T$ is a subspace of $\text{null } S$, we have

$$\dim(\text{null } S \cap \text{range } T) \leq \dim \text{null } S.$$

Therefore,

$$\dim \text{null } ST = \dim \text{null } T + \dim(\text{null } S \cap \text{range } T) \leq \dim \text{null } T + \dim \text{null } S. \quad \blacksquare$$

Exercise 3B23

Suppose U and V are finite-dimensional vector spaces and $S \in \mathcal{L}(V, W)$ and $T \in \mathcal{L}(U, V)$. Prove that

$$\dim \text{range } ST \leq \min(\dim \text{range } S, \dim \text{range } T).$$

Solution. Let $u \in U$ such that $STu \in \text{range } ST$. Then $STu = S(Tu)$, which implies that $Tu \in \text{range } T$. Therefore, we can express $\text{range } ST$ as

$$\text{range } ST = \{S(Tu) \mid u \in U\}.$$

Since $Tu \in \text{range } T$, we can rewrite this as

$$\text{range } ST = \{Sv \mid v \in \text{range } T\}.$$

This shows that $\text{range } ST$ is a subset of $\text{range } S$, and hence

$$\dim \text{range } ST \leq \dim \text{range } S.$$

Now let $v_1, \dots, v_m$ be a basis of $\text{range } T$. Then for any $u \in U$, we can express Tu as a linear combination of the basis vectors:

$$Tu = \alpha_1 v_1 + \dots + \alpha_m v_m$$

for some scalars $\alpha_1, \dots, \alpha_m \in \mathbb{F}$. Applying S to both sides, we have

$$STu = S(Tu) = S(\alpha_1 v_1 + \dots + \alpha_m v_m) = \alpha_1 S v_1 + \dots + \alpha_m S v_m.$$

This shows that $\text{range } ST$ is spanned by the vectors $Sv_1, \dots, Sv_m$. Therefore,

$$\dim \text{range } ST \leq m = \dim \text{range } T.$$

Combining the two inequalities, we have

$$\dim \text{range } ST \leq \min(\dim \text{range } S, \dim \text{range } T). \quad \blacksquare$$

Exercise 3B24

- (a) Suppose $\dim V = 5$ and $S, T \in \mathcal{L}(V)$ are such that $ST = 0$. Prove that $\dim \text{range } TS \leq 2$.
- (b) Give an example of $S, T \in \mathcal{L}(\mathbb{F}^5)$ with $ST = 0$ and $\dim \text{range } TS = 2$.

Solution.

- (a) Since $ST = 0$, then $S(Tv) = 0$ for all $v \in V$. This implies that $\text{range } T \subseteq \text{null } S$. By the rank-nullity theorem, we have

$$5 = \dim \text{null } S + \dim \text{range } S \geq \dim \text{range } T + \dim \text{range } S.$$

Now, by [Exercise 3B23](#), we have

$$\dim \text{range } TS \leq \min(\dim \text{range } T, \dim \text{range } S).$$

Suppose that $\dim \text{range } TS > 2$. Then both $\dim \text{range } T$ and $\dim \text{range } S$ must be greater than 2, or equivalently, $\dim \text{range } T \geq 3$ and $\dim \text{range } S \geq 3$. This implies that

$$\dim \text{range } T + \dim \text{range } S \geq 3 + 3 = 6 > 5,$$

which is a contradiction. Therefore, $\dim \text{range } TS \leq 2$.

- (b) Let v_1, v_2, v_3, v_4, v_5 be a basis of $\mathbb{F}^5$. Define $S, T \in \mathcal{L}(\mathbb{F}^5)$ by

$$Sv_1 = v_1, \quad Sv_2 = v_2, \quad Sv_3 = Sv_4 = Sv_5 = 0,$$

and

$$Tv_1 = v_3, \quad Tv_2 = v_4, \quad Tv_3 = Tv_4 = Tv_5 = 0.$$

Then $ST = 0$ since $S(Tv_i) = S(0) = 0$ for all $i = 1, \dots, 5$. Now, we compute $\text{range } TS$. We have

$$TSv_1 = T(Sv_1) = Tv_1 = v_3, \quad TSv_2 = T(Sv_2) = Tv_2 = v_4, \quad TSv_3 = TSv_4 = TSv_5 = 0.$$

Therefore, $\text{range } TS = \text{span}(v_3, v_4)$, which has dimension 2. Thus, we have constructed an example of $S, T \in \mathcal{L}(\mathbb{F}^5)$ with $ST = 0$ and $\dim \text{range } TS = 2$. ■

Exercise 3B25

Suppose that W is finite-dimensional and $S, T \in \mathcal{L}(V, W)$. Prove that $\text{null } S \subseteq \text{null } T$ if and only if there exists $E \in \mathcal{L}(W)$ such that $T = ES$.

Remark. This exercise requires showing that some $R \in \mathcal{L}(\text{range } S, W)$ is well-defined. An issue arises when T is not injective: for some $w \in \text{range } S$, there may be multiple $v_1, \dots, v_k \in V$ such that $Sv_1 = \dots = Sv_k = w$. In this case, we need to show that $Tv_1 = \dots = Tv_k$.

Solution. Suppose $\text{null } S \subseteq \text{null } T$. Define a linear map $R : \text{range } S \rightarrow W$ by $R(Sv) = Tv$ for all $v \in V$. We need to show that R is well-defined. Suppose $Sv_1 = Sv_2$ for some $v_1, v_2 \in V$. Then $S(v_1 - v_2) = 0$, which implies that $v_1 - v_2 \in \text{null } S$. Since $\text{null } S \subseteq \text{null } T$, we have $v_1 - v_2 \in \text{null } T$, which means that $T(v_1 - v_2) = 0$. Therefore, $Tv_1 = Tv_2$, and hence $R(Sv_1) = R(Sv_2)$. Thus, R is well-defined.

By [Exercise 3A13](#), we can extend R to some $E \in \mathcal{L}(W)$. Then for any $v \in V$, we have

$$Tv = R(Sv) = E(Sv),$$

and hence $T = ES$.

Conversely, suppose there exists $E \in \mathcal{L}(W)$ such that $T = ES$. Let $v \in \text{null } S$. Then $Sv = 0$, and applying E gives

$$Tv = E(Sv) = E(0) = 0.$$

Therefore, $v \in \text{null } T$, and hence $\text{null } S \subseteq \text{null } T$. ■

Exercise 3B26

Suppose that V is finite-dimensional and $S, T \in \mathcal{L}(V, W)$. Prove that $\text{range } S \subseteq \text{range } T$ if and only if there exists $E \in \mathcal{L}(V)$ such that $S = TE$.

Remark. From the previous exercise, the reader may be tempted to show that a map is well-defined again. However, this is not necessary here, since we are defining a map on the entire space V , not just on $\text{range } S$.

Solution. Suppose $\text{range } S \subseteq \text{range } T$. Let $v_1, \dots, v_n$ be a basis of V . For each $i = 1, \dots, n$, since $Sv_i \in \text{range } S \subseteq \text{range } T$, there exists $u_i \in V$ such that $Tu_i = Sv_i$. Define a linear map $E \in \mathcal{L}(V)$ by

$$Ev_i = u_i \quad \text{for } i = 1, \dots, n.$$

Then for any $v \in V$, we can express v as a linear combination of the basis vectors:

$$v = \alpha_1 v_1 + \dots + \alpha_n v_n$$

for some scalars $\alpha_1, \dots, \alpha_n \in \mathbb{F}$. Applying S and TE to both sides, we have

$$Sv = \alpha_1 Sv_1 + \dots + \alpha_n Sv_n = \alpha_1 Tu_1 + \dots + \alpha_n Tu_n = T(\alpha_1 u_1 + \dots + \alpha_n u_n) = TEv.$$

Therefore, $S = TE$.

Conversely, suppose there exists $E \in \mathcal{L}(V)$ such that $S = TE$. Let $w \in \text{range } S$. Then there exists $v \in V$ such that $Sv = w$. Applying the definition of S , we have

$$w = Sv = TE(v) = T(E(v)).$$

Therefore, $w \in \text{range } T$, and hence $\text{range } S \subseteq \text{range } T$. ■

Exercise 3B27

Suppose $P \in \mathcal{L}(V)$ and $P^2 = P$. Prove that $V = \text{null } P \oplus \text{range } P$.

Solution. Let $v \in V$. We can express v as

$$v = (v - Pv) + Pv.$$

Observe that $Pv \in \text{range } P$ by definition. Now, we need to show that $v - Pv \in \text{null } P$. Applying P to $v - Pv$, we have

$$P(v - Pv) = Pv - P^2v = Pv - Pv = 0.$$

Therefore, $v - Pv \in \text{null } P$. This shows that every vector $v \in V$ can be expressed as a sum of a vector in $\text{null } P$ and a vector in $\text{range } P$, which implies that

$$V = \text{null } P + \text{range } P.$$

Next, we need to show that the sum is direct. Suppose $u \in \text{null } P \cap \text{range } P$. Then there exists some $w \in V$ such that $u = Pw$. Since $u \in \text{null } P$, we have

$$Pu = P(Pw) = P^2w = Pw = u.$$

But since $u \in \text{null } P$, we also have $Pu = 0$. Therefore, we have

$$u = Pu = 0.$$

This shows that the intersection of $\text{null } P$ and $\text{range } P$ is trivial, i.e., $\text{null } P \cap \text{range } P = \{0\}$. Hence, the sum is direct, and we conclude that

$$V = \text{null } P \oplus \text{range } P. \quad \blacksquare$$

Exercise 3B28

Suppose $D \in \mathcal{L}(\mathcal{P}(\mathbb{R}))$ is such that $\deg Dp = (\deg p) - 1$ for every nonconstant polynomial $p \in \mathcal{P}(\mathbb{R})$. Prove that D is surjective.

Remark. Note that $\mathcal{P}(\mathbb{R})$ is infinite-dimensional, so we cannot use the rank-nullity theorem directly. Instead, one should consider finite-dimensional subspaces of $\mathcal{P}(\mathbb{R})$.

Solution. Let $p \in \mathcal{P}(\mathbb{R})$ be an arbitrary polynomial. We want to show that there exists a polynomial $q \in \mathcal{P}(\mathbb{R})$ such that $Dq = p$. Consider the subspace $\mathcal{P}_{n+1}(\mathbb{R})$ of polynomials of degree at most $n + 1$, where $n = \deg p$. Since $\deg Dq = (\deg q) - 1$, we have $D : \mathcal{P}_{n+1}(\mathbb{R}) \rightarrow \mathcal{P}_n(\mathbb{R})$. The dimension of $\mathcal{P}_{n+1}(\mathbb{R})$ is $n + 2$, and the dimension of $\mathcal{P}_n(\mathbb{R})$ is $n + 1$. By the rank-nullity theorem, we have

$$\dim \text{null } D + \dim \text{range } D = \dim \mathcal{P}_{n+1}(\mathbb{R}) = n + 2.$$

Since $\deg Dp = (\deg p) - 1$ for every nonconstant polynomial p , the kernel of D consists of constant polynomials, which has dimension 1. Therefore, $\dim \text{null } D = 1$, and we have

$$\dim \text{range } D = n + 2 - 1 = n + 1 = \dim \mathcal{P}_n(\mathbb{R}).$$

Since $\dim \text{range } D = \dim \mathcal{P}_n(\mathbb{R})$, it follows that $\text{range } D = \mathcal{P}_n(\mathbb{R})$. Therefore, for any polynomial $p \in \mathcal{P}_n(\mathbb{R})$, there exists a polynomial $q \in \mathcal{P}_{n+1}(\mathbb{R})$ such that $Dq = p$. Since p was arbitrary, we conclude that D is surjective. $\blacksquare$

Exercise 3B29

Suppose $p \in \mathcal{P}(\mathbb{R})$. Prove that there exists a polynomial $q \in \mathcal{P}(\mathbb{R})$ such that $5q'' + 3q' = p$.

Solution. Let $p \in \mathcal{P}(\mathbb{R})$ be an arbitrary polynomial. We want to show that there exists a polynomial $q \in \mathcal{P}(\mathbb{R})$ such that $5q'' + 3q' = p$. Consider the subspace $\mathcal{P}_{n+2}(\mathbb{R})$ of polynomials of degree at most $n + 2$, where $n = \deg p$. Define the linear map $D : \mathcal{P}_{n+2}(\mathbb{R}) \rightarrow \mathcal{P}_{n+1}(\mathbb{R})$ by $Dq = 5q'' + 3q'$. The

dimension of $\mathcal{P}_{n+2}(\mathbb{R})$ is $n + 3$, and the dimension of $\mathcal{P}_{n+1}(\mathbb{R})$ is $n + 2$. By the rank-nullity theorem, we have

$$\dim \text{null } D + \dim \text{range } D = \dim \mathcal{P}_{n+2}(\mathbb{R}) = n + 3.$$

The kernel of D consists of polynomials q such that $5q'' + 3q' = 0$, or $5q'' = -3q'$. Note that if $\deg q = m$, then $\deg q' = m - 1$ and $\deg q'' = m - 2$. Therefore, the degree of $5q'' + 3q'$ is at most $m - 1$. For this to be zero, we must have $q' = 0$, which implies that q is a constant polynomial. Therefore, $\dim \text{null } D = 1$, and we have

$$\dim \text{range } D = n + 3 - 1 = n + 2 = \dim \mathcal{P}_{n+1}(\mathbb{R}).$$

Since $\dim \text{range } D = \dim \mathcal{P}_{n+1}(\mathbb{R})$, it follows that $\text{range } D = \mathcal{P}_{n+1}(\mathbb{R})$. Therefore, for any polynomial $p \in \mathcal{P}_{n+1}(\mathbb{R})$, there exists a polynomial $q \in \mathcal{P}_{n+2}(\mathbb{R})$ such that $Dq = p$. Since p was arbitrary, we conclude that D is surjective. ■

Exercise 3B30

Suppose $\varphi \in \mathcal{L}(V, \mathbb{F})$ and $\varphi \neq 0$. Suppose $u \in V$ is not in $\text{null } \varphi$. Prove that

$$V = \text{null } \varphi \oplus \{au \mid a \in \mathbb{F}\}.$$

Solution. Let $v \in V$. We want to show that v can be expressed as a sum of a vector in $\text{null } \varphi$ and a scalar multiple of u . Since $\varphi(u) \neq 0$, we can define a scalar

$$a = \frac{\varphi(v)}{\varphi(u)}.$$

Then consider the vector

$$w = v - au.$$

We have

$$\varphi(w) = \varphi(v - au) = \varphi(v) - a\varphi(u) = \varphi(v) - \frac{\varphi(v)}{\varphi(u)}\varphi(u) = 0.$$

Therefore, $w \in \text{null } \varphi$. Thus, we can express v as

$$v = w + au,$$

where $w \in \text{null } \varphi$ and au is a scalar multiple of u . This shows that every vector in V can be expressed as a sum of a vector in $\text{null } \varphi$ and a scalar multiple of u , which implies that

$$V = \text{null } \varphi + \{au \mid a \in \mathbb{F}\}.$$

Next, we need to show that the sum is direct. Suppose $w \in \text{null } \varphi \cap \{au \mid a \in \mathbb{F}\}$. Then there exists a scalar $a \in \mathbb{F}$ such that $w = au$. Since $w \in \text{null } \varphi$, we have

$$0 = \varphi(w) = \varphi(au) = a\varphi(u).$$

Since $\varphi(u) \neq 0$, it follows that $a = 0$. Therefore, $w = 0$, which shows that the intersection of $\text{null } \varphi$ and $\{au \mid a \in \mathbb{F}\}$ is trivial. Hence, the sum is direct, and we conclude that

$$V = \text{null } \varphi \oplus \{au \mid a \in \mathbb{F}\}. \quad \blacksquare$$

Exercise 3B31

Suppose V is finite-dimensional, X is a subspace of V , and Y is a finite-dimensional subspace of W . Prove that there exists $T \in \mathcal{L}(V, W)$ such that $\text{null } T = X$ and $\text{range } T = Y$ if and only if $\dim X + \dim Y = \dim V$.

Solution. Suppose there exists $T \in \mathcal{L}(V, W)$ such that $\text{null } T = X$ and $\text{range } T = Y$. By the rank-nullity theorem, we have

$$\dim V = \dim \text{null } T + \dim \text{range } T = \dim X + \dim Y.$$

Conversely, suppose $\dim X + \dim Y = \dim V$. Let $n = \dim V$, $k = \dim X$, and $m = \dim Y$. Then $n = k + m$. Choose a basis $\{x_1, \dots, x_k\}$ of X and extend it to a basis $\{x_1, \dots, x_k, v_1, \dots, v_m\}$ of V . Let $\{y_1, \dots, y_m\}$ be a basis of Y . Define a linear map $T \in \mathcal{L}(V, W)$ by

$$Tx_i = 0 \quad \text{for } i = 1, \dots, k,$$

and

$$Tv_j = y_j \quad \text{for } j = 1, \dots, m.$$

Then for any $v \in V$, we can express v as a linear combination of the basis vectors:

$$v = \alpha_1 x_1 + \dots + \alpha_k x_k + \beta_1 v_1 + \dots + \beta_m v_m$$

for some scalars $\alpha_1, \dots, \alpha_k, \beta_1, \dots, \beta_m \in \mathbb{F}$. Applying T to both sides gives

$$Tv = \alpha_1 Tx_1 + \dots + \alpha_k Tx_k + \beta_1 Tv_1 + \dots + \beta_m Tv_m = 0 + \beta_1 y_1 + \dots + \beta_m y_m.$$

Therefore, $Tv \in Y$. Now suppose $y = \gamma_1 y_1 + \dots + \gamma_m y_m \in Y$. Then

$$T(\gamma_1 v_1 + \dots + \gamma_m v_m) = \gamma_1 y_1 + \dots + \gamma_m y_m = y.$$

Hence, $\text{range } T = Y$. Moreover, if $Tv = 0$, then $\beta_1 y_1 + \dots + \beta_m y_m = 0$. Since $y_1, \dots, y_m$ are linearly independent, it follows that $\beta_1 = \dots = \beta_m = 0$. Thus, $v \in X$. Additionally, if $v \in X$, then $v = \alpha_1 x_1 + \dots + \alpha_k x_k$ for some scalars $\alpha_1, \dots, \alpha_k \in \mathbb{F}$, and hence $Tv = 0$. Therefore, $\text{null } T = X$. This shows that there exists a linear map $T \in \mathcal{L}(V, W)$ such that $\text{null } T = X$ and $\text{range } T = Y$. ■

Exercise 3B32

Suppose V is finite-dimensional with $\dim V > 1$. Show that if $\varphi : \mathcal{L}(V) \rightarrow \mathbb{F}$ is a linear map such that $\varphi(ST) = \varphi(S)\varphi(T)$ for all $S, T \in \mathcal{L}(V)$, then $\varphi = 0$.

Note. The description of the two-sided ideals of $\mathcal{L}(V)$ given by [Exercise 3A17](#) might be useful.

Solution. Suppose $\varphi : \mathcal{L}(V) \rightarrow \mathbb{F}$ is a linear map such that $\varphi(ST) = \varphi(S)\varphi(T)$ for all $S, T \in \mathcal{L}(V)$. We want to show that $\varphi = 0$.

Let I be the identity operator on V . Then for any $T \in \mathcal{L}(V)$, we have

$$\varphi(T) = \varphi(IT) = \varphi(I)\varphi(T).$$

We now claim that φ is a two-sided ideal of $\mathcal{L}(V)$. Let $S \in \mathcal{L}(V)$ and $T \in \varphi$. Then

$$\varphi(ST) = \varphi(S)\varphi(T) = \varphi(S) \cdot 0 = 0,$$

and similarly,

$$\varphi(TS) = \varphi(T)\varphi(S) = 0 \cdot \varphi(S) = 0.$$

Therefore, $ST, TS \in \varphi$, and hence φ is a two-sided ideal of $\mathcal{L}(V)$. By [Exercise 3A17](#), the only two-sided ideals of $\mathcal{L}(V)$ are $\{0\}$ and $\mathcal{L}(V)$ itself. If $\varphi = \{0\}$, then φ is injective. However, since $\dim V > 1$, observe from [Exercise 3A16](#) that we may construct $S, T \in \mathcal{L}(V)$ such that $ST \neq TS$. If $\alpha S + \beta T = 0$ for some scalars $\alpha, \beta \in \mathbb{F}$, then we have

$$(\alpha S + \beta T)v_1 = \alpha S v_1 + \beta T v_1 = \alpha v_2 + \beta v_1 = 0.$$

This implies that $\alpha = \beta = 0$, and hence S and T are linearly independent. Then $\dim \mathcal{L}(V) \geq 2$. Since $\dim \mathbb{F} = 1$, the linear map φ cannot be injective, and hence $\varphi \neq \{0\}$. Therefore, we must have $\varphi = \mathcal{L}(V)$, which implies that $\varphi = 0$. ■

Exercise 3B33

Suppose that V and W are real vector spaces and $T \in \mathcal{L}(V, W)$. Define $T_{\mathbb{C}} : V_{\mathbb{C}} \rightarrow W_{\mathbb{C}}$ by

$$T_{\mathbb{C}}(u + iv) = Tu + iTv$$

for all $u, v \in V$.

- (a) Show that $T_{\mathbb{C}}$ is a (complex) linear map from $V_{\mathbb{C}}$ to $W_{\mathbb{C}}$.
- (b) Show that $T_{\mathbb{C}}$ is injective if and only if T is injective.
- (c) Show that $\text{range } T_{\mathbb{C}} = W_{\mathbb{C}}$ if and only if $\text{range } T = W$.

Note. See Exercise 8 in Section 1B for the definition of the complexification $V_{\mathbb{C}}$. The linear map $T_{\mathbb{C}}$ is called the **complexification** of the linear map T .

Solution.

- (a) Let $u_1 + iv_1, u_2 + iv_2 \in V_{\mathbb{C}}$ and $\alpha, \beta \in \mathbb{C}$. By the definition of complex scalar multiplication in $V_{\mathbb{C}}$, we have

$$\alpha(u_1 + iv_1) + \beta(u_2 + iv_2) = (\alpha u_1 + \beta u_2) + i(\alpha v_1 + \beta v_2).$$

Applying $T_{\mathbb{C}}$ to both sides, we have

$$\begin{aligned} T_{\mathbb{C}}(\alpha(u_1 + iv_1) + \beta(u_2 + iv_2)) &= T_{\mathbb{C}}((\alpha u_1 + \beta u_2) + i(\alpha v_1 + \beta v_2)) \\ &= T(\alpha u_1 + \beta u_2) + iT(\alpha v_1 + \beta v_2) \\ &= \alpha T u_1 + \beta T u_2 + i(\alpha T v_1 + \beta T v_2) \\ &= \alpha(T u_1 + iT v_1) + \beta(T u_2 + iT v_2) \\ &= \alpha T_{\mathbb{C}}(u_1 + iv_1) + \beta T_{\mathbb{C}}(u_2 + iv_2). \end{aligned}$$

Therefore, $T_{\mathbb{C}}$ is a complex linear map from $V_{\mathbb{C}}$ to $W_{\mathbb{C}}$.

- (b) Suppose $T_{\mathbb{C}}$ is injective. Let $v \in V$ such that $Tv = 0$. Then

$$T_{\mathbb{C}}(v + i0) = Tv + iT0 = 0 + i0 = 0.$$

Since $T_{\mathbb{C}}$ is injective, we have $v + i0 = 0$, which implies that $v = 0$. Therefore, T is injective.

Conversely, suppose T is injective. Let $u + iv \in V_{\mathbb{C}}$ such that $T_{\mathbb{C}}(u + iv) = 0$. Then

$$Tu + iTv = 0.$$

This implies that $Tu = 0$ and $Tv = 0$. Since T is injective, we have $u = 0$ and $v = 0$. Therefore, $u + iv = 0$, and hence $T_{\mathbb{C}}$ is injective.

- (c) Suppose $\text{range } T_{\mathbb{C}} = W_{\mathbb{C}}$. Let $w \in W$. Then $w + i0 \in W_{\mathbb{C}}$, and since $\text{range } T_{\mathbb{C}} = W_{\mathbb{C}}$, there exists $u + iv \in V_{\mathbb{C}}$ such that

$$T_{\mathbb{C}}(u + iv) = Tu + iTv = w + i0.$$

This implies that $Tu = w$ and $Tv = 0$. Therefore, $w \in \text{range } T$, and hence $\text{range } T = W$.

Conversely, suppose $\text{range } T = W$. Let $w_1 + iw_2 \in W_{\mathbb{C}}$. Since $\text{range } T = W$, there exist $u, v \in V$ such that $Tu = w_1$ and $Tv = w_2$. Then

$$T_{\mathbb{C}}(u + iv) = Tu + iTv = w_1 + iw_2.$$

Therefore, $w_1 + iw_2 \in \text{range } T_{\mathbb{C}}$, and hence $\text{range } T_{\mathbb{C}} = W_{\mathbb{C}}$. ■

3C Matrices

Exercise 3C1

Suppose $T \in \mathcal{L}(V, W)$. Show that with respect to each choice of bases of V and W , the matrix of T has at least $\dim \text{range } T$ nonzero entries.

Solution. Let $v_1, \dots, v_n$ be a basis of V and $w_1, \dots, w_m$ be a basis of W . Then the matrix of T with respect to these bases is given by

$$\mathcal{M}(T) = \begin{pmatrix} Tv_1 & Tv_2 & \cdots & Tv_n \end{pmatrix}$$

where each Tv_j is expressed as a linear combination of the basis vectors $w_1, \dots, w_m$. Let $\dim \text{range } T = r$. Then there exist r linearly independent vectors in $\text{range } T$. After possibly reordering the basis vectors of W , we can assume that $Tv_1, \dots, Tv_r$ are linearly independent. Since each Tv_j is a linear combination of the basis vectors of W , each Tv_j must have at least one nonzero entry in its corresponding column of the matrix $\mathcal{M}(T)$. Therefore, there are at least r nonzero entries in the matrix $\mathcal{M}(T)$, which shows that the matrix of T has at least $\dim \text{range } T$ nonzero entries. ■

Exercise 3C2

Suppose V and W are finite-dimensional and $T \in \mathcal{L}(V, W)$. Prove that $\dim \text{range } T = 1$ if and only if there exist a basis of V and a basis of W such that with respect to these bases, all entries of $\mathcal{M}(T)$ equal 1.

Solution. Suppose $\dim \text{range } T = 1$. Then there exists a nonzero vector $y_1 \in W$ such that $\text{range } T = \text{span } y_1$. Let $\dim V = n$ and $\dim W = m$. Extend y_1 to a basis $y_1, \dots, y_m$ of W , and define the following vectors in W :

$$w_1 = y_1 - (y_2 + \cdots + y_m), \quad w_k = y_k \text{ for } k = 2, \dots, m.$$

It is clear that $w_1 + w_2 + \cdots + w_m = y_1$, and hence $w_1, \dots, w_m$ spans W . Now suppose for scalars $\alpha_1, \dots, \alpha_m \in \mathbb{F}$, we have

$$\alpha_1 w_1 + \alpha_2 w_2 + \cdots + \alpha_m w_m = 0.$$

Then we can express this as

$$\alpha_1(y_1 - (y_2 + \cdots + y_m)) + \alpha_2 y_2 + \cdots + \alpha_m y_m = 0.$$

Simplifying, we get

$$(\alpha_1)y_1 + (-\alpha_1 + \alpha_2)y_2 + \cdots + (-\alpha_1 + \alpha_m)y_m = 0.$$

Since $y_1, \dots, y_m$ are linearly independent, we must have $\alpha_1 = \alpha_2 = \cdots = \alpha_m = 0$. Therefore, $w_1, \dots, w_m$ are linearly independent and hence form a basis of W .

We now construct a basis of V . By the rank nullity theorem, we have

$$\dim V = \dim T + \dim \text{range } T = \dim T + 1.$$

Let $u_2, \dots, u_n$ be a basis of T . Then we can extend this to a basis $u_1, u_2, \dots, u_n$ of V , where $u_1 \notin T$. Since $\text{range } T = \text{span } y_1$, we have

$$Tu_1 = \beta y_1$$

for some nonzero scalar $\beta \in \mathbb{F}$. If $\beta \neq 1$, we may replace u_1 with $\beta^{-1}u_1$ to ensure that $Tu_1 = y_1$. We construct the following vectors in V :

$$v_1 = u_1, \quad v_k = u_1 + u_k \text{ for } k = 2, \dots, n.$$

Note that for every $j = 1, \dots, n$, we have $Tv_j = Tu_1 + Tu_j = y_1 + 0 = y_1$. We claim that $v_1, \dots, v_n$ form a basis of V . Suppose for scalars $\alpha_1, \dots, \alpha_n \in \mathbb{F}$, we have

$$\alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_n v_n = 0.$$

Then we can express this as

$$(\alpha_1 + \alpha_2 + \dots + \alpha_n)u_1 + \alpha_2 u_2 + \dots + \alpha_n u_n = 0.$$

Since $u_1, \dots, u_n$ are linearly independent, we must have $\alpha_1 + \alpha_2 + \dots + \alpha_n = 0$ and $\alpha_2 = \dots = \alpha_n = 0$. This implies that $\alpha_1 = 0$, and hence $v_1, \dots, v_n$ are linearly independent and form a basis of V . Then with respect to the bases $v_1, \dots, v_n$ of V and $w_1, \dots, w_m$ of W , we have that $Tv_j = y_1 = w_1 + w_2 + \dots + w_m$ for each $j = 1, \dots, n$. Therefore, the matrix of T with respect to these bases has all entries equal to 1.

Conversely, suppose there exist a basis of V and a basis of W such that with respect to these bases, all entries of $\mathcal{M}(T)$ equal 1. Then for each basis vector v_j of V , we have

$$Tv_j = w$$

for some fixed nonzero vector $w \in W$, where $w = w_1 + \dots + w_m$ is the sum of the basis vectors of W . Since $\text{range } T$ is spanned by the images of the basis vectors of V , then any sum of Tv_j is a scalar multiple of w . Therefore, $\text{range } T = \text{span } w$, and hence $\dim \text{range } T = 1$. ■

Exercise 3C3

Suppose $v_1, \dots, v_n$ is a basis of V and $w_1, \dots, w_m$ is a basis of W .

- (a) Show that if $S, T \in \mathcal{L}(V, W)$, then $\mathcal{M}(S + T) = \mathcal{M}(S) + \mathcal{M}(T)$.
- (b) Show that if $\lambda \in \mathbb{F}$ and $T \in \mathcal{L}(V, W)$, then $\mathcal{M}(\lambda T) = \lambda \mathcal{M}(T)$.

Solution.

- (a) Let $v_1, \dots, v_n$ be a basis of V and $w_1, \dots, w_m$ be a basis of W . Then the matrix of $S + T$ with respect to these bases is given by

$$\begin{aligned} \mathcal{M}(S + T) &= \begin{pmatrix} (S + T)v_1 & (S + T)v_2 & \dots & (S + T)v_n \end{pmatrix} \\ &= \begin{pmatrix} Sv_1 + Tv_1 & Sv_2 + Tv_2 & \dots & Sv_n + Tv_n \end{pmatrix} \\ &= \begin{pmatrix} Sv_1 & Sv_2 & \dots & Sv_n \end{pmatrix} + \begin{pmatrix} Tv_1 & Tv_2 & \dots & Tv_n \end{pmatrix} \\ &= \mathcal{M}(S) + \mathcal{M}(T). \end{aligned}$$

- (b) Let $v_1, \dots, v_n$ be a basis of V and $w_1, \dots, w_m$ be a basis of W . Then the matrix of λT with respect

to these bases is given by

$$\begin{aligned}
 \mathcal{M}(\lambda T) &= \begin{pmatrix} (\lambda T)v_1 & (\lambda T)v_2 & \cdots & (\lambda T)v_n \end{pmatrix} \\
 &= \begin{pmatrix} \lambda T v_1 & \lambda T v_2 & \cdots & \lambda T v_n \end{pmatrix} \\
 &= \lambda \begin{pmatrix} T v_1 & T v_2 & \cdots & T v_n \end{pmatrix} \\
 &= \lambda \mathcal{M}(T).
 \end{aligned}$$

■

Exercise 3C4

Suppose that $D \in \mathcal{L}(\mathcal{P}_3(\mathbb{R}), \mathcal{P}_2(\mathbb{R}))$ is the differentiation map defined by $Dp = p'$. Find a basis of $\mathcal{P}_3(\mathbb{R})$ and a basis of $\mathcal{P}_2(\mathbb{R})$ such that the matrix of D with respect to these bases is

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

Solution. Consider the standard basis of $\mathcal{P}_3(\mathbb{R})$. Reorder it to obtain the basis $x^3, x^2, x, 1$. Consider the standard basis of $\mathcal{P}_2(\mathbb{R})$. Reorder it to obtain the basis $x^2, x, 1$. Observe that $D(x^3) = 3x^2$, $D(x^2) = 2x$, $D(x) = 1$, and $D(1) = 0$. However, we need the matrix of D to have the desired form. To achieve this, we can scale the basis vectors of $\mathcal{P}_3(\mathbb{R})$ and $\mathcal{P}_2(\mathbb{R})$ appropriately. Then we may define the basis for $\mathcal{P}_3(\mathbb{R})$ as

$$\frac{1}{3}x^3, \frac{1}{2}x^2, x, 1$$

while maintaining the standard basis for $\mathcal{P}_2(\mathbb{R})$ as $x^2, x, 1$. Then we have

$$D\left(\frac{1}{3}x^3\right) = x^2, \quad D\left(\frac{1}{2}x^2\right) = x, \quad D(x) = 1, \quad D(1) = 0.$$

Therefore, with respect to the bases, we get the desired matrix representation of D . ■

Exercise 3C5

Suppose V and W are finite-dimensional and $T \in \mathcal{L}(V, W)$. Prove that there exist a basis of V and a basis of W such that with respect to these bases, all entries of $\mathcal{M}(T)$ are 0 except that the entries in row k , column k , equal 1 if $1 \leq k \leq \dim \text{range } T$.

Solution. Let $n = \dim V$ and $m = \dim W$. By the rank-nullity theorem, we have

$$\dim V = \dim \text{null } T + \dim \text{range } T.$$

Let $r = \dim \text{range } T$. Let $u_1, \dots, u_{n-r}$ be a basis of $\text{null } T$. Extend this to a basis $u_1, \dots, u_{n-r}, v_1, \dots, v_r$ of V . We claim that $Tv_1, \dots, Tv_r$ are linearly independent. Suppose for scalars $\alpha_1, \dots, \alpha_r \in \mathbb{F}$, we have

$$\alpha_1 Tv_1 + \cdots + \alpha_r Tv_r = 0.$$

Then

$$T(\alpha_1 v_1 + \cdots + \alpha_r v_r) = 0,$$

which implies that $\alpha_1 v_1 + \cdots + \alpha_r v_r \in \text{null } T$. Since $u_1, \dots, u_{n-r}, v_1, \dots, v_r$ is a basis of V , it follows that $\alpha_1 = \cdots = \alpha_r = 0$. Therefore, $Tv_1, \dots, Tv_r$ are linearly independent and form a basis of $\text{range } T$. Extend this to a basis $Tv_1, \dots, Tv_r, w_1, \dots, w_{m-r}$ of W .

If needed, reorder the basis of V so that $v_1, \dots, v_r$ are the first r basis vectors, and reorder the basis of W so that $Tv_1, \dots, Tv_r$ are the first r basis vectors. Then with respect to these bases, we have

$$Tv_k = Tv_k \quad \text{for } k = 1, \dots, r,$$

and

$$Tu_j = 0 \quad \text{for } j = 1, \dots, n - r.$$

Therefore, the matrix of T with respect to these bases has 1's in the diagonal entries corresponding to Tv_k and 0's elsewhere, as desired. ■

Exercise 3C6

Suppose $v_1, \dots, v_m$ is a basis of V and W is finite-dimensional. Suppose $T \in \mathcal{L}(V, W)$. Prove that there exists a basis $w_1, \dots, w_n$ of W such that all entries in the first column of $\mathcal{M}(T)$ [with respect to the bases $v_1, \dots, v_m$ and $w_1, \dots, w_n$] are 0 except for possibly a 1 in the first row, first column.

Solution. Let $n = \dim W$. If $Tv_1 = 0$, then we can choose any basis of W and the first column of $\mathcal{M}(T)$ will be all zeros. If $Tv_1 \neq 0$, then we can extend Tv_1 to a basis of W . Let $w_1 = Tv_1$ and extend it to a basis $w_1, w_2, \dots, w_n$ of W . Then with respect to the bases $v_1, \dots, v_m$ of V and $w_1, \dots, w_n$ of W , we have

$$Tv_1 = w_1$$

and for each $j = 2, \dots, m$, we can express

$$Tv_j = \alpha_{j1}w_1 + \alpha_{j2}w_2 + \cdots + \alpha_{jn}w_n$$

for some scalars $\alpha_{j1}, \alpha_{j2}, \dots, \alpha_{jn} \in \mathbb{F}$. Therefore, the first column of $\mathcal{M}(T)$ has a 1 in the first row and zeros in all other rows, as desired. ■

Exercise 3C7

Suppose $w_1, \dots, w_n$ is a basis of W and V is finite-dimensional. Suppose $T \in \mathcal{L}(V, W)$. Prove that there exists a basis $v_1, \dots, v_m$ of V such that all entries in the first row of $\mathcal{M}(T)$ [with respect to the bases $v_1, \dots, v_m$ and $w_1, \dots, w_n$] are 0 except for possibly a 1 in the first row, first column.

Solution. Let $m = \dim V$. If T is the zero map, then we can choose any basis of V and the first row of $\mathcal{M}(T)$ will be all zeros. If T is not the zero map, let $u_1, \dots, u_m$ be a basis for V . Then there is some u_j such that $Tu_j \neq 0$. Reorder the basis of V so that u_1 is such that $Tu_1 \neq 0$. Then we can express

$$Tu_1 = \alpha w_1 + \beta_2 w_2 + \cdots + \beta_n w_n$$

for some scalars $\alpha, \beta_2, \dots, \beta_n \in \mathbb{F}$. Since $Tu_1 \neq 0$, we have $\alpha \neq 0$. We can scale u_1 by α^{-1} to obtain a new vector $v_1 = \alpha^{-1}u_1$ such that

$$Tv_1 = w_1 + \frac{\beta_2}{\alpha}w_2 + \cdots + \frac{\beta_n}{\alpha}w_n.$$

With the remaining $u_2, \dots, u_m$ basis vectors of V , note that they can be expressed as

$$Tv_j = \alpha_{j1}w_1 + \alpha_{j2}w_2 + \dots + \alpha_{jn}w_n$$

for some scalars $\alpha_{j1}, \alpha_{j2}, \dots, \alpha_{jn} \in \mathbb{F}$. Consider the set of vectors of V given by

$$v_1 = \alpha^{-1}u_1, \quad v_j = u_j - \alpha_{j1}v_1 \text{ for } j = 2, \dots, m.$$

Note that $Tv_j = Tu_j - \alpha_{j1}Tv_1$ is just a combination of $w_2, \dots, w_n$. We now claim that $v_1, v_2, \dots, v_m$ are linearly independent. Suppose for scalars $\gamma_1, \dots, \gamma_m \in \mathbb{F}$, we have

$$\gamma_1v_1 + \gamma_2v_2 + \dots + \gamma_mv_m = 0.$$

Then we can express this as

$$\gamma_1v_1 + \gamma_2(u_2 - \alpha_{21}v_1) + \dots + \gamma_m(u_m - \alpha_{m1}v_1) = 0.$$

Simplifying, we get

$$(\gamma_1 - \gamma_2\alpha_{21} - \dots - \gamma_m\alpha_{m1})v_1 + \gamma_2u_2 + \dots + \gamma_mu_m = 0.$$

Since $v_1, u_2, \dots, u_m$ are linearly independent, we must have $\gamma_1 - \gamma_2\alpha_{21} - \dots - \gamma_m\alpha_{m1} = 0$ and $\gamma_2 = \dots = \gamma_m = 0$. This implies that $\gamma_1 = 0$, and hence $v_1, v_2, \dots, v_m$ are linearly independent. Then $v_1, v_2, \dots, v_m$ form a basis of V . It follows that Tv_1 has a 1 in the first row, while Tv_j for $j = 2, \dots, m$ have zeros in the first row. ■

Exercise 3C8

Suppose A is an m -by- n matrix and B is an n -by- p matrix. Prove that $(AB)_{j\cdot} = A_{j\cdot}B$ for each $1 \leq j \leq m$. In other words, show that row j of AB equals (row j of A) times B .

Solution. Let A be an m -by- n matrix and B be an n -by- p matrix. The entry in row j , column k , of the product AB is given by

$$(AB)_{jk} = \sum_{r=1}^n A_{jr}B_{rk}.$$

The row vector $A_{j\cdot}$ is the j -th row of A , which can be expressed as

$$A_{j\cdot} = (A_{j1}, A_{j2}, \dots, A_{jn}).$$

When we multiply this row vector by the matrix B , we get a new row vector whose entries are given by

$$(A_{j\cdot}B)_k = \sum_{r=1}^n A_{jr}B_{rk}.$$

Comparing this with the expression for $(AB)_{jk}$, we see that

$$(AB)_{jk} = (A_{j\cdot}B)_k$$

for each column index k . Therefore, the entire row vector $(AB)_{j\cdot}$ is equal to the row vector $A_{j\cdot}B$. This proves that row j of AB equals (row j of A) times B . ■

Exercise 3C9

Suppose $a = (a_1 \ \cdots \ a_n)$ is a 1-by- n matrix and B is an n -by- p matrix. Prove that

$$aB = a_1B_{1,\cdot} + \cdots + a_nB_{n,\cdot}.$$

In other words, show that aB is a linear combination of the rows of B , with the scalars that multiply the rows coming from a .

Solution. Let $a = (a_1 \ \cdots \ a_n)$ be a 1-by- n matrix and B be an n -by- p matrix. The product aB is a 1-by- p matrix, and its entries are given by

$$(aB)_{1,k} = \sum_{j=1}^n a_j B_{j,k}$$

for each column index k .

Now, consider the linear combination of the rows of B given by

$$a_1B_{1,\cdot} + a_2B_{2,\cdot} + \cdots + a_nB_{n,\cdot}.$$

The entry in row 1, column k , of this linear combination is

$$(a_1B_{1,\cdot} + a_2B_{2,\cdot} + \cdots + a_nB_{n,\cdot})_k = a_1B_{1,k} + a_2B_{2,k} + \cdots + a_nB_{n,k} = \sum_{j=1}^n a_j B_{j,k}.$$

Comparing this with the expression for $(aB)_{1,k}$, we see that

$$(aB)_{1,k} = (a_1B_{1,\cdot} + a_2B_{2,\cdot} + \cdots + a_nB_{n,\cdot})_k$$

for each column index k . Therefore, we conclude that

$$aB = a_1B_{1,\cdot} + a_2B_{2,\cdot} + \cdots + a_nB_{n,\cdot},$$

which shows that aB is indeed a linear combination of the rows of B , with the scalars coming from the entries of a . ■

Exercise 3C10

Give an example of 2-by-2 matrices A and B such that $AB \neq BA$.

Solution. Let

$$A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

Then we compute the products AB and BA :

$$AB = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 1 \cdot 0 + 2 \cdot 1 & 1 \cdot 1 + 2 \cdot 0 \\ 3 \cdot 0 + 4 \cdot 1 & 3 \cdot 1 + 4 \cdot 0 \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 4 & 3 \end{pmatrix},$$

and

$$BA = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} = \begin{pmatrix} 0 \cdot 1 + 1 \cdot 3 & 0 \cdot 2 + 1 \cdot 4 \\ 1 \cdot 1 + 0 \cdot 3 & 1 \cdot 2 + 0 \cdot 4 \end{pmatrix} = \begin{pmatrix} 3 & 4 \\ 1 & 2 \end{pmatrix}.$$

By comparing the two products, we see that $AB \neq BA$. ■

Exercise 3C11

Prove that the distributive property holds for matrix addition and matrix multiplication. In other words, suppose A, B, C, D, E , and F are matrices whose sizes are such that $A(B + C)$ and $(D + E)F$ make sense. Explain why $AB + AC$ and $DF + EF$ both make sense and prove that

$$A(B + C) = AB + AC \quad \text{and} \quad (D + E)F = DF + EF.$$

Solution. Let A be an m -by- n matrix, and let B and C be n -by- p matrices. Then the product $A(B + C)$ is defined. By [Exercise 3C8](#) and [Exercise 3C9](#), we have

$$A(B + C)_{j,\cdot} = A_{j,\cdot}(B + C) = A_{j,\cdot}B + A_{j,\cdot}C = (AB)_{j,\cdot} + (AC)_{j,\cdot} = (AB + AC)_{j,\cdot}$$

for all $j = 1, \dots, m$. Therefore, $A(B + C) = AB + AC$.

Similarly, let D and E be m -by- n matrices, and let F be an n -by- p matrix. Then the product $(D + E)F$ is defined. By [Exercise 3C8](#) and [Exercise 3C9](#), we have

$$(D + E)F_{j,\cdot} = (D + E)_{j,\cdot}F = D_{j,\cdot}F + E_{j,\cdot}F = (DF)_{j,\cdot} + (EF)_{j,\cdot} = (DF + EF)_{j,\cdot}$$

for all $j = 1, \dots, m$. Therefore, $(D + E)F = DF + EF$. ■

Exercise 3C12

Prove that matrix multiplication is associative. In other words, suppose A, B , and C are matrices whose sizes are such that $(AB)C$ makes sense. Explain why $A(BC)$ makes sense and prove that

$$(AB)C = A(BC).$$

Remark. Unfortunately, we cannot use the same trick as in [Exercise 3C11](#) to prove this result.

Solution. Let A be an m -by- n matrix, B be an n -by- p matrix, and C be a p -by- q matrix. Then the product $(AB)C$ is defined. Going by the definition of matrix multiplication, we have

$$((AB)C)_{j,k} = \sum_{r=1}^p (AB)_{j,r} C_{r,k} = \sum_{r=1}^p \left(\sum_{s=1}^n A_{j,s} B_{s,r} \right) C_{r,k} = \sum_{r=1}^p \sum_{s=1}^n A_{j,s} B_{s,r} C_{r,k}.$$

On the other hand, the product $A(BC)$ is also defined. Again, by the definition of matrix multiplication, we have

$$(A(BC))_{j,k} = \sum_{s=1}^n A_{j,s} (BC)_{s,k} = \sum_{s=1}^n A_{j,s} \left(\sum_{r=1}^p B_{s,r} C_{r,k} \right) = \sum_{s=1}^n \sum_{r=1}^p A_{j,s} B_{s,r} C_{r,k}.$$

Since the order of summation does not matter, we can interchange the order of the sums in the last expression to get

$$(A(BC))_{j,k} = \sum_{r=1}^p \sum_{s=1}^n A_{j,s} B_{s,r} C_{r,k}.$$

Comparing the expressions for $((AB)C)_{j,k}$ and $(A(BC))_{j,k}$, we see that they are equal for all j and k . ■

Exercise 3C13

Suppose A is an n -by- n matrix and $1 \leq j, k \leq n$. Show that the entry in row j , column k , of A^3 (which is defined to mean AAA) is

$$\sum_{p=1}^n \sum_{r=1}^n A_{j,p} A_{p,r} A_{r,k}.$$

Solution. Note that $A^3 = (AA)A = A^2A$. By the definition of matrix multiplication, we have

$$(A^3)_{j,k} = (A^2A)_{j,k} = \sum_{p=1}^n (A^2)_{j,p} A_{p,k}.$$

Now, we can express $(A^2)_{j,p}$ as

$$(A^2)_{j,p} = (AA)_{j,p} = \sum_{r=1}^n A_{j,r} A_{r,p}.$$

Substituting this back into the expression for $(A^3)_{j,k}$, we get

$$(A^3)_{j,k} = \sum_{p=1}^n \left(\sum_{r=1}^n A_{j,r} A_{r,p} \right) A_{p,k} = \sum_{p=1}^n \sum_{r=1}^n A_{j,r} A_{r,p} A_{p,k}.$$

By renaming the indices r and p , we can rewrite this as

$$(A^3)_{j,k} = \sum_{p=1}^n \sum_{r=1}^n A_{j,p} A_{p,r} A_{r,k},$$

which is the desired expression. ■

Exercise 3C14

Suppose m and n are positive integers. Prove that the function $A \mapsto A^t$ is a linear map from $\mathbb{F}^{m,n}$ to $\mathbb{F}^{n,m}$.

Solution. Let $A, B \in \mathbb{F}^{m,n}$ and let $\lambda \in \mathbb{F}$. We need to show that $(A + B)^t = A^t + B^t$ and $(\lambda A)^t = \lambda A^t$.

First, consider the sum $A + B$. The entry in row j , column k , of $A + B$ is given by

$$(A + B)_{j,k} = A_{j,k} + B_{j,k}.$$

Taking the transpose, we have

$$((A + B)^t)_{k,j} = (A + B)_{j,k} = A_{j,k} + B_{j,k} = (A^t)_{k,j} + (B^t)_{k,j} = (A^t + B^t)_{k,j}.$$

Therefore, $(A + B)^t = A^t + B^t$.

Next, consider the scalar multiplication λA . The entry in row j , column k , of λA is given by

$$(\lambda A)_{j,k} = \lambda A_{j,k}.$$

Taking the transpose, we have

$$((\lambda A)^t)_{k,j} = (\lambda A)_{j,k} = \lambda A_{j,k} = \lambda (A^t)_{k,j} = (\lambda A^t)_{k,j}.$$

Therefore, $(\lambda A)^t = \lambda A^t$.

Since both properties hold, we conclude that the function $A \mapsto A^t$ is a linear map from $\mathbb{F}^{m,n}$ to $\mathbb{F}^{n,m}$. ■

Exercise 3C15

Prove that if A is an m -by- n matrix and C is an n -by- p matrix, then

$$(AC)^t = C^t A^t.$$

Solution. Let A be an m -by- n matrix and C be an n -by- p matrix. The entry in row j , column k , of the product AC is given by

$$(AC)_{j,k} = \sum_{r=1}^n A_{j,r} C_{r,k}.$$

Taking the transpose, we have

$$((AC)^t)_{k,j} = (AC)_{j,k} = \sum_{r=1}^n A_{j,r} C_{r,k}.$$

Now, consider the product $C^t A^t$. The entry in row k , column j , of this product is given by

$$(C^t A^t)_{k,j} = \sum_{r=1}^n (C^t)_{k,r} (A^t)_{r,j} = \sum_{r=1}^n C_{r,k} A_{j,r}.$$

Since multiplication in the field $\mathbb{F}$ is commutative, we can rewrite this as

$$(C^t A^t)_{k,j} = \sum_{r=1}^n A_{j,r} C_{r,k},$$

which is exactly the same as the expression for $((AC)^t)_{k,j}$.

Therefore, we conclude that

$$(AC)^t = C^t A^t. \quad \blacksquare$$

Exercise 3C16

Suppose A is an m -by- n matrix with $A \neq 0$. Prove that the rank of A is 1 if and only if there exist $(c_1, \dots, c_m) \in \mathbb{F}^m$ and $(d_1, \dots, d_n) \in \mathbb{F}^n$ such that $A_{j,k} = c_j d_k$ for every $j = 1, \dots, m$ and every $k = 1, \dots, n$.

Solution. Suppose the rank of A is 1. Then the column rank of A is 1, which means that all columns of A are scalar multiples of a single non-zero column vector. Let $c = (c_1, \dots, c_m)^t$ be this non-zero

column vector. Then for each column k of A , there exists a scalar $d_k \in \mathbb{F}$ such that the k -th column of A is given by $d_k c$. Therefore, we can express the entries of A as

$$A_{j,k} = c_j d_k$$

for all $j = 1, \dots, m$ and $k = 1, \dots, n$.

Conversely, suppose there exist $(c_1, \dots, c_m) \in \mathbb{F}^m$ and $(d_1, \dots, d_n) \in \mathbb{F}^n$ such that $A_{j,k} = c_j d_k$ for all j and k . Since $A \neq 0$, then $(c_1, \dots, c_m) \neq 0$ and $(d_1, \dots, d_n) \neq 0$. This means that all columns of A are scalar multiples of the column vector c , and hence the column rank of A is 1. Therefore, the rank of A is 1. ■

Exercise 3C17

Suppose $T \in \mathcal{L}(V)$, and $u_1, \dots, u_n$ and $v_1, \dots, v_n$ are bases of V . Prove that the following are equivalent.

- (a) T is injective.
- (b) The columns of $\mathcal{M}(T)$ are linearly independent in $\mathbb{F}^{n,1}$.
- (c) The columns of $\mathcal{M}(T)$ span $\mathbb{F}^{n,1}$.
- (d) The rows of $\mathcal{M}(T)$ span $\mathbb{F}^{1,n}$.
- (e) The rows of $\mathcal{M}(T)$ are linearly independent in $\mathbb{F}^{1,n}$.

Solution.

- (a) $(a) \iff (b)$: $(a) \iff \dim \text{null } T = 0 \iff \dim \text{range } T = n \iff \text{column rank } \mathcal{M}(T) = n$. For n column vectors, rank n implies linear independence $\iff (b)$.
- (b) $(b) \iff (c)$: n columns of $\mathcal{M}(T)$ are linearly independent in $\mathbb{F}^{n,1} \iff n$ columns of $\mathcal{M}(T)$ span $\mathbb{F}^{n,1}$.
- (c) $(c) \iff (d)$: columns span $\mathbb{F}^{n,1} \iff \text{rank } \mathcal{M}(T) = n$ by (b) $\iff (c)$. By Theorem 3.57, $\text{rank } \mathcal{M}(T) = \text{row rank } \mathcal{M}(T) \iff \text{rows span } \mathbb{F}^{1,n}$.
- (d) $(d) \iff (e)$: n rows of $\mathcal{M}(T)$ span $\mathbb{F}^{1,n} \iff n$ rows of $\mathcal{M}(T)$ are linearly independent in $\mathbb{F}^{1,n}$. ■