

ASTR480 Progress Report

Brayden Leicester

Supervisors: Michele Bannister and Ryan Ridden-Harper

July 16, 2024

This project aims to find and characterise the lightcurves of all the asteroids seen by the Transiting Exoplanet Survey Satellite (TESS).

1 Background

TESS is a large area, high imaging cadence, space telescope (Ricker et al., 2014). TESS is tasked with observing one piece of sky for 27 d at a time (a sector), delivering $96^\circ \times 24^\circ$ full frame images (FFIs) at regular intervals. With the initial cadence for these full frame images set to 30 min, the time resolution of TESS is unparalleled, with a Nyquist frequency of 1 h^{-1} , as the mission was extended the length of the FFIs has come down to 10 min and then 200 s. This high time resolution and observation area does come at the cost of spatial resolution, as the pixels are each $21''$ square. There have been attempts before to find and classify the asteroids in TESS data before by Pál et al. (2018, 2020). This work aims to extend their study to more sectors, and to use a different data reduction method, the `TESSreduce` package (Ridden-Harper et al., 2021). Because of the survey properties, TESS provides a selfconsistent way to measure the properties of asteroids over the full sky. Another beneficial part of this work is that as part of a full sky transient survey using TESS, `TESSELLATE` (Ridden-Harper and Roxburgh et al., in prep), asteroids are transient objects that spike the brightness of a pixel for only a few frames. The goal of finding all the asteroids will allow for the removal of these spikes from the transient pipeline, as well as to understand the asteroid population better.

Asteroids are a key class of solar system objects. Understanding their rotation properties has long been of interest to astronomers (e.g. Weidenschilling, 1981; Harris, 1994). High amplitude variation has come to the forefront of questions about asteroid properties because of the first interstellar object (ISO) 1I/‘Oumuamua (see Bannister et al., 2019, for a review). ‘Oumuamua was measured to have a rotation period of $8.67 \pm 0.34 \text{ h}$ (Belton et al., 2018) and seemed to be tumbling (e.g. Drahus et al., 2018; Fraser et al., 2018). The peak to peak amplitude variation of 2.5 mag (Meech et al., 2017) of the double peaked light curve is of interest, as this is much higher than most asteroids, and it implies a large axis ratio. With the full sky survey of bright asteroids, we hope to find many asteroids with such a large amplitude variation, and to see just how rare ‘Oumuamua is.

2 Work So Far

To check for asteroids in the TESS data, the positions of the asteroids with time are required. For most asteroids, their orbital elements are well known, so it is a matter of looking them up and cross-matching with transients in the TESS data. Python was used to make API calls to Skybot¹ to get positions of asteroids in a cone search box in RA and Dec space. As TESS

¹ Skybot

sectors are 27 d long, querying every 12 h is manageable. These positions are still very sparsely spaced in time compared to the TESS data, so an interpolation is needed to bridge the gap. With TESS data coming in $\frac{1}{2}$ h chunks, 24 interpolated points are needed between each API call. This interpolation should be accurate, as asteroids move at close to a TESS pixel per TESS frame on average (Pál et al., 2018, 2020), and checking against a higher frequency query to JPL Horizons² confirmed this. For the faster TESS data, more interpolated points are needed, but a smaller the change in position between each point.

Matching these interpolated positions to **TESSELLATE** detections is important to lower the unknown transient outputs of this pipeline. Using the **KDTree** algorithm (Maneewongvatana et al., 1999) as implemented in **SciPy** (Virtanen et al., 2020), the right ascension (RA) and declination (Dec) coordinates of the interpolated points and the detections can be compared and matched together. Filtering this **KDTree** output by not allowing the time between spatially coincident matches to be longer than 0.1 d stops any accidental matches in position from non-asteroids.

There are two sets of points to take lightcurves from. The matches from the detections, which already have a flux calculated, or the interpolated points themselves, which are more numerous but require forcing the photometry. Not every interpolated point gets a match, due to a variety of reasons, so a comparison between the two lightcurves is interesting. There were some challenges getting the flux of the interpolated points, even when **TESSELLATE** had already reduced all the FFIs of interest, due to the timing of the TESS frames, but these were identified and corrected for.

3 Future Work

The next part of my analysis has to do with determining the periods and amplitudes of each asteroid’s lightcurve. For this there are a few methods I can try, using the **Lightkurve** package built for period analysis of TESS (and Kepler) data of variable stars, or peel back a layer of abstraction and use the Lomb-Scargle periodogram as implemented by **Astropy**. Some trialling of both methods is needed, as preliminary testing reveals of interesting similarities and differences between the packages.

The **TESSELLATE** pipeline has been running on the OzSTAR supercomputing facilities. After I am confident that all the parts of the asteroid detection and subsequent lightcurve analysis works as required, the same code can be refactored to work on OzSTAR and a large-scale analysis of all the processed TESS sectors can be run. Only after this has completed can the asteroid population statistics can be computed. I will be looking for completeness of detections of asteroids, as well as accuracy of periods and amplitude variation.

4 Figures

References

- Bannister M T, Bhandare A, Dybczyński P A, Fitzsimmons A, Guilbert-Lepoutre A, Jedicke R, Knight M M, Meech K J, McNeill A, Pfalzner S, Raymond S N, Snodgrass C, Trilling D E and Ye Q 2019 *Nature Astronomy* 2019 3:7 **3**, 594–602.
URL: <https://www.nature.com/articles/s41550-019-0816-x>
- Belton M J S, Hainaut O R, Meech K J, Mueller B E A, Kleyna J T, Weaver H A, Buie M W, Drahus M, Guzik P, Wainscoat R J, Waniak W, Handzlik B, Kurowski S, Xu S, Sheppard S S, Micheli M, Ebeling H and Keane J V 2018 *The Astrophysical Journal Letters* **856**, L21.
URL: <https://ui.adsabs.harvard.edu/abs/2018ApJ...856L..21B/abstract>
- Drahus M, Guzik P, Waniak W, Handzlik B, Kurowski S, Xu S, Drahus M, Guzik P, Waniak W, Handzlik B, Kurowski S and Xu S 2018 *NatAs* **2**, 407–412.
URL: <https://ui.adsabs.harvard.edu/abs/2018NatAs...2..407D/abstract>
- Fraser W C, Pravec P, Fitzsimmons A, Lacerda P, Bannister M T, Snodgrass C and Smolić I 2018 *Nature Astronomy* **2**, 383–386.
URL: <https://ui.adsabs.harvard.edu/abs/2018NatAs...2..383F/abstract>
- Harris A W 1994 *Icarus* **107**, 209–211.

² JPL Horizons

- Maneewongvatana S, Mount D M, Maneewongvatana S and Mount D M 1999 *arXiv* p. cs/9901013.
URL: <https://ui.adsabs.harvard.edu/abs/1999cs.....1013M/abstract>
- Meech K J, Weryk R, Micheli M, Kleyna J T, Hainaut O R, Jedicke R, Wainscoat R J, Chambers K C, Keane J V, Petric A, Denneau L, Magnier E, Berger T, Huber M E, Flewelling H, Waters C, Schunova-Lilly E and Chastel S 2017 *Nature* **552**, 378–381.
URL: <https://www.nature.com/articles/nature25020>
- Pál A, Molnár L and Kiss C 2018 *Publications of the Astronomical Society of the Pacific* **130**, 114503.
URL: <https://iopscience.iop.org/article/10.1088/1538-3873/aae2aa>
- Pál A, Szakáts R, Kiss C, Bódi A, Bognár Z, Kalup C, Kiss L L, Marton G, Molnár L, Plachy E, Sárneczky K, Szabó G M and Szabó R 2020 *The Astrophysical Journal Supplement Series* **247**, 26.
URL: <https://iopscience.iop.org/article/10.3847/1538-4365/ab64f0>
- Ricker G R, Winn J N, Vanderspek R, Latham D W, Bakos G Á, Bean J L, Berta-Thompson Z K, Brown T M, Buchhave L, Butler N R, Butler R P, Chaplin W J, Charbonneau D, Christensen-Dalsgaard J, Clampin M, Deming D, Doty J, Lee N D, Dressing C, Dunham E W, Endl M, Fressin F, Ge J, Henning T, Holman M J, Howard A W, Ida S, Jenkins J M, Jernigan G, Johnson J A, Kaltenegger L, Kawai N, Kjeldsen H, Laughlin G, Levine A M, Lin D, Lissauer J J, MacQueen P, Marcy G, McCullough P R, Morton T D, Narita N, Paegert M, Palles E, Pepe F, Pepper J, Quirrenbach A, Rinehart S A, Sasselov D, Sato B, Seager S, Sozzetti A, Stassun K G, Sullivan P, Szentgyorgyi A, Torres G, Udry S and Villaseñor J 2014 *Journal of Astronomical Telescopes, Instruments, and Systems* **1**, 014003.
URL: <http://astronomicaltelescopes.spiedigitallibrary.org/article.aspx?doi=10.1117/1.JATIS.1.1.014003>
- Ridden-Harper R, Rest A, Hounsell R, Müller-Bravo T E, Wang Q and Villar V A 2021 *arXiv e-prints* p. arXiv:2111.15006.
- Virtanen P, Gommers R, Oliphant T E, Haberland M, Reddy T, Cournapeau D, Burovski E, Peterson P, Weckesser W, Bright J, van der Walt S J, Brett M, Wilson J, Millman K J, Mayorov N, Nelson A R J, Jones E, Kern R, Larson E, Carey C J, Polat İ, Feng Y, Moore E W, VanderPlas J, Laxalde D, Perktold J, Cimrman R, Henriksen I, Quintero E A, Harris C R, Archibald A M, Ribeiro A H, Pedregosa F, van Mulbregt P and SciPy 1.0 Contributors 2020 *Nature Methods* **17**, 261–272.
- Weidenschilling S J 1981 *Icarus* **46**, 124–126.