JuMP and MathOptInterface: An optimization framework extensible by design

Benoît Legat (UCLouvain)
Joint work with:
Joaquim Dias Garcia (PUC-Rio), Oscar Dowson (Northwestern) and Miles Lubin (Google)

30 octobre 2026

30th European Conference in Operational Research, 25th June 2019

Extending MathOptInterface

Extending JuMP

Sum-of-Squares extension

Reshaping

Extending MathOptInterface

MathOptInterface (MOI)

MOI in a nutshell :

- `add_variable(model)`.
- `add_constraint(model, func, set)`, e.g. $2x + 3y = 1 \rightarrow$ `(2*x + 3*y)-in-EqualTo(1.0)`.
- `set`, `get` attributes, e.g., `ObjectiveSense`, `ObjectiveFunction`.

Extensible framework :

- Generic on attribute, function and set types. New ones can be defined independently.
- Solver-specific features easily exposed to JuMP/MOI users through custom attributes.
- Expose specialized problem structure easily through custom functions, sets (e.g. Sum-of-Squares variables/constraints).

Semidefinite programming

$$\begin{array}{lll} \underset{Q \in \mathcal{S}^n}{\text{minimize}} & \langle C, Q \rangle & \underset{y \in \mathbb{R}^n}{\text{maximize}} \quad \langle b, y \rangle \\ \text{subject to} & \langle A_i, Q \rangle = b_i & \text{subject to} \quad \sum_i A_i y_i \preceq C \\ & Q \succeq 0 & \end{array}$$

File format : SDPA

Solvers : CSDP, SDPA, DSDP, SDPLR, ...

Variables : Q block diagonal, nonnegative scalar variables (1×1 blocks) or SDP matrices.

Constraints : Affine equations.

Conic Modelling

```
using JuMP
model = Model(...)
@variable(model, -1 <= x <= 1)
@variable(model, y)
@variable(model, z <= 0)
@constraint(model, [x + y  x
                    y      x - y] in PSDCone())
@constraint(model, [x + y, z, y] in SecondOrderCone())
@objective(model, x^2 - 2x*z + z^2)
```

The gap between models and solvers

The solver interface should only support structures and the algorithm exploits :

- n solvers and m structures $\rightarrow$ mn transformations $\rightarrow$ unscalable for large m, n.
- enables evaluation of formulation quality, e.g. automatic transformation and automatic dualization.

The model should

- be independent from solvers.
- represent the structure exploitable by algorithms.
- allow reprentable structure unknown to solvers, e.g. Sum-of-Squares variables/constraints.

Bridging the gap

$$x \in S_1 \Leftrightarrow Ax \in S_2 \qquad AS_1 = S_2$$
$$A^*y \in S_1^* \Leftrightarrow y \in S_2^* \qquad S_1^* = A^*S_2^*$$

In Lagrangian :

$$\langle Ax, y\rangle_2 = \langle x, A^*y\rangle_1$$

Transformation of variable-in-S_2 to variable-in-S_1.

Primal Transform value v to Av.

Dual Transform dual y to $A^{-*}y$.

Transformation of f-in-S_1 constraint to Af-in-S_2 constraint.

Primal Transform value v of Af to $A^{-1}v$ of f.

Dual Transform dual y of A^*y.

Exemples

FlipSignBridge

- **Variable** $x \geq l$ substituted by $x = -y$ where $y \leq -l$.
- **Constraint** $a^\top x \leq \beta$ transformed into $-a^\top x \geq -\beta$.

VectorizeBridge

- **Variable** $x \geq l$ substituted by $x = y + l$ where $y \in \mathbb{R}^1_+$.
- **Constraint** $a^\top x \leq \beta$ transformed into $[a^\top x - \beta] \in \mathbb{R}^1_-$.

FreeBridge

- **Variable** $x \in \mathbb{R}$ substituted by $x = y + z$ where $y \in \mathbb{R}_+$ and $z \in \mathbb{R}_-$.

SlackBridge

- **Constraint** $f \in S$ transformed into $f = x$ for variable $x \in S$.

Selection of bridges

How to select bridges automatically ?

Example
Free variable for SDP solver :

- FreeBridge : $x \in \mathbb{R} \to y \in \mathbb{R}_+$ (supported) and $z \in \mathbb{R}_-$ (not supported)
- FlipSignBridge : $x \in \mathbb{R}_- \to y \in \mathbb{R}_+$.

Shortest path ?

Shortest path in directed Hypergraph

Nodes
Node for each set S (variable-in-S).

Node for each constraint F-in-S.

Types F and S are not limited to those defined in MOI.

Infinitely many nodes, we need to be lazy.

Edges
Each bridge defined possible infinitely many edges.

For each edge and ingoing node : outgoing nodes are

- variable-in-S created.
- constraints F-in-S created.

Solved by a modified Bellman-Ford algorithm [1].

1. See presentation at the Second Annual JuMP-dev Workshop

Extending JuMP

Extending JuMP macros

```
@constraint(model, [x + 1, x - y] in MOI.Zeros())
```

Implementation :

```
function build_constraint(
    _error::Function,
    func::Vector{<:AbstractJuMPScalar},
    set::MOI.AbstractVectorSet)
    return VectorConstraint(x, set)
end
```

Extending JuMP macros : Custom set

```
@constraint(model, [x + 1, x - y] in SecondOrderCone())
```

Implementation :

```
function build_constraint(_error::Function,
                          f::AbstractVector,
                          s::AbstractVectorSet)
    set = moi_set(s, length(f))
    return build_constraint(_error, f, set)
end
function moi_set(::SecondOrderCone, dim::Int)
  return MOI.SecondOrderCone(dim)
end
```

Extending JuMP macros : PSD cone

```
using LinearAlgebra # For Symmetric
@constraint(model, Symmetric([x + 1 x - y
                              x - y y]) in PSDCone())
```

Implementation :

```
function build_constraint(_error::Function,
                          Q::Symmetric,
                          ::PSDCone)
    n = LinearAlgebra.checksquare(Q)
    func = [Q[i, j] for j in 1:n for i in 1:j]
    set = MOI.PositiveSemidefiniteConeTriangle(n)
    VectorConstraint(func, set,
                     SymmetricMatrixShape(n))
end
```

Sum-of-Squares extension

Sum-of-Squares bridges

Polynomial $p \in \Sigma$ (p is SOS) iff $p = X^\top Q x$ with $Q \in \mathbb{S}_+$ (Q is PSD). Hence $\Sigma = A\mathbb{S}_+$.

SOSPolnomialBridge : Transformation of variable-in-Σ to variable-in-$\mathbb{S}_+$.

Transformation of contraint F-in-Σ : SlackBridge + SOSPolnomialBridge.

Result transformations

Constraint Attribute
Examples : ConstraintPrimal, ConstraintDual, ConstraintFunction, ConstraintSet, ...

Redirected to bridge when constraint is bridged.

New attributes :

- GramMatrixAttribute : Gram matrix Q indexed by X.
- MomentMatrixAttribute : Moment matrix index by X, dual of constraint $Q \in \mathbb{S}_+$.
- MomentsAttribute : Vector of moments, dual of constraint $p = X^\top Q X$.

Sum-of-Squares constraint macro

```
@constraint(model, p in SOSCone())
```

Implementation :

```
function JuMP.build_constraint(_error::Function, p,
                                cone::SOSCone; kws...)
    coefs = coefficients(p)
    monos = monomials(p)
    set = JuMP.moi_set(cone, monos; kws...)
    shape = PolyJuMP.PolynomialShape(monos)
    return PolyJuMP.bridgeable(
        JuMP.VectorConstraint(coefs, set, shape),
        JuMP.moi_function_type(typeof(coefs)),
        typeof(set)
    )
end
```

Reshaping

Reshaping results

```
function reshape_vector(vectorized_form::Vector{T},
        shape::SymmetricMatrixShape) where T
    matrix = Matrix{T}(undef, shape.side_dimension,
                       shape.side_dimension)
    k = 0
    for j in 1:shape.side_dimension
        for i in 1:j
            k += 1
            matrix[j, i] = matrix[i, j] =
                vectorized_form[k]
        end
    end
    return Symmetric(matrix)
end
```

Reshaping sets


```
function reshape_set(set::MOI.AbstractScalarSet,
                     ::ScalarShape)
    return set
end
function reshape_set(
    ::MOI.PositiveSemidefiniteConeTriangle,
    ::SymmetricMatrixShape
)
    return PSDCone()
end
```

Reshaping polynomial results

```
function JuMP.reshape_set(set::SOSPolynomialSet,
                          ::PolyJuMP.PolynomialShape)
    return set.cone
end
function JuMP.reshape_vector(x::Vector,
                             shape::PolynomialShape)
    return polynomial(x, shape.monomials)
end
function JuMP.reshape_vector(x::Vector,
                             shape::MomentsShape)
    return measure(x, shape.monomials)
end
function JuMP.dual_shape(shape::PolynomialShape)
    return MomentsShape(shape.monomials)
end
```

Backup

Nonnegative quadratic forms into sum of squares



$$x_1^2 + 2x_1x_2 + 5x_2^2 + 4x_2x_3 + x_3^2 = x^\top \begin{pmatrix} 1 & 1 & 0 \\ 1 & 5 & 2 \\ 0 & 2 & 1 \end{pmatrix} x$$

$$p(x) \geq 0 \ \forall x \iff Q \succeq 0 \qquad \downarrow \text{ cholesky}$$

$$(x_1 + x_2)^2 + (2x_2 + x_3)^2 \longleftarrow x^\top \begin{pmatrix} 1 & 1 & 0 \\ 0 & 2 & 1 \end{pmatrix}^\top \begin{pmatrix} 1 & 1 & 0 \\ 0 & 2 & 1 \end{pmatrix} x$$

Nonnegative polynomial into sum of squares

(x_1, x_2, x_3) (x_1, x_1x_2, x_2) *not* unique

$$p(x) = X^\top Q X$$

$$x_1^2 + 2x_1^2x_2 + 5x_1^2x_2^2 + 4x_1x_2^2 + x_2^2 = X^\top \begin{pmatrix} 1 & 1 & 0 \\ 1 & 5 & 2 \\ 0 & 2 & 1 \end{pmatrix} X$$

$$p(x) \geq 0 \ \forall x \iff Q \succeq 0 \quad \Big| \ \text{cholesky}$$

$$(x_1 + x_1x_2)^2 + (2x_1x_2 + x_2)^2 \longleftarrow X^\top \begin{pmatrix} 1 & 1 & 0 \\ 0 & 2 & 1 \end{pmatrix}^\top \begin{pmatrix} 1 & 1 & 0 \\ 0 & 2 & 1 \end{pmatrix} X$$

When is nonnegativity equivalent to sum of squares ?

Determining whether a polynomial is nonnegative is NP-hard.

Hilbert 1888
Nonnegativity of $p(x)$ of n variables and degree $2d$ is equivalent to sum of squares in the following three cases :

- $n = 1$: Univariate polynomials
- $2d = 2$: Quadratic polynomials
- $n = 2$, $2d = 4$: Bivariate quartics

Motzkin 1967
First explicit example :

$$x_1^4 x_2^2 + x_1^2 x_2^4 + 1 - 3x_1^2 x_2^2 \geq 0 \quad \forall x$$

but is not a sum of squares.

