2024

Baystate Baby Birth Weights



PYCATS

Welcome To Our Presentation

MARGARET GACHAU
CAMERON MOROSKY
BLENDA MUTUMA
MARCELLINE OUMA



DATA ORIGIN AND OVERVIEW




BAYSTATE DATA SOURCE
https://www.key2stats.com/data-set/view/1131

- **Dataset Context:** Covers 189 newborn births at Baystate Medical Center, 1986.
- **Target Variable:** Focuses on ‘low_birth_weight’ as the primary outcome.
- **Risk Factors:** Includes maternal demographic and health-related factors.
- **Purpose:** Provides a basis for analyzing influences on neonatal weight.


Project Objective

- **Project Goal:** Analyze the relationship between infant birth weight and six predictors (three numerical, three categorical).
- **Analysis Objective:** Identify key factors contributing to "low" birth weight classification.
- **Methodology:** Utilize exploratory data analysis and classification modeling to evaluate variable impacts.
- **Outcome:** Determine significant contributors and identify the best machine learning model for prediction accuracy.



CHECK POINTS

189 0BSERVATIONS



- low_birth_weight: indicator of birth weight less than 2.5kg

1. low_birth_weight
2. age_of_mother
3. last_menstrual_weight

1. mother_race
2. smoking_status
3. hypertension_history
4. uterine_irritability
5. physician_visit
6. premature_labor

Data Preprocessing

- **Renamed variables for improved readability.**
- **Dropped irrelevant columns:**
  - birth_weight_grams: Used to create the boolean target variable.
  - weight: Unclear relevance to analysis.
- **Cleaned data for consistency:**
  - Checked numerical variables for non-numeric entries.
  - Ensured consistent labels in categorical variables (e.g., "Yes"/"No").
  - Removed 6 duplicate records.
- **Verified no missing values in the dataset.**
- **Retained outliers as they did not impact model performance in this small dataset.**
- **Mutated numerical column into new categorical variable**
  - age_category: Grouped mothers into 'Teen,' 'Young Adult,' and 'Mature Adult' age brackets.
- **Converted categorical variables to boolean datatypes using One-hot Encoding**




Exploratory Data Analysis: Numerical Variables


Age of Mother:

- Mean age: 23.42 years, Median: 23 years
- Age range: 14 to 45 years
- Most ages fall within 20 to 26.5 years, indicating a dataset primarily composed of younger mothers.
- Distribution is relatively symmetrical.

Last Menstrual Weight:

- Mean weight: 130.09 pounds, Median: 121 pounds
- Weight range: 80 to 250 pounds
- Distribution shows a slight right skew, suggesting higher weights are less common but present.





Exploratory Data Analysis: Categorical Variables

Low Birth Weight:

- 124 infants did not have LBW, 59 did.
- Indicates a significant absence of LBW among the infants studied.

Smoking Status:

- 70 mothers smoked during pregnancy, while 113 did not.
- Smoking is prevalent, impacting birth outcomes.

Mother's Race:

- Distribution: White (92), Other (66), Black (25)
- Majority are white, providing context for exploring racial disparities.


Health-Related Variables:

- Hypertension History: Only 12 out of 183 mothers reported hypertension.
- Uterine Irritability: Reported by 27 mothers.
- Premature Labors: Occurred in 30 mothers.
- Physician Visit: 88 mothers had visits in the first trimester; 95 did not.

Exploratory Data Analysis: Bivariate Insights


Impact of Smoking on Low Birth Weight:

- 42.86% of smoking mothers had LBW babies, compared to 25.66% of non-smoking mothers.
- Strong association between maternal smoking and LBW.



Last Menstrual Weight and Low Birth Weight:

- Mothers of LBW babies generally had lower last menstrual weights.
- Visualized through a box plot showing mean lines and distribution for LBW and non-LBW groups.



Machine Learning

Model selection Justification

Training Process & Evaluation Metrics

Classification Analysis

Data Preprocessing part 2


One-HotEncoding ¶

```
[138]: # one-hot encoding
       # pd.get_dummies(, drop_first=True)
       df=pd.get_dummies(data=df, drop_first=True)
```




```
[135]: # Make a prediction of y based on the optimal threshold, 0.34
       pred_logr2 = pred_proba[:, [1]] >= 0.34
```

	precision	recall	f1-score	support
False	0.83	0.80	0.82	25
True	0.62	0.67	0.64	12
accuracy			0.76	37
macro avg	0.72	0.73	0.73	37
weighted avg	0.76	0.76	0.76	37



Model Selection


```
The best model is Logistic Regression with the Optimal Thresholds, 0.34 with f1_score of 0.64
```

```
The best model is Logistic Regression with the Optimal Thresholds, 0.34 with recall_score of 0.6666666666666666
```

Discussion: Model Selection


Rationale for Choosing Logistic Regression with Optimal Threshold 0.34


- High Performance: Achieved an F1-score of 0.64 and recall of 0.67, indicating strong predictive capability especially in minimizing false negatives.
- Optimal Balance: Threshold of 0.34 provided the best balance between precision and recall compared to other thresholds tested.




	precision	recall	f1-score	support
False	0.83	0.80	0.82	25
True	0.62	0.67	0.64	12
accuracy			0.76	37
macro avg	0.72	0.73	0.73	37
weighted avg	0.76	0.76	0.76	37

Comparison to other Supervised Machine Learning Model





Importance Insights



Last Menstrual Weight
Importance Score: *0.705*


Hypertension History (Yes)
Importance Score: *0.172*

Smoking Status (Yes)
Importance Score: *0.133*

Recommendations



- **Nutritional Support:** Ensure that expecting mothers receive adequate nutritional support to maintain a healthy weight throughout pregnancy. This can involve counseling on balanced diets and providing access to nutritional supplements if needed.



- **Regular Monitoring:** Implement regular weight monitoring during prenatal visits to detect and address any significant weight changes early.



- **Medical Management**: Provide appropriate medical management and treatment for pregnant women with a history of hypertension. This can include medications, lifestyle changes, and regular monitoring of blood pressure.



- **Education and Counseling**: Offer education on managing hypertension through diet, exercise, and stress reduction techniques.




- **Smoking Cessation Programs**: Implement smoking cessation programs specifically tailored for pregnant women. These programs can include counseling, support groups, and nicotine replacement therapies.




- **Awareness Campaigns**: Launch awareness campaigns highlighting the risks of smoking during pregnancy and promoting the benefits of quitting.

Conclusions

Summary of Findings




Project Limitations





THANK YOU! &HAPPY HOLIDAYS



PYCATS!