

Neurology®

TRIGEMINAL NERVE STIMULATION FOR EPILEPSY: LONG-TERM FEASIBILITY AND EFFICACY

Christopher M. DeGiorgio, Diana Murray, Daniela Markovic, et al.

Neurology 2009;72;936-938

DOI 10.1212/01.wnl.0000344181.97126.b4

This information is current as of March 9, 2009

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://www.neurology.org/content/72/10/936.full.html>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.



Table Laboratory values at admission		
Hematologic and chemical values		Serum virology
Analyte	Value	Results
Hematocrit	25%	HIV negative
Mean corpuscular volume	102 μm^3	HTLV-I negative
White cell count	18,300/mm ³	HCV negative
Neutrophils	77%	HBsAg negative
Lymphocytes	9.2%	EBV negative
Eosinophils	4%	CMV negative
Platelet count	232,000/mm ³	HSV-1 negative
Fibrinogen	390 mg/dL	HSV-2 negative
Prothrombin time	31 s	
Urea nitrogen	19 mg/dL	
Creatinine	1.1 mg/dL	
Ammonia	42 mmol/mL	
Glucose	86 mg/dL	
Total bilirubin	12.8 mg/dL	
Conjugated bilirubin	8.1 mg/dL	
Aspartate aminotransferase	161 U/L	
Alanine aminotransferase	64 U/L	

HTLV = human T-cell lymphotropic virus; HCV = hepatitis C virus; EBV = Epstein-Barr virus; CMV = cytomegalovirus; HSV = herpes simplex virus.

IgG. Third, serum contained increased HHV-6 IgM. Remarkably, HHV-6 is closely related to CMV, the second most commonly found infection preceding Guillain-Barré syndrome, and its most common viral trigger.⁶ The normal CSF protein content in our patient seemingly contrasts with evidence that about 80% of patients with AIDP have raised CSF protein concentrations. One explanation is that we collected our patient's CSF sample in the early days of illness, a time when protein content is often normal.⁷ Measurement of potassium, phosphate, and porphyrin metabolism products helped to exclude alternative diagnoses. Stool sample cultures excluded *Campylobacter jejuni* enteritis. In addition, the neurologic examination on admission failed to disclose the cognitive disturbances typical of some alcohol-related

acute reactions, thus arguing against a history of alcohol abuse, and lacked the signs of distal symmetric axonal polyneuropathy typically found in heavy chronic alcohol users.

Viral reactivation of a latent HHV-6 infection may be among the causative or precipitating factors of acute inflammatory PNS diseases.

From the Infectious Disease Unit (C.D.B., A.V., F.S.) and Neurology Unit (A.B.), Ospedale S. Maria Goretti, Latina; Virology (S.Z., C.M.), Istituto Nazionale per le Malattie Infettive Lazaro Spallanzani, Rome; Department of Neurological Sciences, Neurosurgical Unit, Ospedale S. Maria Goretti (P.M.), Department of ENT and Neurology, Neurorehabilitation Unit, ICOT (F.P.), and Department of Neurological Sciences, Neurology Unit, Ospedale A. Fiorini Terracina (A.C.), Sapienza University of Rome, Polo Pontino, Italy.

Disclosure: The authors report no disclosures.

Received August 19, 2008. Accepted in final form November 10, 2008.

Address correspondence and reprint requests to Dr. Antonio Curra, Department of Neurological Sciences, Neurology Unit, Ospedale A. Fiorini Terracina, LT, Sapienza University of Rome, Polo Pontino, Italy; antonio.curra@uniroma1.it

Copyright © 2009 by AAN Enterprises, Inc.

1. Levy JA. Three new human herpesviruses (HHV6, 7, and 8). *Lancet* 1997;349:558–563.
2. Morimoto T, Sato T, Matsuoka A, et al. Trimethoprim-sulfamethoxazole-induced hypersensitivity syndrome associated with reactivation of human herpesvirus-6. *Intern Med* 2006;45:101–105.
3. Gildea DH, Mahalingam R, Cohrs RJ, Tyler KL. Herpesvirus infections of the nervous system. *Nat Clin Pract Neurol* 2007;3:82–94.
4. Miyake F, Yoshikawa T, Suzuki K, Ohashi M, Suga S, Asano Y. Guillain-Barré syndrome after exanthem subitum. *Pediatr Infect Dis J* 2002;21:569–570.
5. Ward KN, Leong HN, Thiruchelvam AD, Atkinson CE, Clark DA. Human herpesvirus 6 DNA levels in cerebrospinal fluid due to primary infection differ from those due to chromosomal viral integration and have implications for diagnosis of encephalitis. *J Clin Microbiol* 2007;45:1298–1304.
6. Hadden RD, Karch H, Hartung HP, et al. Preceding infections, immune factors, and outcome in Guillain-Barre syndrome. *Neurology* 2001;56:758–765.
7. Paradiso G, Tripoli J, Galicchio S, Fejerman N. Epidemiological, clinical, and electrodiagnostic findings in childhood Guillain-Barré syndrome: a reappraisal. *Ann Neurol* 1999;46:701–707.


Christopher M.
DeGiorgio, MD
Diana Murray
Daniela Markovic, MS
Todd Whitehurst, MD

TRIGEMINAL NERVE STIMULATION FOR EPILEPSY: LONG-TERM FEASIBILITY AND EFFICACY

Neurostimulation has emerged as a viable alternative for intractable epilepsy. Trigeminal nerve stimulation (TNS), a novel form of neurostimulation, has an antiepileptic effect in a rodent model.¹ The superficial location of trigeminal branches allows for minimally invasive approaches, allowing assessment of response prior to a permanent device.² We report the long-term safety and efficacy

of external TNS for epilepsy.

Methods. Research committee approval was obtained for an open study of external TNS in epilepsy. Informed consent was obtained before enrollment. Inclusion/exclusion criteria were age 18–65 years, ≥ 3 complex-partial/generalized tonic-clonic seizures/month, no progressive medical conditions, and exposure to ≥ 2 antiepileptic drugs (AEDs). Subjects enrolled in a 4-week pretreatment baseline, and were



evaluated at 1, 2, 3, 6, and 12 months. AEDs remained unchanged unless essential for patient safety. Neurostimulation was initially supplied using the analog EMS Model 400, and later a digital EMS model 7500.² Stimulation settings were as follows: frequency 120 Hz, 250 μ s, ≤ 30 seconds on, ≤ 30 seconds off for 12–24 hours/day, and 1.25-inch disposable, silver-gel, adhesive electrodes were utilized, spaced 2 inches apart. Subjects 1–3 were initiated with infraorbital stimulation (V_2), but subjects found this awkward for chronic use. Since then, we have stimulated the ophthalmic nerve (V_1) in all subjects, which allows bilateral stimulation and can be concealed by a cap. See figure e-1 on the *Neurology*[®] Web site at www.neurology.org for a diagram of the device and electrode placement.

Statistical analysis. We compared mean seizure rates over time using a repeated measure analysis of variance (ANOVA) model. After confirming that missing data were missing at random, we used multiple imputation to replace missing values for daily seizure rate with a set of imputed values in patients who did not complete the study. The imputation was carried out under the repeated measure ANOVA model. The resulting variances of the means reflect not only the variation in the observed data but also the added uncertainty of using partially imputed information. It is preferable to impute missing values rather than omit patients with incomplete data entirely. Four imputed values were generated for each missing value. All analyses were carried out using SAS version 9.1. A p value of 0.05 was chosen for significance.

Results. Thirteen subjects completed the 4-week prospective baseline. Twelve completed 3 months, 10 completed 6 months, and 7 completed 12 months (table e-1). Short-term results from the first seven subjects reported previously are included in this analysis.² The fig-

ure shows results at 3, 6, and 12 months. At 3 months, mean seizure frequency was reduced from a baseline of 2.1 seizures/day to 0.71 seizures/day (66% reduction, $p = 0.034$, standard error 0.64). At 6 months, mean seizure frequency was 0.92 seizures/day (56% reduction, $p = 0.073$, standard error 0.64). At 12 months, mean seizure frequency was 0.86 seizures/day (59% reduction, $p = 0.058$, standard error 0.64). Five subjects experienced greater than 50% reduction at 6 and 12 months; one subject had >90% reduction at 12 months.

TNS was well tolerated. Side effects included skin irritation in five subjects, which improved by reducing stimulation to 12–16 hours/day, or with hydrocortisone 1% cream. Tingling, forehead pressure, and headache were reported, but improved with reduction of current. No effect on pulse, blood pressure, or ECG was detected.

Discussion. External stimulation of the trigeminal nerve (V_1 and V_2) was well tolerated, with a mean reduction in seizure frequency of 59% at 12 months. The results of this study build upon data in an animal model, and our short-term report in humans.^{1,2}

Why stimulate the trigeminal nerve for epilepsy? The trigeminal nerve has three bilateral cutaneous branches.³ These afferent branches (V_1 – V_3) exit foramen 1.25–1.5 inches lateral of the midline, and project to the trigeminal ganglion and the trigeminal nucleus, which projects to the nucleus tractus solitarius and locus ceruleus, all of which play a role in inhibition of seizures.^{4,7} The antiepileptic effect of the trigeminal nerve and related structures has been confirmed in animal models.^{1,4,5} Stimulation of the locus ceruleus suppresses epileptic discharges induced by cobalt and penicillin, and plays a central role in the mechanism of VNS.^{6,7} In a pentylenetetrazole model, high-frequency stimulation of the trigeminal nerve significantly reduced seizure severity and frequency.¹ Bilateral stimulation was more effective than unilateral stimulation, and high frequency stimulation was more effective than low frequency stimulation.¹

The data provide a foundation for future randomized studies of TNS in humans. Based on the long-term feasibility and efficacy of this study, we have embarked on a randomized controlled study of high vs low frequency stimulation of the ophthalmic branch (V_1) of the trigeminal nerve for intractable epilepsy.

From the Departments of Neurology (C.M.D.) and Biomathematics (D. Markovic), David Geffen School of Medicine at UCLA, Los Angeles; Olive View/UCLA Medical Center (D. Murray), Sylmar; and Boston Scientific (T.W.), Valencia, CA.

Supplemental data at www.neurology.org

Supported by an investigator-initiated grant from Advanced Bionics and Boston Scientific and grants from Mr. James Peters, the Salter, Brill, Jacoby, Lagermeier, Lester, and Johnson Families, AMDG. Disclosure: C.M.D. has received grant support from Advanced Bionics/Boston Scientific for this investigator-initiated study. T.W. is an employee of Boston Scientific, Inc.

Received August 29, 2008. Accepted in final form November 18, 2008.

Address correspondence and reprint requests to Dr. Christopher M. DeGiorgio, 710 Westwood Plaza AMDG, UCLA, Los Angeles, CA 90095; cmd@mednet.ucla.edu

Copyright © 2009 by AAN Enterprises, Inc.

AUTHOR CONTRIBUTIONS

Daniela Markovic performed the statistical analysis.

ACKNOWLEDGMENT

The authors thank Dr. Alan Shewmon for his support and assistance.

1. Fanselow EE, Reid A, Nicoletis AL. Reduction of pentylenetetrazole induced seizure activity in awake rats by

seizure-triggered trigeminal nerve stimulation. *J Neurosci* 2000;20:8160–8168.

2. DeGiorgio CM, Shewmon DA, Whitehurst T. Pilot study of trigeminal nerve stimulation (TNS) for epilepsy: a proof-of-concept trial. *Epilepsia* 2006;47:1213–1215.
3. Clemente C. *Anatomy: A Regional Atlas of the Human Body*, 4th ed. Baltimore: Williams & Wilkins; 1997.
4. Magdaleno-Madrigal VM, Valdes Cruz A, Martinez Vargas, et al. Effect of electrical stimulation of the nucleus of the solitary tract on the development of electrical amygdaloid kindling in the cat. *Epilepsia* 2002;43:964–969.
5. Walker BR, Easton A, Gale K. Regulation of limbic motor seizures by GABA and glutamate transmission in nucleus tractus solitarius. *Epilepsia* 1999;40:1051–1057.
6. Weiss GK, Lewis J, Jimenez-Rivera C, Vigil A, Corcoran ME. Anti-kindling effects of locus coeruleus stimulation: mediation by ascending noradrenergic projections. *Exp Neurol* 1990;108:136–140.
7. Kahl SE, Clark K, Smith D, et al. Locus coeruleus lesions suppress the seizure-attenuating effects of vagus nerve stimulation *Epilepsia* 1998;39:709–714.

TRIGEMINAL NERVE STIMULATION FOR EPILEPSY: LONG-TERM FEASIBILITY AND EFFICACY

Christopher M. DeGiorgio, Diana Murray, Daniela Markovic, et al.

Neurology 2009;72;936-938

DOI 10.1212/01.wnl.0000344181.97126.b4

This information is current as of March 9, 2009

Updated Information & Services	including high resolution figures, can be found at: http://www.neurology.org/content/72/10/936.full.html
Supplementary Material	Supplementary material can be found at: http://www.neurology.org/content/suppl/2009/03/09/72.10.936.DC1.html
References	This article cites 6 articles, 1 of which you can access for free at: http://www.neurology.org/content/72/10/936.full.html#ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://www.neurology.org/content/72/10/936.full.html#otherarticles
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://www.neurology.org/misc/addir.xhtml#reprintsus

