

# Vagus nerve stimulation: from pre-clinical to clinical application: challenges and future directions

Gaetano M. De Ferrari · Peter J. Schwartz

Published online: 17 December 2010  
© Springer Science+Business Media, LLC 2010

**Abstract** Vagus nerve stimulation was performed experimentally for the first time more than 150 years ago. In the 1980s and 1990s, vagus nerve stimulation was shown, both in the anesthetized and in the conscious animal, to exert marked antiarrhythmic effects, particularly during acute myocardial ischemia. There is a strong rationale for a beneficial effect of augmented vagal activity in the setting of chronic heart failure. Studies in experimental models of heart failure showed that chronic vagus nerve stimulation exerts beneficial effects on left ventricular function and on survival. Vagus nerve stimulation is approved in man for refractory epilepsy and depression. The first-in-man study performed in 32 patients with chronic heart failure suggests that vagus nerve stimulation was safe and well tolerated. Six months of open-label

treatment was associated with significant improvements ( $P < 0.001$ ) in NYHA class, quality of life, 6-min walk test, LV ejection fraction (from  $22 \pm 7$  to  $29 \pm 8\%$ ), and LV systolic volumes ( $P = 0.02$ ). These improvements were maintained at 1 year. Mechanisms of action may include the following: heart rate, anti-adrenergic, anti-apoptotic, and anti-inflammatory effects as well as an increase in nitric oxide. Controlled clinical trials will start soon to assess whether vagus nerve stimulation can indeed represent a new non-pharmacological approach for the treatment of symptomatic heart failure.

**Keywords** Heart failure · Vagal stimulation · Ventricular arrhythmia

---

G. M. De Ferrari (✉) · P. J. Schwartz  
Department of Cardiology, Fondazione IRCCS  
Policlinico S. Matteo, Viale Golgi 19, 27100 Pavia, Italy  
e-mail: g.deferrari@smatteo.pv.it

P. J. Schwartz  
Department of Lung, Blood and Heart, Section of Cardiology,  
University of Pavia, Pavia, Italy

P. J. Schwartz  
Laboratory of Cardiovascular Genetics, IRCCS Istituto  
Auxologico Italiano, Milan, Italy

P. J. Schwartz  
Department of Medicine, Cardiovascular Genetics Laboratory,  
Hatter Institute for Cardiovascular Research, University of Cape  
Town, Cape Town, South Africa

P. J. Schwartz  
Chair of Sudden Death, Department of Family and Community  
Medicine, College of Medicine, King Saud University, Riyadh,  
Saudi Arabia

## Introduction

Research into vagus nerve stimulation has a long and storied history. Much of the early work was performed in animal models with the goal of describing its effects on either cardiovascular physiology or cardiac rhythm disturbances. We will therefore examine these latter studies and then focus on heart failure. We will also provide some concluding thoughts that help to synthesize many of the concepts raised in this issue of HF Reviews.

## Vagal stimulation in experimental models of arrhythmias: historical and recent studies

The first study was performed more than 150 years ago. Einbrodt assessed the effects of an inductorium on heart beats and noticed that during vagal stimulation, the distance between the coils could be markedly reduced,

thereby increasing the current delivered, before ventricular fibrillation (VF) occurred and the dog died [1]. Despite this pioneering observation, dating back to 1859, the dominant view [2–4], with few exceptions [5, 6], was that vagal hyperactivity is proarrhythmic and favors the occurrence of VF. The proarrhythmic effect of vagal stimulation was not confirmed in dogs subjected to coronary artery occlusion by Scherlag et al. [7] and by Kerzner et al. [8] and was seriously questioned in the early 1970s [9–11]. In the chloralose-anesthetized cat, bilateral vagotomy increased the incidence of VF produced by occlusion of the left anterior descending (LAD) coronary artery from 20 to 55%, independently of heart rate effects [12]. Subsequently, Yoon et al. [13, 14] found in anesthetized dogs that heart rate reduction mediated the antiarrhythmic effects of vagal stimulation during coronary occlusion and that an anti-adrenergic mechanism played a major role in the increase in VF threshold (VFT). Similar conclusions were reached by Kolman et al. [15] and by Furey and Levy [16], who showed that vagal stimulation also counteracted the effect of isoproterenol infusion suggesting post-junctional antagonism. This was at variance with Kent et al. [9] and Yoon et al. [13], who showed an increase in VFT independent of sympathetic stimulation. Many years later, the question of whether vagal stimulation has biologically significant effects at ventricular level independent of its anti-adrenergic action is still a matter of debate.

Malignant arrhythmias also occur following reperfusion of the ischemic myocardium [17]. We evaluated the effects of bilateral vagotomy and of vagal stimulation on reperfusion arrhythmias in cats [18]. Vagal stimulation decreased the incidence of malignant arrhythmias dramatically when heart rate was allowed to decrease and moderately when heart rate was kept constant. Although reperfusion arrhythmias were considered to have limited clinical relevance, reperfusion of the ischemic myocardium produces a cascade of events resulting in cell death and dysfunction that are grouped under the definition of reperfusion injury [19], and it is likely that the mechanisms leading to electrical or mechanical abnormalities share common pathways that may be favorably affected by vagal stimulation. Indeed, a series of recent experimental studies have found promising results for the potential of vagal stimulation to attenuate reperfusion injury and subsequent cardiac remodeling (see section on future developments).

All experimental studies described above involved anesthetized animals and conditions distant from the one potentially encountered clinically. Therefore, we assessed the effects of vagal stimulation in a conscious animal model for sudden cardiac death [20] in which VF may be reproducibly induced for several months in approximately 55% of the animals (“susceptible”), allowing internal control studies. In this model, vagal activity was essential for the

prevention of VF in 11 of 45 animals originally resistant to VF and tested after atropine [21]. Additionally, susceptible dogs underwent a further exercise and ischemia test with no additional intervention (control group) or with right vagal stimulation started few seconds after the beginning of coronary occlusion [22]. VF occurred in 23 of 25 (92%) control animals, but only in 3 of 26 (11.5%) vagally stimulated dogs. When heart rate was not allowed to decrease, 5 of 9 animals (55%) remained protected from VF. We stimulated the intact cervical vagus, and thus, both efferent and afferent fibers were stimulated. Since vagal afferent stimulation produces a reflex withdrawal of sympathetic efferent activity [23], this withdrawal might have significantly influenced our results. However, this appears unlikely since atropine prevented both the chronotropic and the antiarrhythmic effects of vagal stimulation in this model. Thus, in the exercising dog with high levels of circulating catecholamines, a transient decrease in the firing of sympathetic nerves is unlikely to grossly affect cardiac electrophysiology. Notably, the picture is likely to be different when dealing with long-term studies aiming, for instance, at providing anti-remodeling effects in patients with heart failure: in this setting, simultaneous sympathetic withdrawal may very well contribute to the beneficial effects of chronic vagal stimulation. Summarizing, vagal hyperactivity drastically reduces malignant arrhythmia, and the reduction in heart rate is not a crucial component in half of the protected animals. Favorable effects were also observed in this same animal model with pharmacological vagal activation [24].

In a more recent rat study [25], vagal stimulation confirmed a striking antiarrhythmic effect during myocardial ischemia and prevented >50% of the loss of connexin 43 induced by ischemia, a new potential mechanism involved in the vagally mediated increase in electrical stability.

### Vagal stimulation and arrhythmias in patients

Until very recently, all information regarding vagal activation in patients with cardiac disorders was obtained from indirect evidence. The background regarding the predictive role of a marker of vagal reflexes such as baroreflex sensitivity is discussed elsewhere in this Supplement [26]. More relevant to the current topic are the few clinical studies examining the effects of reflexly induced vagal hyperactivity, mostly on ventricular arrhythmias. Waxman and colleagues provided the first evidence that some ventricular tachycardias (VTs) could respond to vagal activation, contrary to traditional belief [27, 28], and that ventricular automaticity was decreased by vagal activity [29].

We have assessed the antiarrhythmic effect of reflexly induced vagal activation, in patients with frequent, stable premature ventricular contractions (PVCs) [30]. When

heart rate was allowed to decrease, phenylephrine completely abolished ectopic activity in 9 of the 10 patients studied. During atrial pacing at the pre-drug rate, PVCs reappeared in most subjects. Their mean number, however, was still significantly less than the control values, which suggests that the antiarrhythmic effect of reflex vagal activation is only partially rate-dependent.

### Vagal stimulation in experimental models of heart failure

Chronic electrical vagal stimulation has been assessed in three animal models of heart failure. Rats subjected to a large anterior myocardial infarction (MI) leading to heart failure were randomized to vagal and sham-stimulated groups 14 days post-MI. The vagus was stimulated for 10 s/min, with an intensity adjusted to reduce heart rate by 20–30 bpm. Rats randomized to vagal stimulation showed significant improvement in left ventricular (LV) hemodynamics and decreased mortality from 50 to 14% at 140 days (Fig. 1) [31]. In an established canine model of intracoronary microembolization-induced heart failure, vagal stimulation significantly improved LV function compared to sham-operated animals [32]. In the same animal model, therapy with vagus nerve electrical stimulation combined with beta-blockade improved left ventricular systolic function beyond that seen with beta-blockade alone [33].

More recently, the effects of chronic vagal stimulation in a canine pacing model of heart failure (HF) were

presented [34]. These dogs were subjected to 8 weeks of high-rate ventricular pacing with concomitant vagus nerve stimulation in the active group and no stimulation in the control group. At the end of the 12-week study period, vagally stimulated animals had significantly lower LV end-diastolic and end-systolic volumes and higher LV ejection fraction (EF). This result was obtained in the absence of any heart rate effect provided by vagal stimulation since both groups were subjected to the same constant ventricular pacing.


Finally, vagus nerve stimulation at a low intensity, not causing any heart rate reduction, exerted positive effect on LV function and multiple biomarkers in the canine microembolization HF model [35].

### Vagal stimulation and heart failure in patients

In patients, vagus nerve stimulation has been approved for the treatment of drug-refractory epilepsy [36–38] and, subsequently, depression [39, 40].

Given the strong rationale for vagal stimulation in patients with heart failure, we performed a single-arm open-label interventional phase II study that followed a 2-staged approach [41, 42]: a single-center feasibility phase followed by a multicenter phase assessing primarily safety and tolerability of chronic vagus nerve stimulation. A secondary objective of the study was to collect preliminary information on efficacy.


We enrolled 32 patients (94% men; mean age  $56 \pm 11$  years) with a history of chronic heart failure in symptomatic NYHA class II–IV and average LVEF of  $23 \pm 8\%$ . The patients were on a very comprehensive background medical treatment including beta-blockers, ACE inhibitors/ARBs, and loop diuretics in 31 out of 32; implantable cardioverter-defibrillators (ICDs) were previously implanted in 19 out of 32. The main inclusion and exclusion criteria are described in detail elsewhere [41, 42]. These patients underwent implantation of the CardioFit 5000 system (BioControl Medical, Yehud, Israel), including an implantable neurostimulator, capable of delivering low-current electrical pulses, with adjustable parameters, to stimulate the vagus nerve. The stimulator is designed to sense the heart rate (via an intracardiac implanted electrode) and deliver stimulation at preset delays from the R wave as well as to interrupt vagal stimulation when the heart rate reaches the bradycardia limit (e.g., 55 b/min). The stimulation lead is an asymmetric bipolar multi-contact cuff electrode specifically designed for cathodic induction of action potentials in the vagus nerve while simultaneously applying asymmetrical anodal blocks, which are expected to lead to preferential, but not exclusive, activation of efferent fibers.



**Fig. 1** Survival curves of rats subjected to a large anterior myocardial infarction leading to heart failure and randomized 14 days post-MI to vagal stimulation or sham stimulation. Rats randomized to vagal stimulation showed a significantly lower mortality. (Modified from Ref. [31])

Within 4 weeks of implantation, a 3-week stimulation up-titration phase was started with the goal of reaching 5.5 mAmp, a 5–10 b/min heart rate reduction or onset of side effects. The stimulation intensity reached  $4.1 \pm 1.2$  mA and was limited by discomfort or pain in the majority of patients. Two serious adverse events (SAEs) were clearly related to the procedure: one acute pulmonary edema occurring after surgery and treated with iv diuretics and one need of surgical revision of the device. Two SAEs were possibly related to the experimental treatment: an episode of atrial fibrillation leading to electrical cardioversion and an episode of ulcer bleeding in a *Helicobacter pylori*-positive patient.

Heart rate changes observed during vagal stimulation (10-s ON time: red crosses, 30-s OFF time: blue crosses) in one patient are shown in Fig. 2. The average acute heart rate changes observed during vagal stimulation were modest (1.5 b/min). However, baseline resting heart rate decreased significantly during the study from  $82 \pm 13$  to  $76 \pm 13$  b/min. Most patients improved by at least one NYHA class both at 3 months (18/32, 56%) and at 6 months (19/32, 59%). Quality of life markedly improved at 3 months (from  $49 \pm 17$  to  $33 \pm 16$  with the Minnesota Living with Heart Failure® Questionnaire), thereafter remaining almost identical at 6 months ( $32 \pm 19$ ). The same was found for the 6-min walk test with an increase at 3 months from  $411 \pm 76$  to  $470 \pm 99$  m and subsequent stability. The blinded echocardiogram analysis disclosed a non-significant decrease in LV end-diastolic volume, a significant reduction in LV end-systolic volume (see Fig. 3), and a significant increase in LV ejection fraction (from  $22 \pm 7\%$  to  $29 \pm 8\%$ ). A group of 23 patients




**Fig. 3** Left ventricular volume indexes. LVEDVI: left ventricular end-diastolic volume index. LVESVI: left ventricular end-systolic volume index. (From Ref. [42])

continued their follow-up with active vagal stimulation. Analysis of clinical variables in this group of patients revealed maintenance and even magnification of the favorable effects of vagal stimulation at 1 year (especially LVEF from 21 to 34%).

This first human experience of chronic vagal stimulation in patients with heart failure suggested that the treatment is feasible, safe and tolerable and leads to a subjective clinical improvement. The study also provided the first evidence to suggest that this new therapeutic approach may produce favorable and long-lasting effects on left ventricular function. The preservation of these effects at 1-year follow-up strongly argues against a major role of a possible placebo effect.

Thus, this novel therapeutic approach appears promising and warrants further investigation. Accordingly, two trials

**Fig. 2** Heart rate observed during vagal stimulation (5.5 mAmp, 10 s ON time: red crosses, 30 s OFF time: blue crosses) in one patient. A reduction of approximately 10 b/min, starting from a baseline heart rate of 110 b/min, occurred during the 10-s train of pulses. (From Ref. [41])



(a phase III study using this same device and a phase II study with a different device) will start soon. They will provide a more precise definition of the role of vagal stimulation in heart failure.

### Mechanisms of favorable effect

Several mechanisms are likely to be involved in the beneficial effects of vagal stimulation. The first question is as follows: how important is the contribution of the heart rate decrease? In the arrhythmia literature, this issue has been analyzed in detail [43]. Briefly, although few studies indicate it as the sole cause of the antiarrhythmic effect, the majority of the investigations, including our own [18, 21, 22], suggest that heart rate modulation plays a significant but not exclusive role. It should be mentioned that several studies were performed during coronary artery occlusion, a condition in which higher heart rates are accompanied by greater degrees of myocardial ischemia [11, 44], thus obviously facilitating arrhythmias. However, the experience with isolated ventricular arrhythmias in patients in the absence of ischemia [30] is still consistent with this concept.

Among patients with heart failure, a heart-rate-mediated effect is likely to play a significant role. Results from clinical trials have suggested that the heart rate decrease during beta-blocker treatment is a major determinant of the favorable effect of this pharmacological approach [45]. The recently reported beneficial effect of ivabradine, an agent acting almost exclusively by decreasing heart rate, also supports the concept of a favorable role of heart rate reduction [46]. Experimentally, heart rate during vagal stimulation was allowed to decrease in the first two models used [31–33], thus not allowing the assessment of a heart-rate-independent mechanism. However, in two recent studies, namely in dogs subjected to the same high-rate ventricular pacing in both vagal stimulation and control group [34] and in dogs with microembolization-induced HF subjected to vagus nerve stimulation at a low intensity [35], beneficial effects were found in the absence of any heart rate difference.

Several of the potential additional mechanisms contributing to the beneficial effect of vagal stimulation have been recently discussed in detail [47]. These may include anti-adrenergic effects that may occur at central level and at peripheral level (both pre-synaptic and post-synaptic), anti-apoptotic effects, an increase in nitric oxide and anti-inflammatory effects. Indeed, these intriguing additional mechanisms are related to the so-called inflammatory reflex, which shows that vagal stimulation inhibits macrophage activation, the synthesis of TNF, and inflammatory responses [48]. This particular effect is likely to be

involved also in the beneficial results of vagal stimulation against myocardial ischemia/reperfusion injury.

### Present challenges: identification of the best stimulation technique and the ideal patient

The demonstration of a beneficial effect of vagal stimulation in patients with heart failure clearly needs confirmation in a large randomized controlled clinical trial. At least two challenges exist: selection of the stimulation protocol and selection of the patient. It should be emphasized that simply stating that a patient will be assigned to vagus nerve stimulation therapy is very inexact. A list of more than ten parameters that may affect the clinical efficacy of the intervention is presented in the Table 1.

While most pharmacological interventions show a dose–response curve (with some doses ineffective or even detrimental), the same likely applies to neural modulation interventions such as vagal stimulation. To date, experimental studies have not adequately addressed this issue; thus, choices have to be made on the basis of the limited clinical experience.

Equally important is the selection of patients who are likely to respond better to this autonomic intervention. Again, most interventions in patients with heart failure show variable efficacy, and a non-negligible percentage of patients are generally considered to be non-responders. For instance, among candidates to cardiac resynchronization therapy, identification of responders and non-responders (still an open issue) would clearly allow the best use of resources and avoid unnecessary implantation of biven-tricular devices. Very limited data are available so far to

**Table 1** Vagus nerve stimulation parameter that may be regulated

Variables in vagal nerve stimulation
Right vs left vs bilateral stimulation
Electrode and waveform configuration
Bidirectional efferent and afferent (technically easier), preferential efferent or preferential afferent stimulation (technically more complex)
Continuous stimulation versus pulse-synchronous stimulation
With pulse-synchronous stimulation: delay from the R wave (or other trigger) and number of pulses per cycle
With continuous stimulation: frequency of stimulation
Current amplitude, titration protocols, and maximum current
Target: heart rate reduction vs low-intensity vagal stimulation without heart rate target
Duration of the ON/OFF cycles
Presence or absence of long stimulation pauses
Heart-rate-dependent stimulation intensity and limits for stimulation withdrawal (e.g., low heart rate)



help identify the ideal candidate for vagus nerve stimulation. It appears that patients who, despite optimal medical therapy, still show signs of marked sympathetic activation such as higher heart rate and lower heart rate variability tend to show a better response [49]. Patients with diabetic neuropathy were excluded our study, but the question whether any type of diabetes reduces the responsiveness to vagal stimulation is still open. In addition, it appears reasonable that patients who present a very large LV scar and thus have a limited amount of viable tissue are unlikely to show a marked improvement in LV volumes and ejection fraction.

### **Future directions: possible alternative electrical stimulation techniques leading to an augmentation of vagal efferent activity**

A shift in the autonomic balance among subjects with heart disease toward reduced sympathetic drive and increased vagal activity can be obtained both experimentally and clinically by several means including exercise training [50, 51] and pharmacologic modulation [52, 53] as detailed elsewhere in this issue of HF Reviews [54]. Here, we briefly focus on electrical stimulation techniques other than direct vagus nerve stimulation that can lead to increased vagal efferent activity.

Following the experimental demonstration in rats that electroneurostimulation of the ear endings and brainstem structures increases parasympathetic tone and exerts an antiarrhythmic effect [55], Zamotirsky et al. [56, 57] performed clinical studies using acupuncture needles placed at a depth of 0.1–0.3 mm into an acupuncture point known as heart 1 (shin), corresponding to an area near the auditory passage that contains endings of the nerve auricularis, a sensitive ramus of the vagus nerve. The treatment consisted of one daily application of low-intensity current for 15 min. Both studies involved patients with advanced coronary artery disease and severe angina. The first study [56] found a striking reduction in angina and the absence, in atrial tissue taken during the subsequent surgical revascularization intervention, of inducible heat-shock protein, present in all control patients not subjected to electrostimulation. The second paper [57] confirmed and extended these findings, also showing a significant decrease in heart rate and blood pressure, as well as an improvement in LVEF and diastolic properties.

An additional method involves electrical stimulation of carotid baroreceptors. This possibility was first introduced in the 1960s [58, 59] but was abandoned, mainly for technical reasons. More recently, an implantable device has been developed that may overcome most of these problems [60]. Chronic stimulation with this device has been shown

to produce sustained hypotension experimentally [61] and to significantly decrease blood pressure in drug-resistant hypertensive patients in open-label investigations [62]. Although carotid baroreceptor stimulation likely increases significantly vagal efferent activity, the main mechanism for blood pressure reduction may be sympathetic inhibition. Indeed, Heusser et al. [63] measured muscle sympathetic nerve activity by microneurography in 12 hypertensive patients during carotid baroreceptor stimulation and found that sympathetic nerve activity decreased markedly during stimulation and that this decrease significantly correlated with the blood pressure decrease. Experimentally, in a canine model of pacing-induced heart failure, chronic baroreceptor activation reduced neurohormonal activation and prolonged survival from an average of 37 days in the control group to an average of 68 days [64].

A detailed description of the effects of spinal cord stimulation (SCS) goes beyond the scope of this paper. However, it should be mentioned that experimentally SCS has been shown to enhance parasympathetic activity, mediated via the vagus nerve [65]. This technique was shown to reduce ventricular arrhythmia and to markedly improve LV function in a canine post-infarction heart failure model over a several week period [66].

Vagal stimulation can also be accomplished by means of intravascular catheters. Schauerte et al. [67] have shown that cardiac parasympathetic nerves can be stimulated in conscious humans via a multipolar catheter placed in the superior vena cava or in the ostium of the coronary sinus (stimulating during the atrial refractory period to avoid inducing atrial fibrillation). The two distinct sites predominantly slowed sinus rate and AV conduction, respectively. Stimulation duration was very short, and patients perceived pain, particularly during stimulation in the superior vena cava. Despite these limitations, this approach appears interesting. For instance, atrial pacing leads implanted in appropriate areas of the right atrium may provide safe high-frequency stimulation recruiting vagal efferent fibers and slowing the ventricular rate during atrial fibrillation in patients [68].

### **Future directions: potential application in other clinical settings**

At a time when the possibility of chronic, safe, and effective stimulation of the vagus nerve in cardiac patients is coming to reality, it should be emphasized that cardiac conditions other than heart failure may benefit from this technique. Given the wealth of experimental data on the antiarrhythmic potential of vagal stimulation (discussed above), an obvious clinical application may be the prevention of life-threatening arrhythmias in patients at high

risk, particularly if due to ischemic heart disease or the treatment of recurrent episodes of ventricular arrhythmias leading to multiple ICD shocks. In addition, following the pioneering experience of Braunwald [59], vagal stimulation may help patients with intractable angina pectoris. Indeed, the two more recent studies by Zamotrinsky et al. [56, 57] with electroneurostimulation of the ear endings (see above) suggest a striking anti-anginal effect among severely symptomatic patients.

Finally, as discussed earlier in this paper, myocardial ischemia/reperfusion injury may be attenuated by vagal stimulation. Multiple experimental studies demonstrate a beneficial effect of vagal stimulation in the reduction of infarct size following myocardial ischemia and a marked protection against myocardial ischemia/reperfusion injury [69, 70]. At least two clinical settings may be relevant. The first is related to myocardial reperfusion in patients undergoing surgical procedures such as coronary artery bypass grafting (CABG) and heart transplantation. Preliminary canine experimental studies have been performed to assess whether controlled intermittent asystole induced by high-intensity vagal stimulation, potentially useful to facilitate off-pump CABG, would impair post-asystolic coronary blood flow or myocardial function [71]. The authors found no negative acute effect of intermittent asystole; rather, a significant decrease in neutrophil accumulation in the ischemic myocardium was observed, in agreement with the hypothesis of vagally mediated decrease in reperfusion injury. Similarly, other situations requiring transient asystole, such as robotic CABG, endovascular deployment of aortic stents or valves might be applicable.

Reperfusion in the setting of acute MI could also be approached with vagal stimulation, possibly performed by means of an intravascular catheter [67] at the time of primary PCI, resulting in the activation of a protective pathway and in the reduction in myocardial damage following reperfusion.

Additional future applications of chronic vagal stimulation in primarily non-cardiologic conditions are conceivable. For instance, a case can be made for its use in patients with chronic renal failure. These patients have a poor prognosis characterized by a markedly increased cardiovascular mortality. Cardiovascular autonomic dysfunction has been proposed as a major contributing factor [72]. Interestingly, recent experimental data in rats subjected to surgical nephrectomy leading to chronic renal failure found marked autonomic imbalance characterized by diminished cardiac parasympathetic tone but by a completely preserved cardiac response to direct electrical vagus nerve stimulation [73]. Thus, vagus nerve stimulation might help to restore the autonomic imbalance and reduce the unacceptably high cardiovascular risk among this group of patients.

## Conclusions

Vagal stimulation in patients with heart failure has a strong physiologic and experimental rationale and has been recently proven to be feasible and safe. Preliminary data suggest that this intervention provides subjective and objective improvements. These results require appropriate evaluation in randomized controlled clinical trials of adequate size. Should these trials confirm the promising results of the open-label evaluation, vagus nerve stimulation is likely to become an additional important tool for the treatment of patients with heart failure.

As an example of the potential of this new therapeutic strategy, a patient with chronic ischemic heart disease, a reduction in systolic LV function and the persistence of coronary stenoses, and angina might be a candidate for a vagal nerve stimulation device, possibly combined with defibrillator capacity for primary prevention. Indeed, in such a patient, chronic low-intensity vagal stimulation may provide an improvement in LV function, functional capacity, and quality of life, as well as reduce the risk of ventricular tachycardia/fibrillation. Angina, if present, might also respond favorably. In the case of myocardial ischemia (detected by a device sensor or eventually by the patient), the intensity of vagal stimulation could be increased by appropriate programming, thereby providing protection against ischemic arrhythmias, a reduction in the severity of ischemia, and, in case of prolonged ischemia, also in the amount of ischemia/reperfusion injury. Finally, the device could also provide higher-intensity vagal stimulation to reduce ventricular rate in case of fast ventricular response to atrial tachycardia/fibrillation.

This scenario may appear futuristic, but the same criticism could have been made to our statement, more than 15 years ago [43], that “the development of a chronically implantable electrode” was “within current technical possibilities”.

Therefore, although much clinical research is still required, we believe we are now at the beginning of the clinical use of electrical neuromodulation for the treatment of patients with heart failure and other related cardiac conditions.

## References

1. Einbrodt (1859) Ueber Herzreizung und ihr Verhaeltnis zum Blutdruck. Akademie der Wissenschaften (Vienna). Sitzungsberichte 38:345
2. Hering HE (1901) Die Myoerethischen Unregelmassigkeiten des Herzens. Praeger Medizinische Wochenschrift 26:7
3. Scherf D (1929) Untersuchungen ueber die Entstehungsweise der Extrasystolen und der extrasystolischen Allorhythmien. Z Exp Med 65:198

4. Scherf D, Blumenfeld S, Yildiz M (1961) Experimental study on ventricular extrasystoles provoked by vagal stimulation. *Am Heart J* 62:670–675
5. Garrey WE (1908) Some effects of cardiac nerves upon ventricular fibrillation. *Am J Physiol* 21:283–300
6. Shumacker HB Jr, Riberi A, Boone RD, Kajikuri H (1956) Ventricular fibrillation in the hypothermic state. IV. The role of extrinsic cardiac innervation. *Ann Surg* 143:223–229
7. Scherlag BJ, Helfant RH, Haft JJ, Damato AN (1970) Electrophysiology underlying ventricular arrhythmias due to coronary ligation. *Am J Physiol* 219:1665–1671
8. Kerzner J, Wolf M, Kosowsky BD, Lown B (1973) Ventricular ectopic rhythms following vagal stimulation in dogs with acute myocardial infarction. *Circulation* 47:44–50
9. Kent KM, Smith ER, Redwood DR, Epstein SE (1973) Electrical stability of acutely ischemic myocardium: influences of heart rate and vagal stimulation. *Circulation* 47:291–298
10. Goldstein RE, Karsh RB, Smith ER, Orlando M, Norman D, Farnham G, Redwood DR, Epstein SE (1973) Influence of atropine and of vagally mediated bradycardia on the occurrence of ventricular arrhythmias following acute coronary occlusion in closed-chest dogs. *Circulation* 47:1180–1190
11. Myers RW, Pearlman AS, Hyman RM, Goldstein RA, Kent KM, Goldstein RE, Epstein SE (1974) Beneficial effects of vagal stimulation and bradycardia during experimental acute myocardial ischemia. *Circulation* 49:943–947
12. Corr PB, Gillis RA (1974) Role of the vagus nerves in the cardiovascular changes induced by coronary occlusion. *Circulation* 49:86–97
13. Yoon MS, Han J, Tse WW, Rogers R (1977) Effects of vagal stimulation, atropine, and propranolol on fibrillation threshold of normal and ischemic ventricles. *Am Heart J* 93:60–65
14. Yoon MS, Fondacaro JD, Han J (1978) Effects of vagal stimulation and atropine on ventricular arrhythmias during acute coronary occlusion. *J Electrocardiol* 11:27–31
15. Kolman BS, Verrier RL, Lown B (1975) The effect of vagus nerve stimulation upon vulnerability of the canine ventricle: Role of the sympathetic-parasympathetic interactions. *Circulation* 52:578–585
16. Furey SA 3rd, Levy MN (1983) Interactions among heart rate, autonomic activity, and arterial pressure upon the multiple repetitive extrasystole threshold in the dog. *Am Heart J* 106:1112–1120
17. Murdock DK, Loeb JM, Euler DE, Randall WC (1980) Electrophysiology of coronary reperfusion. A mechanism for reperfusion arrhythmias. *Circulation* 61:175–182
18. Zuanetti G, De Ferrari GM, Priori SG, Schwartz PJ (1987) Protective effect of vagal stimulation on reperfusion arrhythmias in cats. *Circ Res* 61:429–435
19. Yellon DM, Hausenloy DJ (2007) Myocardial reperfusion injury. *N Engl J Med* 357:1121–1135
20. Schwartz PJ, Billman GE, Stone HL (1984) Autonomic mechanisms in ventricular fibrillation induced by myocardial ischemia during exercise in dogs with a healed myocardial infarction. An experimental preparation for sudden cardiac death. *Circulation* 69:790–800
21. De Ferrari GM, Vanoli E, Stramba-Badiale M, Hull SS Jr, Foreman RD, Schwartz PJ (1991) Vagal reflexes and survival during acute myocardial ischemia in conscious dogs with a healed myocardial infarction. *Am J Physiol* 261:H63–H69
22. Vanoli E, De Ferrari GM, Stramba-Badiale M, Hull SS Jr, Foreman RD, Schwartz PJ (1991) Vagal stimulation and prevention of sudden death in conscious dogs with a healed myocardial infarction. *Circ Res* 68:1471–1481
23. Schwartz PJ, Pagani M, Lombardi F, Malliani A, Brown AM (1973) A cardiocardiac sympathovagal reflex in the cat. *Circ Res* 32:215–220
24. De Ferrari GM, Salvati P, Grossoni M, Ukmar G, Vaga L, Patrono C, Schwartz PJ (1993) Pharmacologic modulation of the autonomic nervous system in the prevention of sudden cardiac death. A study with propranolol, methacholine and oxotremorine in conscious dogs with a healed myocardial infarction. *J Am Coll Cardiol* 22:283–290
25. Ando M, Katare RG, Kakinuma Y, Zhang D, Yamasaki F, Muramoto K, Sato T (2005) Efferent vagal nerve stimulation protects heart against ischemia-induced arrhythmias by preserving connexin43 protein. *Circulation* 112:164–170
26. Schwartz PJ, De Ferrari GM (2010) Sympathetic-parasympathetic interaction in health and disease: abnormalities and relevance in heart failure. *Heart Fail Rev* doi: [10.1007/s10741-010-9179-1](https://doi.org/10.1007/s10741-010-9179-1). Online First<sup>TM</sup>, 25 June 2010
27. Waxman MB, Downar E, Berman ND, Felderhof CH (1974) Phenylephrine (Neo-synephrine) terminated ventricular tachycardia. *Circulation* 50:656–664
28. Waxman MB, Wald RW (1977) Termination of ventricular tachycardia by an increase in cardiac vagal drive. *Circulation* 56:385–391
29. Waxman MB, Cupps CL, Cameron DA (1988) Modulation of an idioventricular rhythm by vagal tone. *J Am Coll Cardiol* 11:1052–1060
30. Facchini M, De Ferrari GM, Bonazzi O, Weiss T, Schwartz PJ (1991) Effect of reflex vagal activation on frequency of ventricular premature complexes. *Am J Cardiol* 68:349–354
31. Li M, Zheng C, Sato T, Kawada T, Sugimachi M, Sunagawa K (2004) Vagal nerve stimulation markedly improves long-term survival after chronic heart failure in rats. *Circulation* 109:120–124
32. Sabbah HN, Rastogi S, Mishra S, Gupta RC, Ilars I, Imai M, Cohen U, Ben-David T, Ben-Ezra O (2005) Long-term therapy with neuroselective electric vagus nerve stimulation improves LV function and attenuates global LV remodeling in dogs with chronic heart failure. (abstr) *Eur J Heart Fail* (Suppl) 4:166
33. Sabbah HN, Imai M, Zaretsky A, Rastogi S, Wang M, Jiang A, Zaca V (2007) Therapy with vagus nerve electrical stimulation combined with beta-blockade improves left ventricular systolic function in dogs with heart failure beyond that seen with beta-blockade alone. (abstr) *Eur J Heart Fail* (Suppl 1) 6:114
34. Zhang Y, Popovic ZB, Bibevski S, Fakhry I, Sica DA, Van Wagoner DR, Mazgalev TN (2009) Chronic vagus nerve stimulation improves autonomic control and attenuates systemic inflammation and heart failure progression in a canine high-rate pacing model. *Circ Heart Fail* 2:692–699
35. Sabbah HN, Wang M, Jiang A, Ruble SB, Hamann JJ (2010) Right vagus nerve stimulation improves left ventricular function in dogs with heart failure. (abstr) *J Am Coll Cardiol* (Suppl) 55:A16.E151
36. Ben-Menachem E (2001) Vagus nerve stimulation, side effects, and long-term safety. *J Clin Neurophysiol* 18:415–418
37. Schachter SC (2002) Vagus nerve stimulation therapy summary: five years after FDA approval. *Neurology* 59:S15–S20
38. Uthman BM, Reichl AM, Dean JC, Eisenschenk S, Gilmore R, Reid S, Roper SN, Wilder BJ (2004) Effectiveness of vagus nerve stimulation in epilepsy patients, a 12-year observation. *Neurology* 63:1124–1126
39. Nemeroff CB, Mayberg HS, Kahl SE, McNamara J, Frazer A, Henry TR, George MS, Charney DS, Brannan SK (2006) VNS therapy in treatment-resistant depression: clinical evidence and putative neurobiological mechanisms. *Neuropsychopharmacology* 31:1345–1355
40. Shuchman M (2007) Approving the vagus-nerve stimulator for depression. *N Engl J Med* 356:1604–1607
41. Schwartz PJ, De Ferrari GM, Sanzo A, Landolina M, Rordorf R, Raineri C, Campana C, Revera M, Ajmone-Marsan N, Tavazzi L, Otero A (2008) Long term vagal stimulation in patients with



- advanced heart failure: first experience in man. *Eur J Heart Fail* 10:884–891
42. De Ferrari GM, Crijns HJGM, Borggrefe M, Milasinovic G, Smid J, Zabel M, Gavazzi A, Sanzo A, Dennert R, Kuschyk J, Raspopovic S, Klein H, Swedberg K, Schwartz PJ (2010) Chronic vagus nerve stimulation: a new and promising therapeutic approach for chronic heart failure. *Eur Heart J*. doi: [10.1093/eurheartj/ehq391](https://doi.org/10.1093/eurheartj/ehq391)
  43. De Ferrari GM, Vanoli E, Schwartz PJ (1994) Vagal activity and ventricular fibrillation. In: Levy MN, Schwartz PJ (eds) Vagal control of the heart: experimental basis and clinical implications. Futura Publishing Co, Armonk, NY, pp 613–636
  44. Maroko PR, Kjekshus JK, Sobel BE, Watanabe T, Cowell JW, Ross J Jr, Braunwald E (1971) Factors influencing infarct size following experimental coronary artery occlusions. *Circulation* 43:67–82
  45. Lechat P, Hulot J-S, Escolano S, Mallet A, Leizorovicz A, Werhlen-Grandjean M, Pochmalicki G, Dargie H, on behalf of the CIBIS II Investigators (2001) Heart rate and cardiac rhythm relationships with bisoprolol benefit in chronic heart failure in CIBIS II Trial. *Circulation* 103:1428–1433
  46. Swedberg K, Komajda M, Böhm M, Borer JS, Ford I, Dubost-Brama A, Lerebours G, Tavazzi L, on behalf of the SHIFT Investigators (2010) Ivabradine and outcomes in chronic heart failure (SHIFT): a randomised placebo-controlled study. *Lancet* 376:875–885
  47. Olshansky B, Sabbah HN, Hauptman PJ, Colucci WS (2008) Parasympathetic nervous system and heart failure. Pathophysiology and potential implications for therapy. *Circulation* 118:863–871
  48. Tracey KJ (2002) The inflammatory reflex. *Nature* 420:853–859
  49. Borggrefe M, De Ferrari GM, Crijns HJ, Milasinovic G, Klein H, Zabel M, Gavazzi A, Kuschyk J, Schoene A, Sanzo A, Dennert R, Raspopovic S, Smid J, Iacovoni A, Schwartz PJ (2009) Chronic vagus nerve stimulation in patients with heart failure: potential predictors of success. (abstr) *Circulation (Suppl)* 120:S911
  50. Billman GE, Schwartz PJ, Stone HL (1984) The effects of daily exercise on susceptibility to sudden cardiac death. *Circulation* 69:1182–1189
  51. La Rovere MT, Bersano C, Gnemmi M, Specchia G, Schwartz PJ (2002) Exercise-induced increase in baroreflex sensitivity predicts improved prognosis after myocardial infarction. *Circulation* 106:945–949
  52. De Ferrari GM, Mantica M, Vanoli E, Hull SS Jr, Schwartz PJ (1993) Scopolamine increases vagal tone and vagal reflexes in patients after myocardial infarction. *J Am Coll Cardiol* 22:1327–1334
  53. La Rovere MT, De Ferrari GM (1995) New potential uses for transdermal scopolamine (hyoscine). *Drugs* 50:769–776
  54. Desai MY, Watanabe M, Laddu A, Hauptman PJ (2010) Pharmacologic modulation of parasympathetic activity in heart failure. *Heart Fail Rev*
  55. Meerson FZ, Radzievskii SA, Vorontsova E, Golubeva L, Chuvpi'Iskaia LM (1989) Possibility of reflex activation of stress-limiting systems and the prevention of arrhythmia by electroacupuncture. *Kardiologiya* 29:88–90
  56. Zamotinsky A, Afanasiev S, Karpov RS, Cherniavsky A (1997) Effects of electrostimulation of the vagus afferent endings in patients with coronary artery disease. *Coron Artery Dis* 8:551–557
  57. Zamotinsky AV, Kondratiev B, de Jong JW (2001) Vagal neurostimulation in patients with coronary artery disease. *Auton Neurosci* 88:109–116
  58. Schwartz SI, Griffith LS, Neistadt A, Hagfors N (1967) Chronic carotid sinus nerve stimulation in the treatment of essential hypertension. *Am J Sur* 114:5–15
  59. Braunwald E, Epstein SE, Glick G, Wechsler AS, Braunwald NS (1967) Relief of angina pectoris by electrical stimulation of the carotid sinus nerves. *N Eng J Med* 277:1278–1283
  60. Tordoi JH, Schmidli SI, Savolainen H, Liebeskind U, Hansky B, Herold U, Irwin E, Kroon AA, de Leeuw P, Peters TK, Kieval R, Cody R (2007) An implantable carotid sinus baroreflex activating system: surgical technique and short-term outcome from a multicenter feasibility trial for the treatment of resistant hypertension. *Eur J Vasc Endovasc Surg* 33:414–421
  61. Lohmeier TE, Irwin ED, Rossing MA, Serdar DJ, Kieval RS (2004) Prolonged activation of the baroreflex produces sustained hypotension. *Hypertension* 43:306–311
  62. Rheos hypertension system demonstrates feasibility; phase 3 trial under way. 30 March, 2009. <http://www.theheart.org/article/954051.do>. Accessed 10 May 2010
  63. Heusser K, Tank J, Engeli S, Diedrich A, Menne J, Eckert S, Peters T, Sweep FCGJ, Haller H, Pichlmaier AM, Luft FC, Jordan J (2010) Carotid baroreceptor stimulation, sympathetic activity, baroreflex function, and blood pressure in hypertensive patients. *Hypertension* 55:619–626
  64. Zucker IH, Hackley JF, Cornish KG, Hiser BA, Anderson NR, Kieval R, Irwin ED, Serdar DJ, Peuler JD, Rossing MA (2007) Chronic baroreceptor activation enhances survival in dogs with pacing-induced heart failure. *Hypertension* 50:904–910
  65. Olgin JE, Takahashi T, Wilson E, Verecke A, Steinberg H, Zipes DP (2002) Effects of thoracic spinal cord stimulation on cardiac autonomic regulation of the sinus and atrioventricular nodes. *J Cardiovasc Electrophysiol* 13:475–481
  66. Lopshire JG, Zhou X, Dusa C, Ueyama T, Rosenberger J, Courtney N, Ujhelyi M, Mullen T, Das M, Zipes DP (2009) Spinal cord stimulation improves ventricular function and reduces ventricular arrhythmias in a canine postinfarction heart failure model. *Circulation* 120:286–294
  67. Schauerte P, Mischke K, Plisiene J, Waldmann M, Zarse M, Stellbrink C, Schimpf T, Knackstedt C, Sinha A, Hanrath P (2001) Catheter stimulation of cardiac parasympathetic nerves in humans. A novel approach to the cardiac autonomic nervous system. *Circulation* 104:2430–2435
  68. Bianchi S, Rossi O, Della Scala A, Kornet L, Pulvirenti R, Monari G, Di Renzi P, Schauerte P, Azzolinini P (2009) Atrioventricular (AV) node vagal stimulation by transvenous permanent lead implantation to modulate AV node function: safety and feasibility in humans. *Heart Rhythm* 6:1282–1286
  69. Katare RG, Ando M, Kakinuma Y, Arikawa M, Handa T, Yamasaki F, Sato T (2009) Vagal nerve stimulation prevents reperfusion injury through inhibition of opening of mitochondrial permeability transition pore independent of the bradycardiac effect. *J Thorac Cardiovasc Surg* 137:223–231
  70. Uemura K, Zheng C, Li M, Kawada T, Sugimachi M (2010) Early short-term vagal nerve stimulation attenuates cardiac remodeling after reperfused myocardial infarction. *J Cardiac Fail* 16:689–699
  71. Ronson RS, Puskas JD, Thourani VH, Velez DA, Bufkin BL, Glass J, Guyton RA, Vinten-Johansen J (2003) Controlled intermittent asystole cardiac therapy induced by pharmacologically potentiated vagus nerve stimulation in normal and hibernating myocardium. *Ann Thorac Surg* 75:1929–1936
  72. Robinson TG, Carr SJ (2002) Cardiovascular autonomic dysfunction in uremia. *Kidney Int* 62:1921–1932
  73. Kuncová J, Švíglerová J, Kummer W, Rajdl D, Chottová-Dvořáková M, Tonar Z, Nalos L, Štengl M (2009) Parasympathetic regulation of heart rate in rats after 5/6 nephrectomy is impaired despite functionally intact cardiac innervation. *Nephrol Dial Transplant* 24:2362–2370