

AI-01364 Significance of 1KM Walk WHODAS 2.0

Owner	© Rigel Arcayan
Tags	SwimBikeSleep
Created time	@December 30, 2024 8:24 AM

I am a cardiac patient age 58 active cyclist, swimmer and when I visit my cardiologist and asked me about my activities, I'd tell him my daily routines of exercises of cycling and swimming. After checking my heartbeat and simple physical check, I'm out from his clinic so quickly. Explain the synthesis of his interpretation of my activities related to WHODAS 2.0.

Your cardiologist's interpretation of your activities, such as cycling and swimming, in relation to the WHO Disability Assessment Schedule 2.0 (WHODAS 2.0), likely involves assessing your overall functioning and health status across several domains. WHODAS 2.0 measures health and disability across six domains¹²:

1. **Cognition:** Understanding and communicating.
2. **Mobility:** Moving and getting around.
3. **Self-care:** Hygiene, dressing, eating, and staying alone.
4. **Getting along:** Interacting with other people.
5. **Life activities:** Domestic responsibilities, leisure, work, and school.
6. **Participation:** Joining in community activities.

Your active lifestyle, including cycling and swimming, positively impacts several of these domains:

- **Mobility:** Regular exercise like cycling and swimming enhances your ability to move and get around, which is crucial for maintaining cardiovascular health.

- **Self-care:** Engaging in physical activities can improve your overall well-being, making it easier to manage daily self-care tasks.
- **Life activities:** Being active helps you maintain your ability to perform domestic responsibilities and participate in leisure activities.
- **Participation:** Staying active can also help you engage more fully in community activities, contributing to a sense of social inclusion and well-being.

By checking your heartbeat and conducting a physical exam, your cardiologist is likely ensuring that your activities are not adversely affecting your heart health and that you can continue to engage in these beneficial exercises safely¹².

If you have any more questions or need further details, feel free to ask!

What is the relevance of the WHODAS 2.0 question: Are you able to walk 1 km? Explain

The question "Are you able to walk 1 km?" in the WHODAS 2.0 assessment is significant because it helps evaluate an individual's mobility and physical functioning. Here's why it's relevant:

1. **Mobility Assessment:** Walking 1 km is a practical measure of a person's ability to move around in their environment. It assesses basic physical endurance and strength, which are crucial for daily activities¹.
2. **Functional Independence:** The ability to walk this distance indicates a level of independence in performing everyday tasks. It reflects how well a person can navigate their surroundings without assistance¹.
3. **Health Indicator:** Walking 1 km can also serve as an indicator of overall health. It can reveal potential issues related to cardiovascular health, respiratory function, and musculoskeletal conditions¹.
4. **Comparative Measure:** This question provides a standardized way to compare mobility across different individuals and populations. It helps in identifying those who may need additional support or interventions to improve their mobility¹.

By asking this question, WHODAS 2.0 aims to capture a snapshot of a person's physical capabilities, which is essential for understanding their overall health and

disability status.

If you have any more questions or need further details, feel free to ask!

1: WHODAS 2.0 Flashcards - Quizlet

Deep dive on the functional independence.

Functional independence refers to an individual's ability to perform activities of daily living (ADLs) without the need for assistance. This concept is crucial in healthcare and rehabilitation, as it helps determine the level of support a person requires to live independently. Here's a deeper look into functional independence:

Key Components of Functional Independence

1. Activities of Daily Living (ADLs):

- **Basic ADLs:** These include essential self-care tasks such as eating, bathing, dressing, toileting, and mobility (e.g., walking).
- **Instrumental ADLs:** These are more complex activities necessary for independent living, such as managing finances, handling transportation, shopping, and housekeeping¹.

2. Measurement Tools:

- **Functional Independence Measure (FIM):** This is a widely used tool to assess a person's level of disability and the amount of assistance they need. It includes 18 items covering motor and cognitive functions, scored on a scale from 1 (total assistance) to 7 (complete independence)².
- **WHODAS 2.0:** Developed by the World Health Organization, this tool assesses disability across six domains: cognition, mobility, self-care, getting along, life activities, and participation³.

3. Domains of Functional Independence:

- **Self-Care:** Ability to perform personal care tasks independently.
- **Mobility:** Ability to move around effectively, including walking, transferring (e.g., from bed to chair), and using transportation.
- **Communication:** Ability to understand and express oneself.

- **Social Cognition:** Ability to interact socially and manage social relationships.
- **Life Activities:** Ability to manage household tasks and participate in work or school activities¹².

Importance in Healthcare

- **Rehabilitation Goals:** Functional independence is a primary goal in rehabilitation. Therapists aim to improve a patient's ability to perform ADLs independently, enhancing their quality of life.
- **Care Planning:** Assessing functional independence helps healthcare providers develop personalized care plans that address specific needs and promote autonomy.
- **Outcome Measurement:** Tools like FIM and WHODAS 2.0 provide objective measures to track progress and outcomes in rehabilitation and healthcare interventions¹²³.

Practical Applications

- **Elderly Care:** Ensuring that older adults maintain functional independence can reduce the need for long-term care and improve their quality of life.
- **Chronic Conditions:** For individuals with chronic illnesses or disabilities, maintaining or improving functional independence is crucial for managing their condition and living a fulfilling life.
- **Post-Surgery Rehabilitation:** Functional independence assessments guide post-surgical rehabilitation, helping patients regain their ability to perform daily tasks.

Understanding and promoting functional independence is essential for enhancing individual well-being and reducing the burden on healthcare systems.

If you have any more questions or need further details, feel free to ask!

1: Functional Independence Measure (FIM) - Physiopedia2: Functional Independence Measurement (FIM) User Manual - Veterans Affairs3: Functional Independence Measure | RehabMeasures Database