

# PMS – Exercise Sheet 8

## Exercise 1

### Initialisierung

$$A = \begin{pmatrix} 1 & 1 & 4 \\ 1 & 1 & 1 \\ 4 & 1 & 1 \end{pmatrix}$$

$$b = \begin{pmatrix} 6 \\ 6 \\ 6 \end{pmatrix}$$

$$x_0 = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

### Erste Iteration

$$d_0 = b - A * x_0 = \begin{pmatrix} 6 \\ 6 \\ 6 \end{pmatrix} - \begin{pmatrix} 1 & 1 & 4 \\ 1 & 1 & 1 \\ 4 & 1 & 1 \end{pmatrix} * \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 6 \\ 6 \\ 6 \end{pmatrix}$$

$$r_0 = d_0 = \begin{pmatrix} 6 \\ 6 \\ 6 \end{pmatrix}$$

$$\alpha_0 = \frac{r_0^T * r_0}{d_0^T * A * d_0} = \frac{\begin{pmatrix} 6 \\ 6 \\ 6 \end{pmatrix}^T * \begin{pmatrix} 6 \\ 6 \\ 6 \end{pmatrix}}{\begin{pmatrix} 6 \\ 6 \\ 6 \end{pmatrix}^T * \begin{pmatrix} 1 & 1 & 4 \\ 1 & 1 & 1 \\ 4 & 1 & 1 \end{pmatrix} * \begin{pmatrix} 6 \\ 6 \\ 6 \end{pmatrix}} = \frac{1}{5}$$

$$x_1 = x_0 + \alpha_0 d_0 = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} + \frac{1}{5} \begin{pmatrix} 6 \\ 6 \\ 6 \end{pmatrix} = \begin{pmatrix} \frac{6}{5} \\ \frac{6}{5} \\ \frac{6}{5} \end{pmatrix}$$

### Zweite Iteration

$$r_1 = r_0 - \alpha_0 A * d_0 = \begin{pmatrix} 6 \\ 6 \\ 6 \end{pmatrix} - \frac{1}{5} \begin{pmatrix} 1 & 1 & 4 \\ 1 & 1 & 1 \\ 4 & 1 & 1 \end{pmatrix} * \begin{pmatrix} 6 \\ 6 \\ 6 \end{pmatrix} = \begin{pmatrix} -\frac{6}{5} \\ \frac{12}{5} \\ -\frac{6}{5} \end{pmatrix}$$

$$\beta_1 = \frac{r_1^T r_1}{r_0^T r_0} = \frac{\begin{pmatrix} -\frac{6}{5} \\ \frac{12}{5} \\ -\frac{6}{5} \end{pmatrix}^T * \begin{pmatrix} -\frac{6}{5} \\ \frac{12}{5} \\ -\frac{6}{5} \end{pmatrix}}{\begin{pmatrix} 6 \\ 6 \\ 6 \end{pmatrix}^T * \begin{pmatrix} 6 \\ 6 \\ 6 \end{pmatrix}} = \frac{2}{25}$$

$$d_1 = r_1 + \beta_1 d_0 = \begin{pmatrix} -\frac{6}{5} \\ \frac{12}{5} \\ -\frac{6}{5} \end{pmatrix} + \frac{2}{25} \begin{pmatrix} 6 \\ 6 \\ 6 \end{pmatrix} = \begin{pmatrix} -\frac{18}{25} \\ \frac{72}{25} \\ -\frac{18}{25} \end{pmatrix}$$

$$\alpha_1 = \frac{r_1^T * r_1}{d_1^T * A * d_1} = \frac{\begin{pmatrix} -\frac{6}{5} \\ \frac{12}{5} \\ -\frac{6}{5} \end{pmatrix}^T * \begin{pmatrix} -\frac{6}{5} \\ \frac{12}{5} \\ -\frac{6}{5} \end{pmatrix}}{\begin{pmatrix} -\frac{18}{25} \\ \frac{72}{25} \\ -\frac{18}{25} \end{pmatrix}^T * \begin{pmatrix} 1 & 1 & 4 \\ 1 & 1 & 1 \\ 4 & 1 & 1 \end{pmatrix} * \begin{pmatrix} -\frac{18}{25} \\ \frac{72}{25} \\ -\frac{18}{25} \end{pmatrix}} = \frac{5}{3}$$

$$x_2 = x_1 + \alpha_1 d_1 = \begin{pmatrix} 6 \\ 6 \\ 6 \end{pmatrix} + \frac{5}{3} \begin{pmatrix} -\frac{18}{25} \\ \frac{72}{25} \\ -\frac{18}{25} \end{pmatrix} = \begin{pmatrix} 0 \\ 6 \\ 0 \end{pmatrix}$$

$$r_2 = r_1 - \alpha_1 A * d_1 = \begin{pmatrix} -\frac{6}{5} \\ \frac{12}{5} \\ -\frac{6}{5} \end{pmatrix} - \frac{5}{3} \begin{pmatrix} 1 & 1 & 4 \\ 1 & 1 & 1 \\ 4 & 1 & 1 \end{pmatrix} * \begin{pmatrix} -\frac{18}{25} \\ \frac{72}{25} \\ -\frac{18}{25} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

Abbruch, da Residuum  $r_2 = 0 \Rightarrow$  keine weitere Änderung an gefundenem Optimum  $x_2 = \begin{pmatrix} 0 \\ 6 \\ 0 \end{pmatrix}$

## Exercise 2

- As each step of the algorithm finds the local minimum of one dimension, for a 3-dimensional systems it will at most take three iterations.
- In large systems, the conjugate gradient algorithm allows for fast approximation, doing only a few iterations until the solution is "good enough". Gauss elimination does not allow this. TODO: More?
- $A$  has to be a symmetric, positive-definite matrix. All symmetric matrices are also square.
- TODO