

# STATS 205: Final Project Write-Up

Brian Liu

6/14/2019

## 1. Background of the data and why it is interesting or important

The data we are using is the data from WHO suicide statistics from Kaggle. This gives population-based statistics on suicide rate...

## 2. Explanation of the method studied and its properties

## 3. Data analysis or simulation study

We will use the crude rate of suicide per 100,000 people.

This analysis provides information on age-standardized rates...

```
who_suicide_statistics_df <- read.csv("who_suicide_statistics.csv")
head(who_suicide_statistics_df)
```

```
##   country year    sex      age suicides_no population
## 1 Albania 1985 female 15-24 years         NA      277900
## 2 Albania 1985 female 25-34 years         NA      246800
## 3 Albania 1985 female 35-54 years         NA      267500
## 4 Albania 1985 female  5-14 years         NA      298300
## 5 Albania 1985 female 55-74 years         NA      138700
## 6 Albania 1985 female  75+ years         NA       34200
```

```
colnames(who_suicide_statistics_df)
```

```
## [1] "country"    "year"       "sex"        "age"        "suicides_no"
## [6] "population"
```

Filter and save countries with missing suicide rate.

```
library(tidyverse)
```

```
## Registered S3 methods overwritten by 'ggplot2':
```

```
##   method      from
## [.quosures   rlang
## c.quosures   rlang
## print.quosures rlang
```

```
## -- Attaching packages ----- tidyverse 1.2.1 --
```

```
## v ggplot2 3.1.1    v purrr   0.3.2
## v tibble  2.1.1    v dplyr  0.8.1
## v tidyr   0.8.3    v stringr 1.4.0
## v readr   1.3.1    v forcats 0.4.0
```

```
## -- Conflicts ----- tidyverse_conflicts() --
```

```
## x dplyr::filter() masks stats::filter()
## x dplyr::lag()    masks stats::lag()
```

```
filtered_suicide_df <- drop_na(who_suicide_statistics_df, "suicides_no")
head(filtered_suicide_df)
```

```
##   country year  sex      age suicides_no population
## 25 Albania 1987 female 15-24 years         14    289700
## 26 Albania 1987 female 25-34 years          4    257200
## 27 Albania 1987 female 35-54 years          6    278800
## 28 Albania 1987 female  5-14 years          0    311000
## 29 Albania 1987 female 55-74 years          0    144600
## 30 Albania 1987 female  75+ years          1     35600
```

After filtering countries with missing suicide rate, take a random sample of 100 countries and make sure each continent has approximately equal countries.

$$\frac{100 \text{ countries}}{7 \text{ continents}} \approx 14 \text{ to } 15 \text{ countries per continent}$$

Filter countries by continent:

```
library(countrycode)
filtered_suicide_df$continent <- countrycode(sourcevar = filtered_suicide_df[, "country"],
                                             origin = "country.name",
                                             destination = "continent")
```

```
## Warning in countrycode(sourcevar = filtered_suicide_df[, "country"], origin = "country.name", : Some
```

```
## Warning in countrycode(sourcevar = filtered_suicide_df[, "country"], origin = "country.name", : Some
```

```
head(filtered_suicide_df)
```

```
##   country year  sex      age suicides_no population continent
## 25 Albania 1987 female 15-24 years         14    289700    Europe
## 26 Albania 1987 female 25-34 years          4    257200    Europe
## 27 Albania 1987 female 35-54 years          6    278800    Europe
## 28 Albania 1987 female  5-14 years          0    311000    Europe
## 29 Albania 1987 female 55-74 years          0    144600    Europe
## 30 Albania 1987 female  75+ years          1     35600    Europe
```

```
write.csv(filtered_suicide_df, 'filtered_suicide.csv')
```

```
# Get seven dataframes, filtered by list of countries for each continent.
```

```
europe_suicide = filtered_suicide_df[filtered_suicide_df$country == 'Europe']
head(europe_suicide)
```

```
## data frame with 0 columns and 6 rows
```

## 4. Interpretation of the results or discussion