

STATS 205: Homework Assignment 5

Brian Liu

6/10/2019

Solution to Problem 1

We say that two observations X_1 and X_2 are *independent* of one another with respect to a collection of events \mathcal{A} if

$$Pr \{X_1 \in A \text{ and } X_2 \in B\} = Pr \{X_1 \in A\} Pr \{X_2 \in B\}$$

where A and B are any two not necessarily distinct sets of outcomes belonging to \mathcal{A}^3 .

– 2.2.1 Independent Observations; Permutation, Parametric, and Bootstrap Tests of Hypotheses; Good, Phillip I

In deciding whether your own observations are exchangeable and a permutation test applicable, the key question is the one we posed in the very first chapter: Under the null hypothesis of no differences among the various experimental or survey groups, can we exchange the labels on the observations without significantly affecting the results?

– 2.2.2 Exchangeable Observations; Permutation, Parametric, and Bootstrap Tests of Hypotheses; Good, Phillip I

Solution to Problem 2

```
cysticerici <- c(28.9, 32.8, 12.0, 9.9, 15.0, 38.0, 12.5, 36.5, 8.6, 26.8);cysticerici
```

```
## [1] 28.9 32.8 12.0 9.9 15.0 38.0 12.5 36.5 8.6 26.8
```

```
worms_reco <- c(1.0, 7.7, 7.3, 7.9, 1.1, 3.5, 18.9, 33.9, 28.6, 25.0); worms_reco
```

```
## [1] 1.0 7.7 7.3 7.9 1.1 3.5 18.9 33.9 28.6 25.0
```

The null hypothesis is that the mean weight of introduced cysticerici *has no correlation with* the mean weight of worms recovered. That is,

$$H_0 : \tau = 0$$

The alternative hypothesis is that the mean weight of introduced cysticerici is *positively correlated with* the mean weight of worms recovered. That is,

$$H_A : \tau > 0$$

To test the null hypothesis against the alternative hypothesis, we will use the Kendall test, a distribution-free test for independence based on signs.

```
cor.test(x = cysticerici, y = worms_reco, method = "kendall", alt = "greater")
```

```
##
```

```
## Kendall's rank correlation tau
```

```
##
```

```
## data: cysticerci and worms_reco
## T = 19, p-value = 0.7578
## alternative hypothesis: true tau is greater than 0
## sample estimates:
##      tau
## -0.1555556
```

The p -value is 0.7578, which is *not* significant at the $\alpha = 0.05$ level. There is *not enough* evidence that the mean weight of introduced cysticerci is *positively correlated with* the mean weight of worms recovered.

Solution to Problem 3

```
cysticerci <- c(28.9, 32.8, 12.0, 9.9, 15.0, 38.0, 12.5, 36.5, 8.6, 26.8)
worms_reco <- c(1.0, 7.7, 7.3, 7.9, 1.1, 3.5, 18.9, 33.9, 28.6, 25.0)
cor.test(x = cysticerci, y = worms_reco, method = "kendall", alt = "greater")
```

```
##
## Kendall's rank correlation tau
##
## data: cysticerci and worms_reco
## T = 19, p-value = 0.7578
## alternative hypothesis: true tau is greater than 0
## sample estimates:
##      tau
## -0.1555556
```

The estimate for $\tau = -0.1555556$.

Solution to Problem 4

Solution to Problem 5

```
cysticerci <- c(28.9, 32.8, 12.0, 9.9, 15.0, 38.0, 12.5, 36.5, 8.6, 26.8)
worms_reco <- c(1.0, 7.7, 7.3, 7.9, 1.1, 3.5, 18.9, 33.9, 28.6, 25.0)
```

The null hypothesis is that the mean weight of introduced cysticerci *has no correlation with* the mean weight of worms recovered. That is,

$$H_0 : r_s < r_{s,\alpha}$$

The alternative hypothesis is that the mean weight of introduced cysticerci is *positively correlated with* the mean weight of worms recovered. That is,

$$H_A : r_s \geq r_{s,\alpha}$$

Otherwise, do not reject.

To test the null hypothesis against the alternative hypothesis, we will use the Spearman test, a distribution-free test for independence based on ranks.

```
library(SuppDists)
qSpearman(p = 0.05, r = 10)
```

```
## [1] -0.5393939
```

Since $r_{s,\alpha} = -0.5393939$, we will reject the null hypothesis only if $r_s \geq -0.5393939$.

Calculating r_s ,

```
cor(x = cysticerici, y = worms_reco, method = "spearman")
```

```
## [1] -0.2
```

Since $r_s = -0.2$ and $r_{s,\alpha} = -0.5393939$, the statement $r_s \geq r_{s,\alpha}$ is *true*. Thus, we *reject* the null hypothesis. There is *sufficient* evidence that the mean weight of introduced cysticerici is *positively correlated with* the mean weight of worms recovered.

NOTE: At this point, I tried to use `cor.test()` with `method = "spearman"` but I got a different result than I expected, and I'm not sure why. Maybe I'm interpreting the output incorrectly?

```
cor.test(x = cysticerici, y = worms_reco, method = "spearman", alternative = "greater")
```

```
##  
## Spearman's rank correlation rho  
##  
## data: cysticerici and worms_reco  
## S = 198, p-value = 0.72  
## alternative hypothesis: true rho is greater than 0  
## sample estimates:  
## rho  
## -0.2
```

The p -value is 0.72, which is *not* significant at the $\alpha = 0.05$ level. There is *not enough* evidence that the mean weight of introduced cysticerici is *positively correlated with* the mean weight of worms recovered.

Solution to Problem 6

```
x = c(0, 5000, 10000, 15000, 20000, 25000, 30000, 100000)  
y = c(0.924, 0.988, 0.992, 1.118, 1.133, 1.145, 1.157, 1.357)
```

The null hypothesis is that the mean weight of introduced cysticerici *has no correlation with* the mean weight of worms recovered. That is,

$$H_0 : \beta = \beta_0 H_0 : \beta = 0$$


The alternative hypothesis is that the mean weight of introduced cysticerici is *positively correlated with* the mean weight of worms recovered. That is,

$$H_A : \beta > \beta_0 H_A : \beta > 0$$

```
library(NSM3)
```

```
## Loading required package: combinat  
##  
## Attaching package: 'combinat'
```

```
## The following object is masked from 'package:utils':
##
##      combn
## Loading required package: MASS
## Loading required package: partitions
## Loading required package: survival
## fANCOVA 0.5-1 loaded
## Registered S3 methods overwritten by 'ggplot2':
##   method      from
## [.quosures    rlang
## c.quosures    rlang
## print.quosures rlang
theil(x, y, alpha=0.05, beta.0=0, type = "u")
```



```
## Alternative: beta greater than 0
## C = 28, C.bar = 1, P = 0
## beta.hat = 0
## alpha.hat = 0.975
##
## 1 - alpha = 0.95 upper bound for beta:
## -Inf, 0
```