

Name: _____

Section: _____

R.I.T SCHOOL OF MATHEMATICAL SCIENCES

28 - First Order Applications with Laplace Transforms

MATH 211

1. A boat with a mass of 10 slugs is being towed at 3 feet per second. The tow rope is then cut and the motor, that exerts a force of 20 lb on the boat, is started one minute later. If the water exerts a retarding force that numerically equals twice the velocity, what is the velocity of the boat 3 minutes later?

$$mv' = F - kv, \quad v(0) = 3(60) = 180$$

$$mv' + kv = F$$

$$10v' + 2v = 20u(t-1)$$

$$5v' + v = 10u(t-1)$$

$$5[s\mathcal{L}\{v\} - v(0)] + \mathcal{L}\{v\} = \mathcal{L}\{10u(t-1)\}$$

$$5[s\mathcal{L}\{v\} - 180] + \mathcal{L}\{v\} = \frac{10e^{-s}}{s}$$

$$5s\mathcal{L}\{v\} - 900 + \mathcal{L}\{v\} = \frac{10e^{-s}}{s}$$

$$(5s+1)\mathcal{L}\{v\} = \frac{10e^{-s}}{s} + 900$$

$$\mathcal{L}\{v\} = e^{-s} \left[\frac{10}{s(5s+1)} \right]_{\star_1} + \frac{900}{5s+1}$$

$$\mathcal{L}\{v\} = e^{-s} \left[\frac{10}{s} - \frac{50}{5s+1} \right]_{\star_2} + \frac{180}{s+1/5}$$

$$v = \left[10 - 10e^{-(t-1)/5} \right] u(t-1) + 180e^{-t/5}$$

$$\star_1 \frac{10}{s(5s+1)} = \frac{A}{s} + \frac{B}{5s+1}$$

$$10 = 5As + A + Bs$$

$$5A + B = 0 \quad A = 10$$

$$50 + B = 0$$

$$B = -50$$

$$\frac{10}{s(5s+1)} = \frac{10}{s} + \frac{-50}{5s+1}$$

$$\star_2 F(s) = \frac{10}{s} - \frac{10}{s+1/5}$$

$$f(t) = 10 - 10e^{-t/5}$$

$$f(t-1) = 10 - 10e^{-(t-1)/5}$$

$$f(t-1)u(t-1) = \left[10 - 10e^{-(t-1)/5} \right] u(t-1)$$

2. Suppose that in a simple RL circuit the resistance is 10Ω and the inductance is 2 H . If a battery gives a voltage of $E(t) = \sin t \, u(t - \pi) \text{ V}$ and the switch is closed when $t = 0$ so the current starts at 3 mA , find the current as a function of time.

$$Li' + Ri = E(t)$$

$$2i' + 10i = \sin t \, u(t - \pi)$$

$$2[s\mathcal{L}\{i\} - i(0)] + 10\mathcal{L}\{i\} = \mathcal{L}\{\sin t \, u(t - \pi)\}_{\star_1}$$

$$2s\mathcal{L}\{i\} - 2\left(\frac{3}{1000}\right) + 10\mathcal{L}\{i\} = -e^{-\pi s} \cdot \frac{1}{s^2 + 1}$$

$$(2s + 10)\mathcal{L}\{i\} = -e^{-\pi s} \cdot \frac{1}{s^2 + 1} + \frac{3}{500}$$

$$\mathcal{L}\{i\} = -e^{-\pi s} \cdot \frac{1}{(s^2 + 1)(2s + 10)} + \frac{3}{500} \cdot \frac{1}{2s + 10}$$

$$\mathcal{L}\{i\} = -\frac{1}{2}e^{-\pi s} \cdot \frac{1}{(s^2 + 1)(s + 5)}_{\star_2} + \frac{3}{1000} \cdot \frac{1}{s + 5}$$

$$i = -\frac{1}{2} \cdot \frac{1}{26} \left[\cos t - 5 \sin t + e^{-5(t-\pi)} \right] u(t - \pi) + \frac{3}{1000} e^{-5t}$$

$$i = -\frac{1}{52} \left[\cos t - 5 \sin t + e^{-5(t-\pi)} \right] u(t - \pi) + \frac{3}{1000} e^{-5t}$$

$$\star_1 \, a = \pi, \, f(t) = \sin t$$

$$f(t + \pi) = \sin(t + \pi) = -\sin t$$

$$\mathcal{L}\{f(t + \pi)\} = -\mathcal{L}\{\sin t\} = -\frac{1}{s^2 + 1}$$

$$e^{-\pi s} \mathcal{L}\{f(t + \pi)\} = -e^{-\pi s} \cdot \frac{1}{s^2 + 1}$$

$$\star_2 \, a = \pi, \, F(s) = \frac{1}{(s^2 + 1)(s + 5)} = \frac{As + B}{s^2 + 1} + \frac{C}{s + 5}$$

$$1 = As^2 + 5As + Bs + 5B + Cs^2 + C$$

$$\begin{array}{lll} A + C = 0 & 5A + B = 0 & 5B + C = 1 \\ C = -A & B = -5A & 5(-5A) - A = 1 \\ & & -26A = 1 \\ & & A = -\frac{1}{26} \end{array}$$

$$C = \frac{1}{26} \quad B = \frac{5}{26}$$

$$F(s) = \frac{(-1/26)s + 5/26}{s^2 + 1} + \frac{1/26}{s + 5}$$

$$F(s) = \frac{1}{26} \left[-\frac{s}{s^2 + 1} + \frac{5}{s^2 + 1} + \frac{1}{s + 5} \right]$$

$$f(t) = \frac{1}{26} \left[-\cos t + 5 \sin t + e^{-5t} \right]$$

$$f(t - \pi) = \frac{1}{26} \left[-\cos(t - \pi) + 5 \sin(t - \pi) + e^{-5(t-\pi)} \right]$$

$$f(t - \pi) u(t - \pi) = \frac{1}{26} \left[\cos t - 5 \sin t + e^{-5(t-\pi)} \right] u(t - \pi)$$

3. Pure water flows at a constant rate of 5 liters per minute into a large tank that initially held 125 liters of brine solution in which was dissolved 1 kilogram of salt. The solution inside the tank is kept well stirred and flows out at the same rate. At $t = 5$ minutes, 4 kilograms of salt are instantaneously dumped into the tank. Find a formula for the amount of salt in the tank as a function of time t . How much salt is in the tank after 10 minutes?

$$\frac{dx}{dt} = 0 - \frac{5x}{125} + 4\delta(t - 5)$$

$$x' + \frac{1}{25}x = 4\delta(t - 5)$$

$$s\mathcal{L}\{x\} - x(0) + \frac{1}{25}\mathcal{L}\{x\} = 4\mathcal{L}\{\delta(t - 5)\}$$

$$s\mathcal{L}\{x\} - 1 + \frac{1}{25}\mathcal{L}\{x\} = 4e^{-5s}$$

$$\left(s + \frac{1}{25}\right)\mathcal{L}\{x\} = 4e^{-5s} + 1$$

$$\mathcal{L}\{x\} = 4e^{-5s} \cdot \frac{1}{s + 1/25} + \frac{1}{s + 1/25}$$

$$x = 4e^{-(t-5)/25} u(t-5) + e^{-t/25}$$

$$\star a = 5, F(s) = \frac{1}{s + 1/25}$$

$$f(t) = e^{-t/25}$$

$$f(t-5) = e^{-(t-5)/25}$$

$$f(t-5) u(t-5) = e^{-(t-5)/25} u(t-5)$$