

Question. What does math/doing math mean to you?

- Why do you want to study math?
- Do you like math? Why?
- How could you learn about math outside of school/VMT?
- What does being “good” at math mean?
- How diverse is the math community?
- Does doing math make you “smarter?”
- Why should people study math?

What's Next?

math stories from beyond VMT

Bryan Lu

June 5, 2023

Your Landscape

Competition math categories:

- algebra (inequalities, FEs)
- geometry (mostly Euclidean)
- combinatorics
- number theory (elementary)

Your Landscape

Competition math categories:

- algebra (inequalities, FEs)
- geometry (mostly Euclidean)
- combinatorics
- number theory (elementary)

TJ Math (engineering reqs):

- calculus (BC + multi)
- linear algebra
- differential equations
- statistics

Your Landscape

Competition math categories:

- algebra (inequalities, FEs)
- geometry (mostly Euclidean)
- combinatorics
- number theory (elementary)

TJ Math (engineering reqs):

- calculus (BC + multi)
- linear algebra
- differential equations
- statistics

This is **VERY FAR** from everything that's out there!

Further Fields of Study

Typical undergraduate categories:

Main categories of study:

- (abstract) algebra
- analysis/measure
- topology/geometry

Further Fields of Study

Typical undergraduate categories:

Main categories of study:

- (abstract) algebra
- analysis/measure
- topology/geometry

Less central categories:

- combinatorics
- logic/set theory
- probability/statistics

Further Fields of Study

Typical undergraduate categories:

Main categories of study:

- (abstract) algebra
- analysis/measure
- topology/geometry


Less central categories:

- combinatorics
- logic/set theory
- probability/statistics

Broader fields (incomplete/ill-defined list; historically important):

- | | | |
|---------------------------|-------------------------|------------------------|
| • algebraic number theory | • harmonic analysis | • algebraic geometry |
| • analytic number theory | • operator theory | • algebraic topology |
| • dynamical systems | • computer science | • foundations |
| • Lie theory | • game theory/economics | • mathematical physics |

Dependency Graph (*Napkin, Evan Chen*)



Myths/Misconceptions

- ➊ Proving really **hard** theorems/solving really **hard** problems (misplaced emphasis)

Myths/Misconceptions

- ① Proving really **hard** theorems/solving really **hard** problems (misplaced emphasis)
- ② Formalism vs. pedantry

Myths/Misconceptions

- ① Proving really **hard** theorems/solving really **hard** problems (misplaced emphasis)
- ② Formalism vs. pedantry
- ③ Working on your own vs. community of ideas

Myths/Misconceptions

- ① Proving really **hard** theorems/solving really **hard** problems (misplaced emphasis)
- ② Formalism vs. pedantry
- ③ Working on your own vs. community of ideas
- ④ Difficulty as a barrier to growth

Myths/Misconceptions

- ➊ Proving really **hard** theorems/solving really **hard** problems (misplaced emphasis)
- ➋ Formalism vs. pedantry
- ➌ Working on your own vs. community of ideas
- ➍ Difficulty as a barrier to growth
- ➎ (there are probably tons more . . .)

Core Ideas

- ① Connecting seemingly unrelated objects/procedures.

Core Ideas

- ① Connecting seemingly unrelated objects/procedures.
- ② Extending/generalizing ideas we already know.

Core Ideas

- ① Connecting seemingly unrelated objects/procedures.
- ② Extending/generalizing ideas we already know.
- ③ Analyzing when/why patterns break and seeking complexity.

Core Ideas

- ① Connecting seemingly unrelated objects/procedures.
- ② Extending/generalizing ideas we already know.
- ③ Analyzing when/why patterns break and seeking complexity.
- ④ Staying grounded with “standard” examples/calculations.

the rest of this lecture: choose your own adventure

- for each Core Idea I present two fields of study that I think exemplify this idea

the rest of this lecture: choose your own adventure

- for each Core Idea I present two fields of study that I think exemplify this idea
- read the descriptions of what I would talk to you about for each field of study I've selected – *a star means a subject is less accessible*

the rest of this lecture: choose your own adventure

- for each Core Idea I present two fields of study that I think exemplify this idea
- read the descriptions of what I would talk to you about for each field of study I've selected – *a star means a subject is less accessible*
- crowd vote!!

the rest of this lecture: choose your own adventure


- for each Core Idea I present two fields of study that I think exemplify this idea
- read the descriptions of what I would talk to you about for each field of study I've selected – *a star means a subject is less accessible*
- crowd vote!! ⇒ this talk could go in 8 possible ways

Idea 1 – Connections

- *How to Tell Spaces Apart (Simplicial Homology)* – We will use graphs as a model for describing various topological spaces (spheres, toruses, Klein bottles) and use an extension of the Euler characteristic with linear algebra to distinguish spaces from each other.
- (★) *Word Problems (Geometric Group Theory)* – We will take an alternate view of groups and study graphs induced by a group's structure, and use these graphs to explain why deciding whether an element in a group is the identity element is unsolvable.



1.1 – Euler Characteristic

Euler characteristic of a graph G with V vertices, E edges, and F faces:
 $\chi(G) = V - E + F$.



$$\chi(G) = 2$$

1.1 – Example Spaces



1.1 – Chain Complexes

$$\sigma = [v_0, v_1, \dots, v_n]$$

$$\Rightarrow \partial\sigma = \sum_{i=0}^n (-1)^i \sigma|_{[v_0, v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_n]}$$

1.1 – Chain Complexes

$$\sigma = [v_0, v_1, \dots, v_n]$$

$$\Rightarrow \partial\sigma = \sum_{i=0}^n (-1)^i \sigma|_{[v_0, v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_n]}$$

$$\cdots \rightarrow C_n(G) \xrightarrow{\partial_n} \cdots \rightarrow C_2(G) \xrightarrow{\partial_2} C_1(G) \xrightarrow{\partial_1} C_0(G) \xrightarrow{\partial_0} 0$$

Holes have no boundary, but are not the boundary of any simplex.

$$\boxed{\partial^2 = 0}$$

1.1 – Chain Complexes

$$\sigma = [v_0, v_1, \dots, v_n]$$

$$\Rightarrow \partial\sigma = \sum_{i=0}^n (-1)^i \sigma|_{[v_0, v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_n]}$$

$$\cdots \rightarrow C_n(G) \xrightarrow{\partial_n} \cdots \rightarrow C_2(G) \xrightarrow{\partial_2} C_1(G) \xrightarrow{\partial_1} C_0(G) \xrightarrow{\partial_0} 0$$

Holes have no boundary, but are not the boundary of any simplex.

$$\boxed{\partial^2 = 0} \Rightarrow \boxed{H_n(G) = \frac{\partial}{\partial}}$$

1.1 – Homology Groups

G	S^2	\mathbb{T}^2	$\mathbb{R}P^2$	K
$H_0(G)$	\mathbb{Z}	\mathbb{Z}	\mathbb{Z}	\mathbb{Z}
$H_1(G)$	0	\mathbb{Z}^2	$\mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$
$H_2(G)$	\mathbb{Z}	\mathbb{Z}	0	0

1.1 – Identical Homology

$$H_0 = \mathbb{Z} \quad H_1 = \mathbb{Z}^2 \quad H_2 = \mathbb{Z}$$
$$\mathbb{T}^2 \quad S^1 \vee S^1 \vee S^2$$

1.2 – Presentations

$$G = \langle S \mid R \rangle$$

1.2 – Presentations

$$G = \langle S \mid R \rangle$$

S – set of symbols S^{-1} – each symbol tagged with an inverse

1.2 – Presentations

$$G = \langle S \mid R \rangle$$

S – set of symbols S^{-1} – each symbol tagged with an inverse

$L(S \cup S^{-1})$ – all strings in letters of $S \cup S^{-1}$

1.2 – Presentations

$$G = \langle S \mid R \rangle$$

S – set of symbols S^{-1} – each symbol tagged with an inverse

$L(S \cup S^{-1})$ – all strings in letters of $S \cup S^{-1}$

R – words $w \in L(S \cup S^{-1})$ in $S \cup S^{-1}$ equal to e

1.2 – Presentations

$$G = \langle S \mid R \rangle$$

S – set of symbols S^{-1} – each symbol tagged with an inverse

$L(S \cup S^{-1})$ – all strings in letters of $S \cup S^{-1}$

R – words $w \in L(S \cup S^{-1})$ in $S \cup S^{-1}$ equal to e

$$xss^{-1}y \sim xs^{-1}sy \sim xy \sim xry$$

1.2 – Presentations

$$G = \langle S \mid R \rangle$$

S – set of symbols S^{-1} – each symbol tagged with an inverse

$L(S \cup S^{-1})$ – all strings in letters of $S \cup S^{-1}$

R – words $w \in L(S \cup S^{-1})$ in $S \cup S^{-1}$ equal to e

$$xss^{-1}y \sim xs^{-1}sy \sim xy \sim xry$$

$\langle S|R \rangle$ is the set of equivalence classes of $L(S \cup S^{-1})$ under \sim

1.2 – Presentations

$$G = \langle S \mid R \rangle$$

S – set of symbols S^{-1} – each symbol tagged with an inverse

$L(S \cup S^{-1})$ – all strings in letters of $S \cup S^{-1}$

R – words $w \in L(S \cup S^{-1})$ in $S \cup S^{-1}$ equal to e

$$xss^{-1}y \sim xs^{-1}sy \sim xy \sim xry$$

$\langle S | R \rangle$ is the set of equivalence classes of $L(S \cup S^{-1})$ under \sim

- | | |
|--|--|
| ① $\langle a \mid \rangle$ | ④ $\langle a, b \mid aba^{-1}b^{-1} \rangle$ |
| ② $\langle a \mid a^n \rangle$ | ⑤ $\langle a, b \mid a^n, b^2, (ab)^2 \rangle$ |
| ③ $\langle a, b \mid a^3, b^2, (ab)^2 \rangle$ | ⑥ $\langle a, b \mid a^2, b^2 \rangle$ |

1.2 – Cayley Graphs

Definition

The **Cayley graph** of a group $G = \langle S \mid R \rangle$, $\text{Cay}(G)$, has vertices g for every $g \in G$, and directed edges $g \rightarrow gs$.

1.2 – Cayley Graphs

Definition


The **Cayley graph** of a group $G = \langle S \mid R \rangle$, $\text{Cay}(G)$, has vertices g for every $g \in G$, and directed edges $g \rightarrow gs$.



1.2 – Cayley Graphs

Definition


The **Cayley graph** of a group $G = \langle S \mid R \rangle$, $\text{Cay}(G)$, has vertices g for every $g \in G$, and directed edges $g \rightarrow gs$.



1.2 – Cayley Graphs

Definition

The **Cayley graph** of a group $G = \langle S \mid R \rangle$, $\text{Cay}(G)$, has vertices g for every $g \in G$, and directed edges $g \rightarrow gs$.



1.2 – Cayley Graphs


Definition

The **Cayley graph** of a group $G = \langle S \mid R \rangle$, $\text{Cay}(G)$, has vertices g for every $g \in G$, and directed edges $g \rightarrow gs$.

1.2 – Cayley Graphs

Definition



The **Cayley graph** of a group $G = \langle S \mid R \rangle$, $\text{Cay}(G)$, has vertices g for every $g \in G$, and directed edges $g \rightarrow gs$.



1.2 – Cayley Graphs

Definition



The **Cayley graph** of a group $G = \langle S \mid R \rangle$, $\text{Cay}(G)$, has vertices g for every $g \in G$, and directed edges $g \rightarrow gs$.



1.2 – Cayley Graphs

Definition


The **Cayley graph** of a group $G = \langle S \mid R \rangle$, $\text{Cay}(G)$, has vertices g for every $g \in G$, and directed edges $g \rightarrow gs$.



1.2 – The Word Problem

Problem (The Word Problem)

Given a group G with finite presentation $\langle S \mid R \rangle$ (i.e. S and R are both finite), for any word $w \in L(S \cup S^{-1})$, decide whether $[w]$ is the identity element e in a finite amount of time.



1.2 – Bad News

Theorem (Novikov-Boone)

*The word problem for finitely presented groups is **undecidable** – in particular, there exists a finitely presented group G for which the word problem is undecidable.*

1.2 – Bad News

Theorem (Novikov-Boone)


*The word problem for finitely presented groups is **undecidable** – in particular, there exists a finitely presented group G for which the word problem is undecidable.*

Theorem (Higman's Embedding Theorem)

*Every finitely generated, **recursively presented** group H can be embedded as a subgroup of some finitely presented group G , i.e. H 's set of relations is **recursively enumerable**.*


1.2 – Free Groups

$$F_n = \langle x_1, \dots, x_n \mid \rangle$$



1.2 – Free Groups

$$F_n = \langle x_1, \dots, x_n \mid \rangle$$



Theorem (Nielsen-Schreier)

Every subgroup of a free group is free.

Idea 2 – Generalizations

- (★) *Doing Calculus Anywhere (Manifolds, Differential Forms)* – One of the staples of calculus is the Fundamental Theorem of Calculus, which has many further generalizations further afield in multivariable calculus. Let's see why these are actually all the same thing!
- *A Case Study in Volume (Measure Theory)* – So you think you know what volume is? Here we test your intuitive understanding of volume and use edge cases as a case study for generalizing what you might think of as “volume.”

2.1 – Fundamental Theorems of (Multivariable) Calculus

Theorem (Fundamental Theorem of Calculus)

$$\int_a^b f(x) dx = F(b) - F(a).$$

2.1 – Fundamental Theorems of (Multivariable) Calculus

Theorem (Fundamental Theorem of Calculus)

$$\int_a^b f(x) dx = F(b) - F(a).$$

Theorem (Fund. Thm. of Line Integrals)

$$\int_{\gamma} \nabla \varphi(\vec{r}) \cdot d\vec{r} = \varphi(\gamma(1)) - \varphi(\gamma(0)).$$

Theorem (Green's Theorem)

$$\oint_C (L dx + M dy) = \iint_D \left(\frac{\partial M}{\partial x} - \frac{\partial L}{\partial y} \right) dx dy$$

2.1 – Fundamental Theorems of (Multivariable) Calculus

Theorem (Kelvin-Stokes' Theorem)

$$\iint_S (\nabla \times \vec{F}) \cdot d\vec{S} = \oint_{\partial S} \vec{F} \cdot d\vec{r}$$

Theorem (Divergence Theorem)

$$\iint_V (\nabla \cdot \vec{F}) dV = \iint_{\partial V} \vec{F} \cdot d\vec{S}$$

2.1 – Fundamental Theorems of (Multivariable) Calculus

Theorem (Kelvin-Stokes' Theorem)

$$\iint_S (\nabla \times \vec{F}) \cdot d\vec{S} = \oint_{\partial S} \vec{F} \cdot d\vec{r}$$

Theorem (Divergence Theorem)

$$\iint_V (\nabla \cdot \vec{F}) dV = \iint_{\partial V} \vec{F} \cdot d\vec{S}$$

What do these theorems all have in common?

*Integrating a **function** over the **boundary of a space** is equal to integrating the **derivative** over the **whole space**.*

2.1 – Generalized Stokes' Theorem

Theorem (Generalized Stokes' Theorem)

Let M be a k -dimensional **oriented smooth manifold-with-boundary** in \mathbb{R}^n , and give the boundary ∂M the **boundary orientation**. Let φ be a smooth **$(k - 1)$ -form** defined on an open set containing M . Then


$$\int_{\partial M} \varphi = \int_M d\varphi.$$

2.1 – Example

Problem

Verify that Stokes' Theorem holds for the surface S given by

$z = 5 - x^2 - y^2$ above the plane $z = 1$ (oriented upwards), and vector field $\vec{F} = z^2\hat{i} - 3xy\hat{j} + x^3y^3\hat{k}$.



2.2 – Warmups

I hear y'all know how to find volumes in geometry – let's test that!
What's the volume of:

2.2 – Warmups

I hear y'all know how to find volumes in geometry – let's test that!
What's the volume of:

- A three-dimensional ball of radius 1? ($\{\vec{x} \in \mathbb{R}^3 : |\vec{x}| \leq 1\}$)

2.2 – Warmups

I hear y'all know how to find volumes in geometry – let's test that!
What's the volume of:

- A three-dimensional ball of radius 1? ($\{\vec{x} \in \mathbb{R}^3 : |\vec{x}| \leq 1\}$)
- A two-dimensional disk of radius 1? ($\{\vec{x} \in \mathbb{R}^2 : |\vec{x}| \leq 1\}$)

2.2 – Warmups

I hear y'all know how to find volumes in geometry – let's test that!
What's the volume of:

- A three-dimensional ball of radius 1? ($\{\vec{x} \in \mathbb{R}^3 : |\vec{x}| \leq 1\}$)
- A two-dimensional disk of radius 1? ($\{\vec{x} \in \mathbb{R}^2 : |\vec{x}| \leq 1\}$)
- The interval $[0, 1]$ in \mathbb{R} ?

2.2 – Warmups

I hear y'all know how to find volumes in geometry – let's test that!
What's the volume of:

- A three-dimensional ball of radius 1? ($\{\vec{x} \in \mathbb{R}^3 : |\vec{x}| \leq 1\}$)
- A two-dimensional disk of radius 1? ($\{\vec{x} \in \mathbb{R}^2 : |\vec{x}| \leq 1\}$)
- The interval $[0, 1]$ in \mathbb{R} ?
- The singleton point $\{0\}$?

2.2 – Linear Exercises (Part I)

For the following, tell me whether the volume of the following sets are 0, 1, or not defined, in \mathbb{R} :

2.2 – Linear Exercises (Part I)

For the following, tell me whether the volume of the following sets are 0, 1, or not defined, in \mathbb{R} :

- ① The two points $\{0, 1\}$.

2.2 – Linear Exercises (Part I)

For the following, tell me whether the volume of the following sets are 0, 1, or not defined, in \mathbb{R} :

- ① The two points $\{0, 1\}$.
- ② For $n \in \mathbb{Z}^+$, $\left\{ \frac{k}{n} : 0 \leq k \leq n, k \in \mathbb{Z} \right\}$ (fractions with denominator n).

2.2 – Linear Exercises (Part I)

For the following, tell me whether the volume of the following sets are 0, 1, or not defined, in \mathbb{R} :

- ① The two points $\{0, 1\}$.
- ② For $n \in \mathbb{Z}^+$, $\left\{ \frac{k}{n} : 0 \leq k \leq n, k \in \mathbb{Z} \right\}$ (fractions with denominator n).
- ③ The converging sequence $\left\{ \frac{1}{2^k} : k \in \mathbb{Z}_{\geq 0} \right\}$.

2.2 – Linear Exercises (Part I)

For the following, tell me whether the volume of the following sets are 0, 1, or not defined, in \mathbb{R} :

- ① The two points $\{0, 1\}$.
- ② For $n \in \mathbb{Z}^+$, $\left\{ \frac{k}{n} : 0 \leq k \leq n, k \in \mathbb{Z} \right\}$ (fractions with denominator n).
- ③ The converging sequence $\left\{ \frac{1}{2^k} : k \in \mathbb{Z}_{\geq 0} \right\}$.
- ④ All dyadic fractions between 0 and 1, $\left\{ \frac{m}{2^n} : m, n \in \mathbb{Z} \right\} \cap [0, 1]$ (fractions whose denominator is a power of 2).

2.2 – Linear Exercises (Part I)

For the following, tell me whether the volume of the following sets are 0, 1, or not defined, in \mathbb{R} :

- ① The two points $\{0, 1\}$.
- ② For $n \in \mathbb{Z}^+$, $\left\{ \frac{k}{n} : 0 \leq k \leq n, k \in \mathbb{Z} \right\}$ (fractions with denominator n).
- ③ The converging sequence $\left\{ \frac{1}{2^k} : k \in \mathbb{Z}_{\geq 0} \right\}$.
- ④ All dyadic fractions between 0 and 1, $\left\{ \frac{m}{2^n} : m, n \in \mathbb{Z} \right\} \cap [0, 1]$ (fractions whose denominator is a power of 2).
- ⑤ The rational numbers between 0 and 1, $\mathbb{Q} \cap [0, 1]$.

2.2 – Linear Exercises (Part II)

For the following, tell me whether the volume of the following sets are 0, 1, or not defined:

2.2 – Linear Exercises (Part II)

For the following, tell me whether the volume of the following sets are 0, 1, or not defined:

- ⑥ Let $C_0 = [0, 1]$, and for $n \geq 1$, assuming C_{n-1} is a disjoint union of closed intervals, remove the middle third of each interval to get C_n . Let $C = \bigcap_{k=0}^{\infty} C_k$. (This set is *uncountable*, unlike any of the examples up until this point.)

2.2 – Linear Exercises (Part II)

For the following, tell me whether the volume of the following sets are 0, 1, or not defined:

- ⑥ Let $C_0 = [0, 1]$, and for $n \geq 1$, assuming C_{n-1} is a disjoint union of closed intervals, remove the middle third of each interval to get C_n . Let $C = \bigcap_{k=0}^{\infty} C_k$. (This set is *uncountable*, unlike any of the examples up until this point.)
- ⑦ Let $F_0 = [0, 2]$. To generate F_1 , remove the middle $\frac{1}{4}$ of each closed interval in F_0 . In general, remove the middle $\frac{1}{4^i}$ of each interval from the intervals in F_{i-1} to get F_i . Let $F = \bigcap_{k=0}^{\infty} F_k$.

2.2 – Linear Exercises (Part II)

For the following, tell me whether the volume of the following sets are 0, 1, or not defined:

- ⑥ Let $C_0 = [0, 1]$, and for $n \geq 1$, assuming C_{n-1} is a disjoint union of closed intervals, remove the middle third of each interval to get C_n . Let $C = \bigcap_{k=0}^{\infty} C_k$. (This set is *uncountable*, unlike any of the examples up until this point.)
- ⑦ Let $F_0 = [0, 2]$. To generate F_1 , remove the middle $\frac{1}{4}$ of each closed interval in F_0 . In general, remove the middle $\frac{1}{4^i}$ of each interval from the intervals in F_{i-1} to get F_i . Let $F = \bigcap_{k=0}^{\infty} F_k$.
- ⑧ Consider $[0, 1]$ under the equivalence relation \sim such that $x \sim y$ if $|x - y| \in \mathbb{Q}$. Partition $[0, 1]$ into equivalence classes, and let V consist of one representative from each equivalence class.

2.2 – Dimension Warmups

Now, let's warm up for a different question:

2.2 – Dimension Warmups

Now, let's warm up for a different question:

- What's the 1-D volume of the sphere of radius $\frac{1}{2\sqrt{\pi}}$ in \mathbb{R}^3 ?

2.2 – Dimension Warmups

Now, let's warm up for a different question:

- What's the 1-D volume of the sphere of radius $\frac{1}{2\sqrt{\pi}}$ in \mathbb{R}^3 ?
- What's the 2-D volume of the sphere of radius $\frac{1}{2\sqrt{\pi}}$ in \mathbb{R}^3 ?

2.2 – Dimension Warmups

Now, let's warm up for a different question:

- What's the 1-D volume of the sphere of radius $\frac{1}{2\sqrt{\pi}}$ in \mathbb{R}^3 ?
- What's the 2-D volume of the sphere of radius $\frac{1}{2\sqrt{\pi}}$ in \mathbb{R}^3 ?
- What's the 3-D volume of the sphere of radius $\frac{1}{2\sqrt{\pi}}$ in \mathbb{R}^3 ?

2.2 – Dimension Warmups

Now, let's warm up for a different question:



- What's the 1-D volume of the sphere of radius $\frac{1}{2\sqrt{\pi}}$ in \mathbb{R}^3 ?
- What's the 2-D volume of the sphere of radius $\frac{1}{2\sqrt{\pi}}$ in \mathbb{R}^3 ?
- What's the 3-D volume of the sphere of radius $\frac{1}{2\sqrt{\pi}}$ in \mathbb{R}^3 ?

Note that for only one of these is the answer not 0 or ∞ – it's when the dimension is 2! We can actually define the dimension in this way...

2.2 – Dimensionality Exercises

For these spaces, compute the definition in this way, assuming there is some $d \in \mathbb{R}^+$ for which the volume is positive and finite (it's not going to be integral!):

- ① The Cantor set C .
- ② The Sierpinski triangle.
- ③ The Koch snowflake.
- ④ The Menger sponge.



Idea 3 – Breaking and Repairing

- *I Forgot How to Factor (Algebraic Number Theory)* – We will try solving some Diophantine equations in some creative ways, and realize that I don't really know how to prime-factorize things anymore. Can we fix that?
- *The Answer Is Not True (Analysis/Foundations)* – Because everyone loves the true/false section of a math exam, we're going to have a true/false party!
Wait, I forgot to take the spoiler out of the section header... fine, to make it more interesting, be prepared to explain your answers!

3.1 – A Diophantine Equation

Problem

Find all solutions in integers to $y^2 = x^3 - 1$.

3.1 – A Diophantine Equation

Problem

Find all solutions in integers to $y^2 = x^3 - 1$.

$$\Rightarrow x^3 = y^2 + 1$$

3.1 – A Diophantine Equation

Problem

Find all solutions in integers to $y^2 = x^3 - 1$.

$$\Rightarrow x^3 = y^2 + 1 = (y + i)(y - i)$$

y is not odd – if it were, then $y^2 + 1 \equiv 2 \pmod{4}$ and $x^3 \equiv 0 \pmod{4}$.

3.1 – A Diophantine Equation

Problem

Find all solutions in integers to $y^2 = x^3 - 1$.

$$\Rightarrow x^3 = y^2 + 1 = (y + i)(y - i)$$

y is not odd – if it were, then $y^2 + 1 \equiv 2 \pmod{4}$ and $x^3 \equiv 0 \pmod{4}$.

Claim. $y + i$ and $y - i$ have no non-unit common divisors.

3.1 – A Diophantine Equation

Problem

Find all solutions in integers to $y^2 = x^3 - 1$.

$$\Rightarrow x^3 = y^2 + 1 = (y + i)(y - i)$$

y is not odd – if it were, then $y^2 + 1 \equiv 2 \pmod{4}$ and $x^3 \equiv 0 \pmod{4}$.

Claim. $y + i$ and $y - i$ have no non-unit common divisors.

Proof. $\pi \mid y + i, y - i \Rightarrow \pi \mid 2i = (1 + i)^2 \Rightarrow \pi = (1 + i)$.

3.1 – A Diophantine Equation

Problem

Find all solutions in integers to $y^2 = x^3 - 1$.

$$\Rightarrow x^3 = y^2 + 1 = (y + i)(y - i)$$

y is not odd – if it were, then $y^2 + 1 \equiv 2 \pmod{4}$ and $x^3 \equiv 0 \pmod{4}$.

Claim. $y + i$ and $y - i$ have no non-unit common divisors.

Proof. $\pi | y + i, y - i \Rightarrow \pi | 2i = (1 + i)^2 \Rightarrow \pi = (1 + i)$. Then

$$y + i = (1 + i)(a + bi), y - i = (1 - i)(a - bi) \Rightarrow y^2 + 1 = 2(a^2 + b^2).$$

Contradiction, $y^2 + 1$ is odd.

3.1 – A Diophantine Equation

Problem

Find all solutions in integers to $y^2 = x^3 - 1$.

$$\Rightarrow x^3 = y^2 + 1 = (y + i)(y - i)$$

y is not odd – if it were, then $y^2 + 1 \equiv 2 \pmod{4}$ and $x^3 \equiv 0 \pmod{4}$.

Claim. $y + i$ and $y - i$ have no non-unit common divisors.

Proof. $\pi | y + i, y - i \Rightarrow \pi | 2i = (1 + i)^2 \Rightarrow \pi = (1 + i)$. Then

$$y + i = (1 + i)(a + bi), y - i = (1 - i)(a - bi) \Rightarrow y^2 + 1 = 2(a^2 + b^2).$$

Contradiction, $y^2 + 1$ is odd.

$$(y + i) = (a + bi)^3 = (a^3 - 3ab^2) + (3a^2b - b^3), \text{ so } b(3a^2 - b^2) = 1, \text{ and } b = \pm 1.$$

3.1 – A Diophantine Equation

Problem

Find all solutions in integers to $y^2 = x^3 - 1$.

$$\Rightarrow x^3 = y^2 + 1 = (y + i)(y - i)$$

y is not odd – if it were, then $y^2 + 1 \equiv 2 \pmod{4}$ and $x^3 \equiv 0 \pmod{4}$.

Claim. $y + i$ and $y - i$ have no non-unit common divisors.

Proof. $\pi | y + i, y - i \Rightarrow \pi | 2i = (1 + i)^2 \Rightarrow \pi = (1 + i)$. Then

$$y + i = (1 + i)(a + bi), y - i = (1 - i)(a - bi) \Rightarrow y^2 + 1 = 2(a^2 + b^2).$$

Contradiction, $y^2 + 1$ is odd.

$(y + i) = (a + bi)^3 = (a^3 - 3ab^2) + (3a^2b - b^3)$, so $b(3a^2 - b^2) = 1$, and $b = \pm 1$. If $b = 1$, no sol'ns, if $b = -1$, then $a = 0$, so $y = 0, x = 1$.

3.1 – A Diophantine Equation

Problem

Find all solutions in integers to $y^2 = x^3 - 1$.

$$\Rightarrow x^3 = y^2 + 1 = (y + i)(y - i)$$

y is not odd – if it were, then $y^2 + 1 \equiv 2 \pmod{4}$ and $x^3 \equiv 0 \pmod{4}$.

Claim. $y + i$ and $y - i$ have no non-unit common divisors.

Proof. $\pi | y + i, y - i \Rightarrow \pi | 2i = (1 + i)^2 \Rightarrow \pi = (1 + i)$. Then

$$y + i = (1 + i)(a + bi), y - i = (1 - i)(a - bi) \Rightarrow y^2 + 1 = 2(a^2 + b^2).$$

Contradiction, $y^2 + 1$ is odd.

$(y + i) = (a + bi)^3 = (a^3 - 3ab^2) + (3a^2b - b^3)$, so $b(3a^2 - b^2) = 1$, and $b = \pm 1$. If $b = 1$, no sol'ns, if $b = -1$, then $a = 0$, so $y = 0, x = 1$.

$\Rightarrow (x, y) = (1, 0)$ is the only solution.

3.1 – Another Diophantine Equation

Problem

Find all solutions in integers to $y^2 = x^3 - 61$.

3.1 – Another Diophantine Equation

Problem

Find all solutions in integers to $y^2 = x^3 - 61$.

$$\Rightarrow x^3 = y^2 + 61$$

3.1 – Another Diophantine Equation

Problem

Find all solutions in integers to $y^2 = x^3 - 61$.

$$\Rightarrow x^3 = y^2 + 61 = (y + \sqrt{-61})(y - \sqrt{-61})$$

y is not odd – if it were, then $y^2 + 61 \equiv 2 \pmod{4}$ and $x^3 \equiv 0 \pmod{4}$.

3.1 – Another Diophantine Equation

Problem

Find all solutions in integers to $y^2 = x^3 - 61$.

$$\Rightarrow x^3 = y^2 + 61 = (y + \sqrt{-61})(y - \sqrt{-61})$$

y is not odd – if it were, then $y^2 + 61 \equiv 2 \pmod{4}$ and $x^3 \equiv 0 \pmod{4}$.

Claim. $y + \sqrt{-61}$ and $y - \sqrt{-61}$ have no non-unit common divisors.

3.1 – Another Diophantine Equation

Problem

Find all solutions in integers to $y^2 = x^3 - 61$.

$$\Rightarrow x^3 = y^2 + 61 = (y + \sqrt{-61})(y - \sqrt{-61})$$

y is not odd – if it were, then $y^2 + 61 \equiv 2 \pmod{4}$ and $x^3 \equiv 0 \pmod{4}$.

Claim. $y + \sqrt{-61}$ and $y - \sqrt{-61}$ have no non-unit common divisors.

Proof. $\pi \mid y + \sqrt{-61}, y - \sqrt{-61} \Rightarrow \pi \mid 2\sqrt{-61}$.

3.1 – Another Diophantine Equation

Problem

Find all solutions in integers to $y^2 = x^3 - 61$.

$$\Rightarrow x^3 = y^2 + 61 = (y + \sqrt{-61})(y - \sqrt{-61})$$

y is not odd – if it were, then $y^2 + 61 \equiv 2 \pmod{4}$ and $x^3 \equiv 0 \pmod{4}$.

Claim. $y + \sqrt{-61}$ and $y - \sqrt{-61}$ have no non-unit common divisors.

Proof. $\pi \mid y + \sqrt{-61}, y - \sqrt{-61} \Rightarrow \pi \mid 2\sqrt{-61}$. $|\pi|^2 \mid y^2 + 61$ odd so $|\pi|^2$ odd, and therefore $|\pi|^2 = 61$.

3.1 – Another Diophantine Equation

Problem

Find all solutions in integers to $y^2 = x^3 - 61$.

$$\Rightarrow x^3 = y^2 + 61 = (y + \sqrt{-61})(y - \sqrt{-61})$$

y is not odd – if it were, then $y^2 + 61 \equiv 2 \pmod{4}$ and $x^3 \equiv 0 \pmod{4}$.

Claim. $y + \sqrt{-61}$ and $y - \sqrt{-61}$ have no non-unit common divisors.

Proof. $\pi \mid y + \sqrt{-61}, y - \sqrt{-61} \Rightarrow \pi \mid 2\sqrt{-61}$. $|\pi|^2 \mid y^2 + 61$ odd so $|\pi|^2$ odd, and therefore $|\pi|^2 = 61$. Then $61 \mid y^2 + 61$, $61 \mid y$, and so $61 \mid x$ and so $61^2 \mid x^3 - y^2 = 61$, contradiction.

3.1 – Another Diophantine Equation

Problem

Find all solutions in integers to $y^2 = x^3 - 61$.

$$\Rightarrow x^3 = y^2 + 61 = (y + \sqrt{-61})(y - \sqrt{-61})$$

y is not odd – if it were, then $y^2 + 61 \equiv 2 \pmod{4}$ and $x^3 \equiv 0 \pmod{4}$.

Claim. $y + \sqrt{-61}$ and $y - \sqrt{-61}$ have no non-unit common divisors.

Proof. $\pi | y + \sqrt{-61}, y - \sqrt{-61} \Rightarrow \pi | 2\sqrt{-61}$. $|\pi|^2 | y^2 + 61$ odd so $|\pi|^2$ odd, and therefore $|\pi|^2 = 61$. Then $61 | y^2 + 61$, $61 | y$, and so $61 | x$ and so $61^2 | x^3 - y^2 = 61$, contradiction.

$(y + \sqrt{-61}) = (a + b\sqrt{-61})^3 = (a^3 - 183ab^2) + (3a^2b - 61b^3)$, so $b(3a^2 - 61b^2) = 1$, and $b = \pm 1$.

3.1 – Another Diophantine Equation

Problem

Find all solutions in integers to $y^2 = x^3 - 61$.

$$\Rightarrow x^3 = y^2 + 61 = (y + \sqrt{-61})(y - \sqrt{-61})$$

y is not odd – if it were, then $y^2 + 61 \equiv 2 \pmod{4}$ and $x^3 \equiv 0 \pmod{4}$.

Claim. $y + \sqrt{-61}$ and $y - \sqrt{-61}$ have no non-unit common divisors.

Proof. $\pi | y + \sqrt{-61}, y - \sqrt{-61} \Rightarrow \pi | 2\sqrt{-61}$. $|\pi|^2 | y^2 + 61$ odd so $|\pi|^2$ odd, and therefore $|\pi|^2 = 61$. Then $61 | y^2 + 61$, $61 | y$, and so $61 | x$ and so $61^2 | x^3 - y^2 = 61$, contradiction.

$(y + \sqrt{-61}) = (a + b\sqrt{-61})^3 = (a^3 - 183ab^2) + (3a^2b - 61b^3)$, so $b(3a^2 - 61b^2) = 1$, and $b = \pm 1$. In either case for b , no solutions.

3.1 – Another Diophantine Equation

Problem

Find all solutions in integers to $y^2 = x^3 - 61$.

$$\Rightarrow x^3 = y^2 + 61 = (y + \sqrt{-61})(y - \sqrt{-61})$$

y is not odd – if it were, then $y^2 + 61 \equiv 2 \pmod{4}$ and $x^3 \equiv 0 \pmod{4}$.

Claim. $y + \sqrt{-61}$ and $y - \sqrt{-61}$ have no non-unit common divisors.

Proof. $\pi | y + \sqrt{-61}, y - \sqrt{-61} \Rightarrow \pi | 2\sqrt{-61}$. $|\pi|^2 | y^2 + 61$ odd so $|\pi|^2$ odd, and therefore $|\pi|^2 = 61$. Then $61 | y^2 + 61$, $61 | y$, and so $61 | x$ and so $61^2 | x^3 - y^2 = 61$, contradiction.

$(y + \sqrt{-61}) = (a + b\sqrt{-61})^3 = (a^3 - 183ab^2) + (3a^2b - 61b^3)$, so $b(3a^2 - 61b^2) = 1$, and $b = \pm 1$. In either case for b , no solutions.

There are no solutions in integers to this equation. Except...

3.1 – Formalizing Primes and Factoring

Definition

An element π is **prime** if for any a and b such that $\pi \mid ab$, then either $\pi \mid a$ or $\pi \mid b$.

3.1 – Formalizing Primes and Factoring

Definition

An element π is **prime** if for any a and b such that $\pi \mid ab$, then either $\pi \mid a$ or $\pi \mid b$.

Definition

An element π is **irreducible** if it is not invertible in the ring of integers (a **unit**) and it is not the product of two non-invertible elements.

3.1 – Formalizing Primes and Factoring

Definition

An element π is **prime** if for any a and b such that $\pi \mid ab$, then either $\pi \mid a$ or $\pi \mid b$.

Definition

An element π is **irreducible** if it is not invertible in the ring of integers (a **unit**) and it is not the product of two non-invertible elements.

Definition

A **unique factorization domain** (or UFD) is a ring where every element can be uniquely written as a product of irreducible elements and a unit, up to ordering and multiplication by units.

3.1 – Ideals

Definition

An **ideal** I is an additively closed subset of a ring R such that for any $r \in R$, $rI = I$.

3.1 – Ideals

Definition

An **ideal** I is an additively closed subset of a ring R such that for any $r \in R$, $rI = I$.

Definition

An ideal $P \subset R$ is **prime** if for any $a, b \in R$ such that $ab \in P$, then either $a \in P$ or $b \in P$.

3.1 – Ideals

Definition

An **ideal** I is an additively closed subset of a ring R such that for any $r \in R$, $rI = I$.

Definition

An ideal $P \subset R$ is **prime** if for any $a, b \in R$ such that $ab \in P$, then either $a \in P$ or $b \in P$.

Theorem

Every nonzero ideal I factors uniquely as a product of prime ideals in a Dedekind domain, where the product of ideals I and J , written IJ , is defined as the closure of the set $\{ab : a \in I, b \in J\}$ under addition.

3.1 – Class Group

Definition

For a ring R , consider its field of fractions K . A **fractional ideal** $J \subseteq K$ is closed under addition and multiplication by R , and there exists some nonzero $r \in R$ such that $rJ \subseteq R$. These ideals form a group in a Dedekind domain.

3.1 – Class Group

Definition

For a ring R , consider its field of fractions K . A **fractional ideal** $J \subseteq K$ is closed under addition and multiplication by R , and there exists some nonzero $r \in R$ such that $rJ \subseteq R$. These ideals form a group in a Dedekind domain.

Definition

An ideal I is **principal** if it consists of all the multiples of some element π in the ring.

3.1 – Class Group

Definition

For a ring R , consider its field of fractions K . A **fractional ideal** $J \subseteq K$ is closed under addition and multiplication by R , and there exists some nonzero $r \in R$ such that $rJ \subseteq R$. These ideals form a group in a Dedekind domain.

Definition

An ideal I is **principal** if it consists of all the multiples of some element π in the ring.

Definition

For a ring of integers R with corresponding field of fractions K , the **class group** is the quotient group J_K/P_K , where J_K is the collection of fractional ideals, and P_K is the subgroup of principal ideals.

3.2 – True/False Party (Smoothness, Part I)

- ① Every function that is continuous everywhere is also differentiable everywhere.
- ② The derivative of a function is always continuous.
- ③ If a function's derivative is finite everywhere, the derivative must be bounded everywhere.
- ④ Every function that is continuous everywhere must be differentiable at some point.
- ⑤ If a continuous function has derivative zero almost everywhere, it is constant.
- ⑥ A function continuous on the irrationals must be continuous at some rational.

3.2 – True/False Party (Integration, Part II)

- ① Every set that has a positive volume must contain some open interval.
- ② A function cannot be integrable if its set of points of discontinuity is dense in \mathbb{R} .
- ③ Every bounded subset of \mathbb{R} can be given a measure.
- ④ Regardless of your definition of the integral, the integral $\int_{-\infty}^{\infty} \frac{\sin x}{x}$ exists and is equal to π .
- ⑤ A function that integrates to 0 on any open interval is identically zero.

3.2 – True/False Party (Multivar/Series, Part III)

- ① A function $\mathbb{R}^2 \rightarrow \mathbb{R}$ that has first partial derivatives everywhere must be differentiable.
- ② For a differentiable function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$, its mixed second-order partials are equal.
- ③ For a function f that is positive and continuous on $x \geq 1$, $\int_1^\infty f(x) dx$ converges iff $\sum_{n=1}^\infty f(n)$ converges.
- ④ A function whose Taylor series at a point converges everywhere must converge to the function.

3.2 – True/False Party (General, Part IV)

- ① Every Cartesian product of non-empty sets is non-empty.
- ② Every vector space has a basis.
- ③ Every surjective function has a right inverse.
- ④ Every linear function $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous.
- ⑤ Induction can only be done over finite sets, such as \mathbb{N} .

3.2 – True/False Party (General, Part IV)

- ① Every Cartesian product of non-empty sets is non-empty.
- ② Every vector space has a basis.
- ③ Every surjective function has a right inverse.
- ④ Every linear function $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous.
- ⑤ Induction can only be done over finite sets, such as \mathbb{N} .

Theorem (Axiom of Choice)

Let $\mathcal{A} = \{A_i\}_{i \in I}$ be a set of non-empty sets. There is a function $f : \mathcal{A} \rightarrow \bigcup_{i \in I} A_i$ such that $f(A_i)$ is an element of A_i for all $i \in I$.