

HW1.2

SCIE 001 Math

October 31, 2025

1. Show that if $x > 0$ and $x \neq 1$ then,

$$\ln(x) < x - 1$$

Justify your claims.

Start of proof:

Lemma 1: $\ln(x) < (x - 1)$ for all $x > 1$

$\ln(x)$ is continuous on $[1, +\infty)$ and differentiable on $(1, +\infty)$

By the Mean Value Theorem there exists some $c > 1$ for all $x > 1$ such that:

$$\begin{aligned}\frac{d}{dc}(\ln(c)) &= \frac{\ln(x) - \ln(1)}{x - 1} \\ \frac{1}{c} &= \frac{\ln(x) - \ln(1)}{x - 1}\end{aligned}$$

Since we are only looking at $c > 1$:

$$\begin{aligned}0 &< \frac{1}{c} < 1 \\ \frac{\ln(x) - \ln(1)}{x - 1} &= \frac{1}{c} < 1 \\ \frac{\ln(x) - \ln(1)}{x - 1} &< 1\end{aligned}$$

Since $x > 1$, $x - 1 > 0$:

$$\ln(x) - \ln(1) < x - 1$$

$\ln(1) = 0$ therefore:

For all $x > 1$: $\ln(x) < x - 1$

Lemma 2: $\ln(x) < (x - 1)$ for all $0 < x < 1$

$\ln(x)$ is continuous on $[0, 1]$ and differentiable on $(0, 1)$

By the Mean Value Theorem there exists some $0 < c < 1$ for all $0 < x < 1$ such that:

$$\begin{aligned}\frac{d}{dc}(\ln(c)) &= \frac{\ln(1) - \ln(x)}{1 - x} \\ \frac{1}{c} &= \frac{\ln(x) - \ln(1)}{x - 1}\end{aligned}$$

Since we are only looking at $0 < c < 1$:

$$\begin{aligned}\frac{1}{c} &> 1 \\ \frac{\ln(x) - \ln(1)}{x - 1} &= \frac{1}{c} > 1 \\ \frac{\ln(x) - \ln(1)}{x - 1} &> 1\end{aligned}$$

Since $0 < x < 1$, $x - 1 < 0$, so we must flip the comparison when we multiply.

$$\ln(x) - \ln(1) < x - 1$$

$\ln(1) = 0$ therefore:

For all $0 < x < 1$: $\ln(x) < x - 1$
--

Combining the comparisons from **Lemma 1** and **Lemma 2** we get:

For all $x > 0, x \neq 1$: $\ln(x) < x - 1$
--

Q.E.D

2. Consider the “Gompertz” model of population dynamics,

$$\frac{dP}{dt} = cP \ln\left(\frac{K}{P}\right), \quad P(0) = P_0$$

where c and K are positive constants. Verify that the functions $P(t) = Ke^{(-be^{(-ct)})}$, where b is a constant, are solutions of the Gompertz equation. Then find an expression for b in terms of P_0 .

For a function P to be a solution of the Gompertz equation, it must satisfy the ordinary differential equation:

$$\frac{dP}{dt} = cP \ln\left(\frac{K}{P}\right)$$

To prove that the functions $P(t) = Ke^{(-be^{(-ct)})}$ are solutions, we can rearrange to show equality:

$$\frac{dP}{dt} = \frac{dP}{dt}$$

Left Side

Right Side

$ \begin{aligned} &= \frac{d}{dt} \left(Ke^{(-be^{(-ct)})} \right) \\ &= Ke^{(-be^{(-ct)})} (-be^{(-ct)}) (-c) \\ &= Kcbe^{(-ct)} e^{(-be^{(-ct)})} \end{aligned} $	$ \begin{aligned} &= cP \ln\left(\frac{K}{P}\right) \\ &= cKe^{(-be^{(-ct)})} \ln\left(e^{-(-be^{(-ct)})}\right) \\ &= cKe^{(-be^{(-ct)})} be^{(-ct)} \\ &= Kcbe^{(-ct)} e^{(-be^{(-ct)})} \end{aligned} $
--	---

$$Kcbe^{(-ct)} e^{(-be^{(-ct)})} = Kcbe^{(-ct)} e^{(-be^{(-ct)})}$$

Left Side = Right Side

Therefore $P(t) = Ke^{(-be^{(-ct)})}$ are a family of solutions.

To find b in terms of P_0 , we just need to solve $P(0)$ for b :

$$\begin{aligned}
 P_0 &= P(0) \\
 &= Ke^{(-be^{(-c(0))})} \\
 P_0 &= Ke^{(-b)}
 \end{aligned}$$

$$\boxed{\therefore b = -\ln\left(\frac{P_0}{K}\right)}$$

3. **Let**

$$f(x) = \begin{cases} x^n \cos\left(\frac{1}{x}\right) & \text{if } x > 0 \\ x^n & \text{if } x \leq 0 \end{cases}$$

for $n > 0$.

(a) **For what values of n is f' differentiable at $x = 0$?**

For f' to be differentiable at $x = 0$:

$$\lim_{h \rightarrow 0} \frac{f'(0+h) - f'(0)}{h}$$

must exist.

Consequently, to find the values of n for which f' is differentiable, we must first find the derivative of f .

The derivative of f can be seen as a piecewise function for values of a below 0, at 0, or above 0. Below and above zero we can just take the derivative of the respective piecewise cases of f , but at zero we need to use the definition of the derivative.

$$\begin{array}{cc} a > 0 & a < 0 \\ f'(a) = (na^{n-1}) \cos\left(\frac{1}{a}\right) + (x^n) \left(-\sin\left(\frac{1}{a}\right)\right) (-1a^{-2}) & \left| \right. f'(a) = na^{n-1} \end{array}$$

$$a = 0$$

$$\begin{aligned} f'(a) = f'(0) &= \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(h) - 0^n}{h} \end{aligned}$$

This splits to left and right hand limits:

$$\begin{aligned} &\lim_{h \rightarrow 0^-} \frac{f(h) - 0^n}{h} \\ &= \lim_{h \rightarrow 0^-} \frac{h^n - 0^n}{h} \end{aligned}$$

For all $n \neq 0$:

$$\begin{aligned} &= \lim_{h \rightarrow 0^-} \frac{h^n}{h} \\ &= \lim_{h \rightarrow 0^-} (h^{n-1}) \end{aligned}$$

$n < 1$	$n = 1$	$n > 1$
$n - 1 < 0$	$n - 1 = 0$	$n - 1 > 0$
$\lim_{h \rightarrow 0^-} (h^{n-1}) = D.N.E$	$\lim_{h \rightarrow 0^-} (h^{n-1}) = 1$	$\lim_{h \rightarrow 0^-} (h^{n-1}) = 0$
		For our purposes,
		h^{n-1} is undefined
		for $h < 0$ when $n \notin \mathbb{Z}$
		$\therefore$ This only holds for
		$n \in \mathbb{Z}$

$$\lim_{h \rightarrow 0^+} \frac{f(h) - 0^n}{h}$$

$$= \lim_{h \rightarrow 0^+} \frac{h^n \cos\left(\frac{1}{h}\right) - 0^n}{h}$$

For all $n \neq 0$:

$$= \lim_{h \rightarrow 0^+} h^{n-1} \cos\left(\frac{1}{h}\right)$$

$n < 1$	$n = 1$
$n - 1 < 0$	
$\therefore \lim_{h \rightarrow 0^+} h^{n-1} \cos\left(\frac{1}{h}\right)$	$\lim_{h \rightarrow 0^+} h^{n-1} \cos\left(\frac{1}{h}\right)$
$= D.N.E$	$= \lim_{h \rightarrow 0^+} h^0 \cos\left(\frac{1}{h}\right)$
	$= \lim_{h \rightarrow 0^+} \cos\left(\frac{1}{h}\right)$
	$= D.N.E$

$n > 1$

$$-1 \leq \cos\left(\frac{1}{h}\right) \leq 1$$

For all $h > 0 : h^{n-1} > 0$

$$-h^{n-1} \leq h^{n-1} \cos\left(\frac{1}{h}\right) \leq h^{n-1}$$

$$\lim_{h \rightarrow 0^+} -h^{n-1} \leq \lim_{h \rightarrow 0^+} h^{n-1} \cos\left(\frac{1}{h}\right) \leq \lim_{h \rightarrow 0^+} h^{n-1}$$

$$0 \leq \lim_{h \rightarrow 0^+} h^{n-1} \cos\left(\frac{1}{h}\right) \leq 0$$

$\therefore$ by The Squeeze Theorem

$$\lim_{h \rightarrow 0^+} h^{n-1} \cos\left(\frac{1}{h}\right) = 0$$

The left and right hand limits exist and are equal only when $n > 1$.

Therefore:

$$\text{When } n > 1, n \in \mathbb{Z} : \quad f'(0) = 0$$

Now we have the tools to solve the original limit.

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{f'(0+h) - f'(0)}{h} & \quad \text{must exist} \\ = \lim_{h \rightarrow 0} \frac{f'(h) - 0}{h} & \quad \text{for } n > 1, n \in \mathbb{Z} \end{aligned}$$

This splits to left and right hand limits:

$$\begin{aligned} & \lim_{h \rightarrow 0^-} \frac{f'(h)}{h} \\ &= \lim_{h \rightarrow 0^-} \frac{nh^{n-1}}{h} \\ &= \lim_{h \rightarrow 0^-} nh^{n-2} \end{aligned}$$

$n < 2$ $n - 2 < 0$ $\lim_{h \rightarrow 0^-} nh^{n-2} = D.N.E$	$n = 2$ $\lim_{h \rightarrow 0^-} nh^{n-2}$ $= \lim_{h \rightarrow 0^-} 2h^0$ $= 2$	$n > 2$ $n - 2 > 0$ $\lim_{h \rightarrow 0^-} nh^{n-2}$ $= 0$ For our purposes, h^{n-2} is undefined for $h < 0$ when $n \notin \mathbb{Z}$ $\therefore$ Again this only holds for $n \in \mathbb{Z}$
---	--	---

$$\begin{aligned} & \lim_{h \rightarrow 0^+} \frac{f'(h)}{h} \\ &= \lim_{h \rightarrow 0^+} \frac{(nh^{n-1}) \cos\left(\frac{1}{h}\right) + (h^n)(-\sin\left(\frac{1}{h}\right))(-1h^{-2})}{h} \\ &= \lim_{h \rightarrow 0^+} (nh^{n-2}) \cos\left(\frac{1}{h}\right) + (h^{n-1}) \left(-\sin\left(\frac{1}{h}\right)\right)(-1h^{-2}) \\ &= \lim_{h \rightarrow 0^+} nh^{n-2} \cos\left(\frac{1}{h}\right) + h^{n-3} \sin\left(\frac{1}{h}\right) \end{aligned}$$

For $n < 3$:

$$n - 3 < 0$$

$$\lim_{h \rightarrow 0^+} h^{n-3} \sin\left(\frac{1}{h}\right) = D.N.E$$

$$\therefore \lim_{h \rightarrow 0^+} nh^{n-2} \cos\left(\frac{1}{h}\right) + h^{n-3} \sin\left(\frac{1}{h}\right) = D.N.E$$

For $n = 3$:

$$\begin{aligned}
& \lim_{h \rightarrow 0^+} nh^{n-2} \cos\left(\frac{1}{h}\right) + h^{n-3} \sin\left(\frac{1}{h}\right) \\
&= \lim_{h \rightarrow 0^+} 3h^1 \cos\left(\frac{1}{h}\right) + h^0 \sin\left(\frac{1}{h}\right) \\
&= \lim_{h \rightarrow 0^+} 3h \cos\left(\frac{1}{h}\right) + \sin\left(\frac{1}{h}\right) \\
& \quad \lim_{h \rightarrow 0^+} \sin\left(\frac{1}{h}\right) = D.N.E \\
&\therefore \lim_{h \rightarrow 0^+} nh^{n-2} \cos\left(\frac{1}{h}\right) + h^{n-3} \sin\left(\frac{1}{h}\right) = D.N.E
\end{aligned}$$

For $n > 3$:

$$-1 \leq \cos\left(\frac{1}{h}\right) \leq 1$$

For all $h > 0 : h^{n-2} > 0$

$$-h^{n-2} \leq h^{n-2} \cos\left(\frac{1}{h}\right) \leq h^{n-2}$$

$$\lim_{h \rightarrow 0^+} -h^{n-2} \leq \lim_{h \rightarrow 0^+} h^{n-2} \cos\left(\frac{1}{h}\right) \leq \lim_{h \rightarrow 0^+} h^{n-2}$$

$$n - 2 > 0$$

$$0 \leq \lim_{h \rightarrow 0^+} h^{n-2} \cos\left(\frac{1}{h}\right) \leq 0$$

$\therefore$ by The Squeeze Theorem

$$\lim_{h \rightarrow 0^+} h^{n-2} \cos\left(\frac{1}{h}\right) = 0$$

$$-1 \leq \sin\left(\frac{1}{h}\right) \leq 1$$

For all $h > 0 : h^{n-3} > 0$

$$-h^{n-3} \leq h^{n-3} \sin\left(\frac{1}{h}\right) \leq h^{n-3}$$

$$\lim_{h \rightarrow 0^+} -h^{n-3} \leq \lim_{h \rightarrow 0^+} h^{n-3} \sin\left(\frac{1}{h}\right) \leq \lim_{h \rightarrow 0^+} h^{n-3}$$

$$n - 3 > 0$$

$$0 \leq \lim_{h \rightarrow 0^+} h^{n-3} \sin\left(\frac{1}{h}\right) \leq 0$$

$\therefore$ by The Squeeze Theorem

$$\lim_{h \rightarrow 0^+} h^{n-3} \sin\left(\frac{1}{h}\right) = 0$$

$$\begin{aligned}
& \lim_{h \rightarrow 0^+} nh^{n-2} \cos\left(\frac{1}{h}\right) + h^{n-3} \sin\left(\frac{1}{h}\right) \\
&= \left(\lim_{h \rightarrow 0^+} n\right) \left(\lim_{h \rightarrow 0^+} h^{n-2} \cos\left(\frac{1}{h}\right)\right) + \lim_{h \rightarrow 0^+} \left(h^{n-3} \sin\left(\frac{1}{h}\right)\right) \\
&= n(0) + 0 \\
&\therefore \lim_{h \rightarrow 0^+} nh^{n-2} \cos\left(\frac{1}{h}\right) + h^{n-3} \sin\left(\frac{1}{h}\right) = 0
\end{aligned}$$

The only values of n for which the left and right hand limit both exist and are equal are $n > 3, n \in \mathbb{Z}$.

$$\boxed{\therefore f' \text{ is differentiable at } x = 0 \text{ for all } n > 3, n \in \mathbb{Z}}$$

(b) **For what values of n is f' is continuous at $x = 0$?**

For f' to be continuous at $x = 0$:

$$\lim_{x \rightarrow 0} f'(x) = f'(0)$$

Using our previously derived definition of f' :

$$f'(0) = 0 \text{ when } n > 1, n \in \mathbb{Z}$$

$\lim_{x \rightarrow 0} f'(x)$ splits to right and left hand limits:

$$\begin{aligned}
& \lim_{x \rightarrow 0^-} f'(x) \\
&= \lim_{h \rightarrow 0^-} nh^{n-1}
\end{aligned}$$

$n < 1$ $n - 1 < 0$ $\lim_{h \rightarrow 0^-} nh^{n-1} = D.N.E$	$n = 1$ $\lim_{h \rightarrow 0^-} nh^{n-1}$ $= \lim_{h \rightarrow 0^-} 1h^0$ $= 1$	$n > 1$ $n - 1 > 0$ $\lim_{h \rightarrow 0^-} nh^{n-1}$ $= 0$ For our purposes, h^{n-1} is undefined for $h < 0$ when $n \notin \mathbb{Z}$ $\therefore$ This only holds for $n \in \mathbb{Z}$
---	--	---

$$\begin{aligned}
& \lim_{h \rightarrow 0^+} f'(h) \\
&= \lim_{h \rightarrow 0^+} (nh^{n-1}) \cos\left(\frac{1}{h}\right) + (h^n) \left(-\sin\left(\frac{1}{h}\right)\right) (-1h^{-2}) \\
&= \lim_{h \rightarrow 0^+} nh^{n-1} \cos\left(\frac{1}{h}\right) + h^{n-2} \sin\left(\frac{1}{h}\right)
\end{aligned}$$

For $n < 2$:

$$n - 2 < 0$$

$$\lim_{h \rightarrow 0^+} h^{n-2} \sin\left(\frac{1}{h}\right) = D.N.E$$

$$\therefore \lim_{h \rightarrow 0^+} nh^{n-1} \cos\left(\frac{1}{h}\right) + h^{n-2} \sin\left(\frac{1}{h}\right) = D.N.E$$

For $n = 2$:

$$\lim_{h \rightarrow 0^+} nh^{n-1} \cos\left(\frac{1}{h}\right) + h^{n-2} \sin\left(\frac{1}{h}\right)$$

$$= \lim_{h \rightarrow 0^+} 3h^1 \cos\left(\frac{1}{h}\right) + h^0 \sin\left(\frac{1}{h}\right)$$

$$= \lim_{h \rightarrow 0^+} 3h \cos\left(\frac{1}{h}\right) + \sin\left(\frac{1}{h}\right)$$

$$\lim_{h \rightarrow 0^+} \sin\left(\frac{1}{h}\right) = D.N.E$$

$$\therefore \lim_{h \rightarrow 0^+} nh^{n-2} \cos\left(\frac{1}{h}\right) + h^{n-3} \sin\left(\frac{1}{h}\right) = D.N.E$$

For $n > 2$:

$$-1 \leq \cos\left(\frac{1}{h}\right) \leq 1$$

$$\text{For all } h > 0 : h^{n-1} > 0$$

$$-h^{n-1} \leq h^{n-1} \cos\left(\frac{1}{h}\right) \leq h^{n-1}$$

$$\lim_{h \rightarrow 0^+} -h^{n-1} \leq \lim_{h \rightarrow 0^+} h^{n-1} \cos\left(\frac{1}{h}\right) \leq \lim_{h \rightarrow 0^+} h^{n-1}$$

$$n - 2 > 0$$

$$0 \leq \lim_{h \rightarrow 0^+} h^{n-1} \cos\left(\frac{1}{h}\right) \leq 0$$

$\therefore$ by The Squeeze Theorem

$$\lim_{h \rightarrow 0^+} h^{n-1} \cos\left(\frac{1}{h}\right) = 0$$

$$-1 \leq \sin\left(\frac{1}{h}\right) \leq 1$$

$$\text{For all } h > 0 : h^{n-2} > 0$$

$$-h^{n-2} \leq h^{n-2} \sin\left(\frac{1}{h}\right) \leq h^{n-2}$$

$$\lim_{h \rightarrow 0^+} -h^{n-2} \leq \lim_{h \rightarrow 0^+} h^{n-2} \sin\left(\frac{1}{h}\right) \leq \lim_{h \rightarrow 0^+} h^{n-2}$$

$$n - 3 > 0$$

$$0 \leq \lim_{h \rightarrow 0^+} h^{n-2} \sin\left(\frac{1}{h}\right) \leq 0$$

$\therefore$ by The Squeeze Theorem

$$\lim_{h \rightarrow 0^+} h^{n-2} \sin\left(\frac{1}{h}\right) = 0$$

$$\begin{aligned}
& \lim_{h \rightarrow 0^+} nh^{n-1} \cos\left(\frac{1}{h}\right) + h^{n-2} \sin\left(\frac{1}{h}\right) \\
&= \left(\lim_{h \rightarrow 0^+} n\right) \left(\lim_{h \rightarrow 0^+} h^{n-1} \cos\left(\frac{1}{h}\right)\right) + \lim_{h \rightarrow 0^+} \left(h^{n-2} \sin\left(\frac{1}{h}\right)\right) \\
&= n(0) + 0 \\
&\therefore \lim_{h \rightarrow 0^+} nh^{n-1} \cos\left(\frac{1}{h}\right) + h^{n-2} \sin\left(\frac{1}{h}\right) = 0
\end{aligned}$$

The only values of n for which the left and right hand limit both exist and are equal are when $n > 2, n \in \mathbb{Z}$.

Therefore:


$$\text{When } n > 2, n \in \mathbb{Z} : \quad \lim_{x \rightarrow 0} f'(x) = 0$$

$$\text{When } n > 1, n \in \mathbb{Z} : \quad f'(0) = 0$$

So $\lim_{x \rightarrow 0} f'(x) = f'(0)$ holds only when $n > 2, n \in \mathbb{Z}$.

$$\therefore f' \text{ is continuous at } x = 0 \text{ for all } n > 2, n \in \mathbb{Z}$$

4. Escaping a predator. A large fish (predator) of (vertical) size S is approaching a small fish (prey) at a constant speed v . Let $x(t)$ be the distance between the predator and the prey at time t .



- (a) Find an expression for the rate of change of the visual angle θ perceived by the prey in terms of the size and speed of the approaching predator, and the predator's distance away from its prey. The visual angle is the angle subtended by an object at the eye of the observer.

$$\begin{aligned}
\theta &= 2 \tan^{-1} \left(\frac{\frac{S}{2}}{x(t)} \right) \\
&= 2 \tan^{-1} \left(\left(\frac{S}{2} \right) x(t)^{-1} \right)
\end{aligned}$$

Since we are **approaching** at speed v , the change in distance over time would be negative v .

$$\frac{dx}{dt} = -v$$

Maybe add in a bit here about how this formula was derived, maybe an image

$$\begin{aligned}
\frac{d\theta}{dt} &= 2 \left(\left(\frac{1}{1 + \left(\left(\frac{S}{2} \right) x(t)^{-1} \right)^2} \right) \left(\frac{S}{2} \right) (-1) x(t)^{-2} \left(\frac{dx}{dt} \right) \right) \\
&= \frac{-Sx(t)^{-2}(-v)}{1 + \left(\frac{S^2}{4} \right) x(t)^{-2}} \\
&= \frac{Sx(t)^{-2}v}{\left(\frac{4x(t)^2 + S^2}{4x(t)^2} \right)} \\
&= \frac{Sx(t)^{-2}v(4x(t)^2)}{4x(t)^2 + S^2} \\
&= \frac{4Sv}{4x(t)^2 + S^2}
\end{aligned}$$

$$\therefore \frac{d\theta}{dt} = \frac{4Sv}{4x(t)^2 + S^2}$$

- (b) **As the predator approaches its prey, we expect the visual angle perceived by the prey to increase, as objects appear bigger when they are closer. Verify this is consistent with your result from part (a). Is the visual angle getting bigger faster or slower as the predator approaches its prey?**

$$\begin{aligned}
\frac{d\theta}{dt} &= \frac{4Sv}{4x(t)^2 + S^2} \\
&= (4Sv)(4x(t)^2 + S^2)^{-1}
\end{aligned}$$

Since $S > 0, v > 0, x^2 > 0$ the change in visual angle over time is always positive. In other words, the visual angle is always increasing with time. And since x is decreasing with time ($\frac{dx}{dt} < 0$), the visual angle is getting larger as the predator approaches.

Therefore, our results are consistent: as the predator approaches the prey the visual angle gets bigger.

$$\begin{aligned}
\frac{d^2\theta}{d^2t} &= (4Sv)(-1)(4x(t)^2 + S^2)^{-2}(8x(t))(-v) \\
&= (32Sv^2x(t))(4x(t)^2 + S^2)^{-2} \\
&= \frac{32Sv^2x(t)}{(4x(t)^2 + S^2)^2}
\end{aligned}$$

For all $x > 0, \frac{d^2\theta}{d^2t} > 0$. In other words $\theta(t)$ is concave up on $x > 0$.

Therefore, the visual angle is getting bigger faster as the predator approaches the prey.

- (c) **Visual angles are important factors in studies of predator avoidance. A model proposed by L. Dill, a behavioural ecologist at Simon Fraser University, about the behaviour of the Zebra Danio (a small tropical fish) suggested that “Zebra Danios react to an approaching predator when the rate of change of the angle subtended by the predator at the prey’s eye exceeds some threshold level” (L. Dill (1974) “The escape response of the Zebra Danio (*Brachydanio rerio*). I. The stimulus for escape.”,**

Animal Behaviour, 22, 711-722). Suppose an escape reaction is triggered when the rate of change of the visual angle reaches the threshold value r_{crit} . Using the model developed in part (a), find the distance x_{react} between prey and predator at which an escape response is triggered.

An escape response is triggered when the derivative of the visual angle reaches r_{crit} , at which point the predator is a distance x_{react} from the prey.

$$\begin{aligned}
 r_{\text{crit}} &= \frac{d\theta}{dt} \\
 r_{\text{crit}} &= \frac{4Sv}{4x_{\text{react}}^2 + S^2} \\
 r_{\text{crit}}(4x_{\text{react}}^2 + S^2) &= 4Sv \\
 4x_{\text{react}}^2 r_{\text{crit}} + S^2 r_{\text{crit}} &= 4Sv \\
 x_{\text{react}} &= \pm \sqrt{\frac{4Sv - S^2 r_{\text{crit}}}{4r_{\text{crit}}}}
 \end{aligned}$$

Since we only care if the prey is in front of the predator:

$$x_{\text{react}} = \sqrt{\frac{4Sv - S^2 r_{\text{crit}}}{4r_{\text{crit}}}}$$

- (d) Using the model developed in part (a) and (c), explain why for a predator of a given size S your model requires a “hunting speed” $v > \frac{Sr_{\text{crit}}}{4}$ in order to trigger an escape response by the prey.

$$x_{\text{react}} = \sqrt{\frac{4Sv - S^2 r_{\text{crit}}}{4r_{\text{crit}}}}$$

For x_{react} to exist and not be 0 (the distance at which the predator eats the prey):

$$\begin{aligned}
 \frac{4Sv - S^2 r_{\text{crit}}}{4r_{\text{crit}}} &> 0 \\
 r_{\text{crit}} &> 0 \\
 4Sv &> S^2 r_{\text{crit}} \\
 v &> \frac{S^2 r_{\text{crit}}}{4S} \\
 v &> \frac{Sr_{\text{crit}}}{4}
 \end{aligned}$$

Therefore, using the model derived in (c), for the animal to react:

$$v > \frac{Sr_{\text{crit}}}{4}$$

- (e) Again, using the model developed in part (a) and (c), for a given “hunting speed” v , what size S would not result in an escape response by the prey?

$$x_{\text{react}} = \sqrt{\frac{4Sv - S^2 r_{\text{crit}}}{4r_{\text{crit}}}}$$

For x_{react} to exist and not be 0 (the distance at which the predator eats the prey):

$$\frac{4Sv - S^2 r_{\text{crit}}}{4r_{\text{crit}}} > 0$$

$$r_{\text{crit}} > 0$$

$$S(4v - Sr_{\text{crit}}) > 0$$

$$S > 0$$

$$4v - Sr_{\text{crit}} > 0$$

$$4v > Sr_{\text{crit}}$$

$$\frac{4v}{r_{\text{crit}}} > S$$

$$S < \frac{4v}{r_{\text{crit}}}$$

To determine the speed that will **not** trigger an escape response, we can just negate the less than symbol (again ignoring the point at which $x_{\text{react}} = 0$).

Therefore, using the model derived in (c), to **not** trigger an escape response in the prey :

$$S > \frac{4v}{r_{\text{crit}}}$$

- (f) **Using your results found in parts (d) and (e), explain why large predators and slowing moving predators would have higher success rate at eating the small Zebra Danio.**

If you are slow moving enough you can sneak up on the small Zebra Danio without the change in visual angle over time reaching the critical level r_{crit} and triggering its escape response. This is because when:

$$v < \frac{Sr_{\text{crit}}}{4}$$

x_{react} does not exist, as developed in (d).

Similarly if you are large enough you can sneak up on the small Zebra Danio without the change in visual angle over time reaching the critical level r_{crit} and triggering its escape response. This is because when:

$$S > \frac{4v}{r_{\text{crit}}}$$

x_{react} does not exist, as developed in (e).

Therefore, slow predators and large predators can approach the Zebra Danio without triggering its escape response, and in turn, they would have a higher success rate at eating it.