

Quiz 5 Reflection

SCIE 001 Physics

Noah Virjee

45515863

December 12, 2025

Contents

1. Collision Ball and Box 1	2
1.1. Specify Question	2
1.2. Diagnosis Phase	2
1.3. Correction Phase	2
2. Angular Velocity of Disk	3
2.1. Specify Question	3
2.2. Diagnosis Phase	3
2.3. Correction Phase	3

1. Collision Ball and Box 1

1.1. Specify Question

Question 4: Hockey Physicist

1

2

Two identical pucks are laying flat on an icy surface and both are pulled via a thin string by a hockey player (who is a physicist at heart). The hockey player pulls with the same constant horizontal force F on both pucks for 2 seconds. In the first scenario, the thin string is wound around the outside of the puck, while in the second it is attached to its center.

a) Which puck achieves a higher speed?

Puck #2

×

0%

b) In which scenario does the hockey player do more work as they pull on the string?

Scenario #2

×

0%

This question is complete and cannot be answered again.

Correct answer

Same for both

Scenario #1

1.2. Diagnosis Phase

I treated speed as the linear speed of the puck (not the angular speed), and I incorrectly thought the energy imparted by the player would be the same in both cases.

After the pull, the first puck has rotational and linear kinetic energy, while the second puck only has linear energy.

If the energy was the same, puck 2 would move faster linearly, while puck one would be spinning while moving slower.

If work is force $\times$ distance, and force is constant, puck two would travel farther since it was moving faster, and therefore had more work done on it.

Words: 99

1.3. Correction Phase

Energy is not the same among cases, but the force, mass, and radius is, so we can start to represent the systems.

Considering from a momentum perspective:

$$F = \frac{dp}{dt}$$
$$p = \int F \, dt = Ft$$

Force is the same, so momentum is the same.

$$p_1 = p_2$$
$$p_{1x} = p_{2x}$$
$$p = mv_x$$
$$mv_{1x} = mv_{2x}$$
$$v_{1x} = v_{2x}$$

So speed is the same.

For the work, we must consider the amount of rope pulled:

For the second puck:

$$F = ma$$
$$a = \frac{F}{m}$$
$$a = \frac{dv}{dt}$$
$$v_2 = \int a \, dt = \int \frac{F}{m} \, dt = \frac{Ft}{m}$$

Work is equal to force times distance, so after 2 seconds:

$$v = \frac{dx}{dt}$$
$$x_2 = \int v_2 \, dt = \int \frac{Ft}{m} \, dt = \frac{Ft^2}{2m}$$
$$x_2(2) = \frac{F2^2}{2m} = \frac{2F}{m} = x_{\text{pulled}}$$
$$W_2 = \left| \vec{F} \times \vec{x} \right| = F(x_{\text{pulled}}) = \frac{2F^2}{m}$$

For the first puck, it is the same as the second, but we must account for the puck un-spooling.

Considering torque:

$$\tau = \left| \vec{r} \times \vec{F} \right| = rF$$
$$\tau = I\alpha$$
$$\alpha = \frac{\tau}{I} = \frac{rF}{I}$$

Treating the puck as a disk:

$$I = \frac{1}{2}mr^2$$
$$\alpha = \frac{rF}{\frac{1}{2}mr^2} = 2Fm^{-1}r^{-1}$$
$$\alpha = \frac{d\omega}{dt}$$
$$\omega = \int \alpha \, dt = \int 2Fm^{-1}r^{-1} \, dt = 2Fm^{-1}r^{-1}t$$
$$v_{\text{un-spooling}} = \omega r = 2Fm^{-1}r^{-1}tr = 2Fm^{-1}t$$

So the rope velocity is:


$$v_{\text{un-spooling}} = 2\left(\frac{Ft}{m}\right)$$
$$v_{\text{pulling}} = v_{\text{linear}} + v_{\text{un-spooling}}$$
$$v_{\text{pulling}} = \frac{Ft}{m} + 2\left(\frac{Ft}{m}\right) = \frac{3Ft}{m} \quad \text{with } v_{\text{linear}} = \frac{Ft}{m} \text{ from 2}$$
$$x_{\text{pulled}} = \int v_{\text{pulling}} \, dt = \int \frac{3Ft}{m} \, dt = \frac{3Ft^2}{2m}$$
$$x_{\text{pulled}}(2) = \frac{3F2^2}{2m} = \frac{6F}{m}$$
$$W_1 = \left| \vec{F} \times \vec{x} \right| = F(x_{\text{pulled}}) = \frac{6F^2}{m}$$
$$\frac{6F^2}{m} > \frac{2F^2}{m}$$
$$\therefore W_1 > W_2$$

Words: 99

2. Angular Velocity of Disk

2.1. Specify Question

Question 5: Angular Velocity of Disk



A solid uniform disk with mass $M = 10\text{ kg}$ and radius $R = 30\text{ cm}$ is attached to a fixed axle going through its center of mass. It is initially at rest when the following forces $F_1 = 16\text{ N}$, $F_2 = 20\text{ N}$ and $F_3 = 32\text{ N}$ are applied to it. If $\theta = 30^\circ$, what is the disk's angular velocity after 2 s (in rad/s)?

Assume that "out of the page" is positive and give your answer to 2 decimal places.

$\omega =$

42.666 rad / s

?

✖

0%

This question is complete and cannot be answered again.

Correct answer

$\omega = 0.0$ radian / second

2.2. Diagnosis Phase

I misread the diagram and mistakenly thought that the $\vec{F}_3$ was pointing up instead of down so when I went to calculate velocity I got:

$$\vec{\tau} = \vec{F} \times \vec{R}$$
$$\tau_1 = F_1 R \sin(90^\circ) = (16\text{N})(0.3\text{m})(1) = 4.8\text{N}$$
$$\tau_2 = F_2 R \sin(180^\circ) = (20\text{N})(0.3\text{m})(0) = 0\text{ N/m}$$

I incorrectly thought the $\vec{F}_3$ was pointing up, so I calculated its torque using $\sin(180^\circ - 30^\circ)$

$$\tau_3 = F_3 R \sin(180 - 30^\circ) = (32\text{N})(0.3\text{m})(0.5) = 4.8\text{ N/m}$$
$$\tau_{\text{total}} = \tau_1 + \tau_2 + \tau_3 = 9.6\text{ Nm}$$
$$\tau = I\alpha$$

As a disk:

$$I = \frac{1}{2}mr^2$$
$$\alpha = \frac{\tau}{I} = \frac{2\tau}{mr^2}$$
$$\alpha = \frac{d\omega}{dt}$$
$$\omega = \int \alpha \, dt = \int \frac{2\tau}{mr^2} \, dt = \frac{2\tau t}{mr^2}$$
$$\omega(2) = \frac{2(9.6)(2)}{(10)(0.3)^2} = 42.67$$

Words: 40

2.3. Correction Phase

Interpreting the direction of $\vec{F}_3$ correctly.

$$\tau = \vec{F} \times R$$
$$\tau_1 = \vec{F}_1 R \sin(90^\circ) = (16\text{N})(32\text{ cm})(1) = 0$$

$90^\circ + \theta = 120^\circ$ “ is the angle between radius and force

$$\tau_2 = \vec{F}_2 R \sin(120^\circ) = (20\text{N})(32\text{ cm})$$
$$\tau_3 = \vec{F}_1 R \sin(90^\circ)$$
$$\tau_{\text{total}} = \tau_1 + \tau_2 + \tau_3 = ()$$

Words: 12