## 0.1 Ring of Fractions

**Theorem 0.1.1.** Let $R$ be a Commutative Ring, $D \subset R$ be a subset such that $\begin{cases} \text{no zero, no zero divisors} \\ \text{closed under multiplication} \end{cases}$.

Then, there exists a Commutative Ring $Q$ with identity satisfies:

1. $R$ can be embedded in $Q$, and every element of $D$ becomes unit in $Q$.
2. $Q$ is the smallest Ring containg $R$ with identity such that every element of $D$ becomes unit in $Q$.

*Proof.* Let $\mathcal{F} \overset{\text{def}}{=} \{(r,d) \mid r \in R,\ d \in D\}$ and the relation $\sim$ on $\mathcal{F}$ by $(r_1,d_1) \sim (r_2,d_2) \iff r_1d_2 = r_2d_1$.
Then, $\sim$ is equivalent relation: reflexive and symmetirc are clear, and Suppose that $(r_1,d_1) \sim (r_2,d_2)$ and $(r_2,d_2) \sim (r_3,d_3)$.

$$r_2d_3 = r_3d_2 \implies r_2d_1d_3 = r_3d_1d_2 \implies r_1d_2d_3 = r_3d_1d_2 \implies d_2(r_1d_3 - r_3d_1) \implies r_1d_3 = r_3d_1$$

Thus transitivity shown. Define

$$\frac{r}{d} \overset{\text{def}}{=} [(r,d)] = \{(a,b) \mid (a,b) \sim (r,d)\},\ Q \overset{\text{def}}{=} \left\{ \frac{r}{d} \,\middle|\, r \in R,\ d \in D \right\}$$

And define operations $+, \times$ on $Q$:

$$\frac{r_1}{d_1} + \frac{r_2}{d_2} \overset{\text{def}}{=} \frac{r_1d_2 + r_2d_1}{d_1d_2}, \quad \frac{r_1}{d_1} \times \frac{r_2}{d_2} \overset{\text{def}}{=} \frac{r_1r_2}{d_1d_2}$$

Well-Definedness: If $\dfrac{r_1}{d_1} = \dfrac{r_1'}{d_1'}$ and $\dfrac{r_2}{d_2} = \dfrac{r_2'}{d_2'}$,

$$\frac{r_1d_2 + r_2d_1}{d_1d_2} = \frac{r_1d_2d_1'd_2' + r_2d_1d_1'd_2'}{d_1d_2d_1'd_2'} = \frac{(r_1d_1')d_2d_2' + (r_2d_2')d_1d_1'}{d_1d_2d_1'd_2'} = \frac{(r_1'd_1)d_2d_2' + (r_2'd_2)d_1d_1'}{d_1d_2d_1'd_2'} = \frac{(r_1'd_2' + r_2'd_1')d_1d_2}{d_1d_2d_1'd_2'} = \frac{r_1'd_2' + r_2'd_1'}{d_1'd_2'}$$

$$\frac{r_1r_2}{d_1d_2} = \frac{r_1r_2d_1'd_2'}{d_1d_2d_1'd_2'} = \frac{(r_1d_1')(r_2d_2')}{d_1d_2d_1'd_2'} = \frac{(r_1'd_1)(r_2'd_2)}{d_1d_2d_1'd_2'} = \frac{r_1'r_2'd_1d_2}{d_1d_2d_1'd_2'} = \frac{r_1'r_2'}{d_1'd_2'}$$

Now, $(Q,+,\times)$ constructs Commutative Ring with identity: for any $d \in D$, put $0_Q \overset{\text{def}}{=} \dfrac{0}{d},\ 1_Q \overset{\text{def}}{=} \dfrac{d}{d}$. Then,

1. $(R,+,\times)$ closed under the operations since $D$ is closed under the multiplication.
2. $(R,+)$ has a zero: $\dfrac{r_1}{d_1} + 0_Q = \dfrac{r_1}{d_1} + \dfrac{0}{d} = \dfrac{r_1d + 0d_1}{d_1d} = \dfrac{r_1d}{d_1d} = \dfrac{r_1}{d_1}$.
3. $(R,+)$ has an inverse: $\dfrac{r_1}{d_1} + \dfrac{-r_1}{d_1} = \dfrac{r_1d_1 + (-r_1)d_1}{d_1d_1} = \dfrac{[(r_1) + (-r_1)]d_1}{d_1d_1} = \dfrac{0d_1}{d_1d_1} = \dfrac{0}{d_1d_1} = 0_Q$.
4. $(R,+,\times)$ satisfies distributive law:

   4-1. The left law:

$$\begin{aligned} \frac{r_1}{d_1} \times \left( \frac{r_2}{d_2} + \frac{r_3}{d_3} \right) &= \frac{r_1}{d_1} \times \frac{r_2d_3 + r_3d_2}{d_2d_3} = \frac{r_1r_2d_3 + r_1r_3d_2}{d_1d_2d_3} = \frac{r_1r_2d_1d_3 + r_1r_3d_1d_2}{d_1d_2d_1d_3} = \frac{r_1r_2}{d_1d_2} + \frac{r_2r_3}{d_2d_3} \\ &= \frac{r_1}{d_1} \times \frac{r_2}{d_2} + \frac{r_2}{d_2} \times \frac{r_3}{d_3} \end{aligned}$$

   4-2. The right law:

$$\begin{aligned} \left( \frac{r_1}{d_1} + \frac{r_2}{d_2} \right) \times \frac{r_3}{d_3} &= \frac{r_1d_2 + r_2d_1}{d_1d_2} \times \frac{r_3}{d_3} = \frac{r_1r_3d_2 + r_2r_3d_1}{d_1d_2d_3} = \frac{r_1r_3d_2d_3 + r_2r_3d_1d_3}{d_1d_3d_2d_3} = \frac{r_1r_3}{d_1d_3} + \frac{r_2r_3}{d_2d_3} \\ &= \frac{r_1}{d_1} \times \frac{r_3}{d_3} + \frac{r_2}{d_2} \times \frac{r_3}{d_3} \end{aligned}$$

5. $(R,\times)$ has an identity: $\dfrac{r_1}{d_1} \times 1_Q = \dfrac{r_1}{d_1} \times \dfrac{d}{d} = \dfrac{r_1d}{d_1d} = \dfrac{r_1}{d_1}$.
6. Elements of $D$ become unit in $Q$: Define $\iota : R \to Q : r \mapsto \dfrac{rp}{p}$ where $p \in D$ is any fixed element in $D$.
   Then, $\iota$ is Ring-Monomorphsim because:

6-1. Well-Defined and Injective: $\iota(r_1) = \iota(r_2) \iff \dfrac{r_1 p}{p} = \dfrac{r_2 f}{f} \iff (r_1 - r_2)pp = 0 \iff r_1 = r_2$

6-2. For any $d \in D$, $\iota(d)$ is a unit of $Q$: Put $(\iota(d))^{-1} \overset{\text{def}}{=} \dfrac{p}{dp}$, then

$$\iota(d) \times (\iota(d))^{-1} = \frac{dp}{p} \times \frac{p}{dp} = \frac{dpp}{dpp} = 1_Q$$

That is, $\iota$ is embedding from $R$ into $Q$ such that $\iota[D]$ becomes units of $Q$ except zero.
Moreover, if $D = R \setminus \{0\}$, then $Q$ is field.

7. $Q$ is the *smallest* ring containing $R$ with identity such that every element of $D$ becomes units in $Q$.
Let $S$ be an any commutative ring with identity,
and assume that $\varphi : R \to S$ is a Ring-Monomorphism such that for any $d \in D$, $\varphi(d)$ is unit in $S$.
Define $\phi : Q \to S : \dfrac{r}{d} \mapsto \varphi(r)\varphi(d)^{-1}$. Then, this $\phi$ is well-defined and injective:

$$\phi\left(\frac{r_1}{d_1}\right) = \phi\left(\frac{r_2}{d_2}\right) \iff \varphi(r_1)\varphi(d_1)^{-1} = \varphi(r_2)\varphi(d_2)^{-1} \iff \varphi(r_1)\varphi(d_2) = \varphi(r_2)\varphi(d_1)$$
$$\overset{\text{homomor.}}{\iff} \varphi(r_1 d_2) = \varphi(r_2 d_1) \overset{\text{one-to-one}}{\iff} r_1 d_2 = r_2 d_1 \iff \frac{r_1}{d_1} = \frac{r_2}{d_2}$$

That is, if a commutative ring $S$ with identity contains a copy of $R$ such that the denominator set $D$ of $R$ becomes unit in $S$, then $S$ contains ring of fractions $Q$ of $R$. Thus $S = Q$ is the smallest ring that satisfies these conditions.

□