

Galois group of $(x^2-2)(x^2-3)$ over $\mathbb{Q}$.

Since the polynomial $(x^2-2)(x^2-3)$ has four distinct roots $\pm\sqrt{2}, \pm\sqrt{3}$, it is separable.

And, since $\sqrt{3} \notin \mathbb{Q}(\sqrt{2})$, $[\mathbb{Q}(\sqrt{2}, \sqrt{3}) : \mathbb{Q}] = [\mathbb{Q}(\sqrt{2}, \sqrt{3}) : \mathbb{Q}(\sqrt{3})] \cdot [\mathbb{Q}(\sqrt{3}) : \mathbb{Q}] = 4$.

So, $|\text{Aut}(K/\mathbb{Q})| = [K : \mathbb{Q}] = 4$ where $K = \mathbb{Q}(\sqrt{2}, \sqrt{3})$, the splitting field over $\mathbb{Q}$.

Since the automorphism leaving $\mathbb{Q}$ fixed must be conjugate the roots,

there are four possibilities:

$$\begin{cases} \sqrt{2} \mapsto \sqrt{2} \\ \sqrt{3} \mapsto \sqrt{3} \end{cases}, \begin{cases} \sqrt{2} \mapsto -\sqrt{2} \\ \sqrt{3} \mapsto \sqrt{3} \end{cases}, \begin{cases} \sqrt{2} \mapsto \sqrt{2} \\ \sqrt{3} \mapsto -\sqrt{3} \end{cases}, \begin{cases} \sqrt{2} \mapsto -\sqrt{2} \\ \sqrt{3} \mapsto -\sqrt{3} \end{cases}.$$

Let $\sigma : K \rightarrow K; a + b\sqrt{2} + c\sqrt{3} + d\sqrt{6} \mapsto a - b\sqrt{2} + c\sqrt{3} - d\sqrt{6}$


$\tau : K \rightarrow K; a + b\sqrt{2} - c\sqrt{3} - d\sqrt{6} \mapsto a + b\sqrt{2} - c\sqrt{3} - d\sqrt{6}.$

(That is, $\sigma : \begin{cases} \sqrt{2} \mapsto -\sqrt{2} \\ \sqrt{3} \mapsto \sqrt{3} \end{cases}$ and $\tau : \begin{cases} \sqrt{2} \mapsto \sqrt{2} \\ \sqrt{3} \mapsto -\sqrt{3} \end{cases}$, and we can observe that

$$\sigma\tau : \begin{cases} \sqrt{2} \mapsto -\sqrt{2} \\ \sqrt{3} \mapsto -\sqrt{3} \end{cases}.$$

Now, $\text{Gal}(K/\mathbb{Q}) = \{1, \sigma, \tau, \sigma\tau\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$, corresponding σ to $(1, 2)$ and τ to $(2, 1)$.

Moreover,



Galois group of x^3-2 over $\mathbb{Q}$.

Denote $\zeta_3 = e^{i\frac{2\pi}{3}} = \frac{1}{2}(-1 + i\sqrt{3})$, the 3rd primitive root of unity.

Since $\zeta_3^3 = 1$, the set $\{\sqrt[3]{2}, \sqrt[3]{2}\zeta_3, \sqrt[3]{2}\zeta_3^2\}$ are all roots of x^3-2 over $\mathbb{Q}$.
So, the splitting field is $K = \mathbb{Q}(\sqrt[3]{2}, \zeta_3)$.

Observe that: the minimal polynomial of $\sqrt[3]{2}$ over $\mathbb{Q}$ is x^3-2 , and

" " ζ_3 over $\mathbb{Q}$ is x^2+x+1 .

And, since x^2+x+1 is irreducible over $\mathbb{Q}(\sqrt[3]{2})$, (it cannot have root in $\mathbb{Q}(\sqrt[3]{2})$)

$$[\mathbb{Q}(\sqrt[3]{2}, \zeta_3) : \mathbb{Q}] = [\mathbb{Q}(\sqrt[3]{2}, \zeta_3) : \mathbb{Q}(\sqrt[3]{2})] \cdot [\mathbb{Q}(\sqrt[3]{2}) : \mathbb{Q}] = 2 \cdot 3 = 6.$$

the tower lemma gave. And, since x^3-2 is separable, $|\text{Aut}(K/\mathbb{Q})| = [K:\mathbb{Q}] = 6$.

Now, construct explicitly the $\text{Aut}(K/\mathbb{Q})$.

Since any automorphism of K leaving $\mathbb{Q}$ fixed must conjugate the roots,

it must map $\sqrt[3]{2}$ to $\sqrt[3]{2}$ or $\sqrt[3]{2}\zeta_3$ or $\sqrt[3]{2}\zeta_3^2$

and maps ζ_3 to ζ_3 or ζ_3^2 .

So, there are exactly 6 possibilities. Consider two automorphisms

$$\sigma \text{ satisfying } \sigma(\sqrt[3]{2}) = \sqrt[3]{2}\zeta_3 \text{ and } \sigma(\zeta_3) = \zeta_3$$

and τ satisfying $\tau(\sqrt[3]{2}) = \sqrt[3]{2}$ and $\tau(\zeta_3) = \zeta_3^2 = -1 - \zeta_3$.

$$\text{Then, } \sigma^3: \begin{cases} \sqrt[3]{2} \mapsto \sqrt[3]{2}\zeta_3 \mapsto \sqrt[3]{2}\zeta_3^2 \mapsto \sqrt[3]{2}\zeta_3^3 = \sqrt[3]{2} \\ \zeta_3 \mapsto \zeta_3 \end{cases}, \tau^2: \begin{cases} \sqrt[3]{2} \mapsto \sqrt[3]{2} \\ \zeta_3 \mapsto \zeta_3^2 \mapsto \zeta_3^4 = 1 - \zeta_3 = \zeta_3 \end{cases}$$

are actually identities. Moreover,

$$\sigma^2: \begin{cases} \sqrt[3]{2} \mapsto \sqrt[3]{2}\zeta_3^2 \\ \zeta_3 \mapsto \zeta_3 \end{cases}, \sigma\tau: \begin{cases} \sqrt[3]{2} \mapsto \sqrt[3]{2}\zeta_3 \\ \zeta_3 \mapsto \zeta_3^2 \end{cases}, \sigma^2\tau: \begin{cases} \sqrt[3]{2} \mapsto \sqrt[3]{2}\zeta_3^2 \\ \zeta_3 \mapsto \zeta_3^2 \end{cases}$$

So, $\text{Aut}(K/\mathbb{Q}) = \{1, \sigma, \sigma^2, \tau, \sigma\tau, \sigma^2\tau\} \cong S_3$, corresponding σ to $(1\ 2\ 3)$, τ to $(1\ 2)$