

0.0.1 Inequality of Riemann-Stieltjes Integral

Let $p, q \geq 1$ such that $\frac{1}{p} + \frac{1}{q} = 1$, and functions be defined on $[a, b]$.

Lemma 0.0.1. Let $f, g \in \mathcal{R}(\alpha)$ with $f, g \geq 0$, and $\int_a^b [f(x)]^p d\alpha = \int_a^b [g(x)]^q d\alpha = 1$. Then, $\int_a^b f(x)g(x) d\alpha \leq 1$.

Proof. For any $x \in [a, b]$, the Young's Inequality gives

$$0 \leq f(x)g(x) \leq \frac{[f(x)]^p}{p} + \frac{[g(x)]^q}{q}$$

Now,

$$\int_a^b f(x)g(x) d\alpha \leq \int_a^b \frac{[f(x)]^p}{p} + \frac{[g(x)]^q}{q} d\alpha = \frac{1}{p} \int_a^b [f(x)]^p d\alpha + \frac{1}{q} \int_a^b [g(x)]^q d\alpha = \frac{1}{p} + \frac{1}{q} = 1$$

□

Definition 0.0.2. Let $f \in \mathcal{R}(\alpha)$. Define a Norm of f :

$$\|f\|_p \stackrel{\text{def}}{=} \left(\int_a^b |f(x)|^p d\alpha \right)^{\frac{1}{p}}$$

It is a Norm of a set of Stieltjes Integrable functions, $\mathcal{F} \stackrel{\text{def}}{=} \{f : [a, b] \rightarrow \mathbb{C} \mid f \in \mathcal{R}(\alpha)\}$.

Lemma 0.0.3. Hölder's Inequality

Let $f, g \in \mathcal{F}$. Then,

$$\left| \int_a^b f(x)g(x) d\alpha \right| \leq \left[\int_a^b |f(x)|^p d\alpha \right]^{\frac{1}{p}} \cdot \left[\int_a^b |g(x)|^q d\alpha \right]^{\frac{1}{q}}$$

Proof. Rewrite:

$$\|f\|_p^p = \int_a^b |f(x)|^p d\alpha, \quad \|g\|_q^q = \int_a^b |g(x)|^q d\alpha$$

Now, we can make the condition of above lemma,

$$\int_a^b \left[\frac{|f(x)|}{\|f\|_p} \right]^p d\alpha = \frac{1}{\|f\|_p^p} \cdot \int_a^b |f(x)|^p d\alpha = 1, \quad \int_a^b \left[\frac{|g(x)|}{\|g\|_q} \right]^q d\alpha = \frac{1}{\|g\|_q^q} \cdot \int_a^b |g(x)|^q d\alpha = 1$$

And apply this,

$$\int_a^b \frac{|f(x)| \cdot |g(x)|}{\|f\|_p \|g\|_q} d\alpha \leq 1 \implies \int_a^b |f(x)| |g(x)| d\alpha \leq \|f\|_p \|g\|_q = \left[\int_a^b |f(x)|^p d\alpha \right]^{\frac{1}{p}} \cdot \left[\int_a^b |g(x)|^q d\alpha \right]^{\frac{1}{q}}$$

Finally, the general property of integral of product gives

$$\left| \int_a^b f(x)g(x) d\alpha \right| \leq \int_a^b |f(x)| |g(x)| d\alpha \leq \|f\|_p \|g\|_q = \left[\int_a^b |f(x)|^p d\alpha \right]^{\frac{1}{p}} \cdot \left[\int_a^b |g(x)|^q d\alpha \right]^{\frac{1}{q}}$$

□

Theorem 0.0.4. Minkowski inequality

Let $f, g \in \mathcal{F}$. Then, for any $p \geq 1$, $\|f + g\|_p \leq \|f\|_p + \|g\|_p$.

Proof.

$$\begin{aligned}
\|f + g\|_p^p &= \int_a^b |f + g|^p d\alpha = \int_a^b |f + g| |f + g|^{p-1} d\alpha \\
&\leq \int_a^b (|f| + |g|) |f + g|^{p-1} d\alpha \\
&= \int_a^b |f| |f + g|^{p-1} d\alpha + \int_a^b |g| |f + g|^{p-1} d\alpha \\
&\stackrel{\text{H\"older}}{\leq} \left[\int_a^b |f|^p d\alpha \right]^{\frac{1}{p}} \left[\int_a^b |f + g|^{(p-1)\frac{p}{p-1}} d\alpha \right]^{\frac{p-1}{p}} + \left[\int_a^b |g|^p d\alpha \right]^{\frac{1}{p}} \left[\int_a^b |f + g|^{(p-1)\frac{p}{p-1}} d\alpha \right]^{\frac{p-1}{p}} \\
&= \left[\int_a^b |f + g|^p d\alpha \right]^{\frac{p-1}{p}} \left(\left[\int_a^b |f|^p d\alpha \right]^{\frac{1}{p}} + \left[\int_a^b |g|^p d\alpha \right]^{\frac{1}{p}} \right) = \|f + g\|_p^{p-1} \cdot (\|f\|_p + \|g\|_p)
\end{aligned}$$

Now,

$$\|f + g\|_p^p \cdot \|f + g\|_p^{1-p} = \|f + g\|_p \leq \|f\|_p + \|g\|_p$$

□