

† Singular value decomposition.

Let  $V, W$  be inner product space and let  $T: V \rightarrow W$  be a linear.

Then, there exists orthonormal basis  $\{v_1, \dots, v_n\}$  and  $\{u_1, \dots, u_n\}$  of  $V, W$ , respectively and positive scalars  $\sigma_1 \geq \dots \geq \sigma_r$  s.t

$$T(v_i) = \begin{cases} \sigma_i u_i & (1 \leq i \leq \text{rank } T) \\ 0 & (i > \text{rank } T) \end{cases}$$

Conversely,

proof. Let  $x \in \ker(T^* \circ T)$ .  $(T^* \circ T)(x) = 0$  implies

$$0 = \langle 0, x \rangle = \langle (T^* \circ T)x, x \rangle = \langle T(x), T(x) \rangle \Rightarrow T(x) = 0.$$

Therefore,  $\ker T^* \subseteq \ker T$ , and the reverse inclusion is clear.

By the Dimension Thm,  $\text{rank } T = \text{rank } T^* T$ . Moreover, for any non-zero  $x \in V$ ,

$$\langle T^* T(x), x \rangle = \langle T(x), T(x) \rangle = \|T(x)\|^2 \geq 0$$

Thus  $T^* T$  is positive semidefinite. Equivalently every eigenvalue of  $T^* T$  is non-negative and has an orthonormal basis of eigenvectors. Let  $r = \text{rank } T$ . Then,

denote the positive eigenvalues as

$$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_r > 0$$

and corresponding eigenvectors as

$$\{v_1, \dots, v_r, \dots, v_n\}$$

For each  $i$ , define  $\sigma_i = \sqrt{\lambda_i}$  and  $u_i = \frac{1}{\sigma_i} T(v_i)$ . Then for any  $1 \leq i, j \leq r$ ,

$$\langle u_i, u_j \rangle = \left\langle \frac{1}{\sigma_i} T(v_i), \frac{1}{\sigma_j} T(v_j) \right\rangle = \frac{1}{\sigma_i \sigma_j} \langle T^* T(v_i), v_j \rangle = \frac{\sigma_i^2}{\sigma_i \sigma_j} \langle v_i, v_j \rangle = \delta_{i,j}.$$

Therefore  $\{u_1, \dots, u_r\}$  is orthonormal. Extending this to  $\{u_1, \dots, u_n\}$ , orthonormal basis for  $W$ .

And  $T^* T(v_i) = 0 \Rightarrow T(v_i) = 0$  was proven.

⌈ Singular Value Decomposition.

Let  $A \in M_{m,n}$  with  $\text{rank } A = r$ , with singular values  $\sigma_1 \geq \dots \geq \sigma_r$ .

Define a Matrix in  $M_{m,n}$  as

$$\Sigma = \begin{cases} \sigma_i & \text{if } i \leq r \\ 0 & \text{otherwise} \end{cases}$$

Then, there exists unitary  $U \in M_{m,m}$  and unitary  $V \in M_{n,n}$  s.t.

$$A = U \Sigma V^*$$

$$U = \begin{pmatrix} u_1 & \dots & u_m \end{pmatrix}, \quad V = \begin{pmatrix} v_1 & \dots & v_n \end{pmatrix}$$

⌈ The Polar Decomposition.

For any  $A \in M_{n,n}$ , there exists unitary  $W$  and  $P \geq 0$  s.t.

$$A = WP$$

Furthermore, if  $A$  is invertible, this representation is unique.

Proof. By the SVD, there exists unitary  $U, V$  and diagonal  $\Sigma \geq 0$  s.t.

$$A = U \Sigma V = \underbrace{UV}_{W} \underbrace{V^* \Sigma V}_P$$

To show uniqueness, suppose  $A$  is invertible and

$$A = W_1 P_1 = W_2 P_2 \quad W_1, W_2 \text{ unitary, } P_1, P_2 \geq 0$$

Then,  $W_2^* W_1 = P_2 P_1^{-1}$  unitary,

$$I = (P_2 P_1^{-1})^* (P_2 P_1^{-1}) = P_1^{-1} P_2^2 P_1^{-1} \Rightarrow P_1^2 = P_2^2 \Rightarrow P_1 = P_2.$$

