[Lemma.
Let K/F be a field extension, and let $\alpha \in K$ be algebraic over F. Then,
For any $\sigma \in Aut(K/F)$, $\sigma\alpha$ is a root of the minimal polynomial for α over F.

Proof. Let $f(x) = x^n + a_{n-1}x^{n-1} + \cdots + a_1 x + a_0 \in F[x]$ be the minimal polynomial for α.
Then, $f(\alpha) = 0$, so $\sigma(f(\alpha)) = 0$. (The field Homomorphism preserves zero).
Now, $0 = \sigma(f(\alpha)) = \sigma\left(\alpha^n + a_{n-1}\alpha^{n-1} + \cdots + a_1\alpha + a_0\right)$

σ fixes F $\leftarrow$ $= (\sigma\alpha)^n + \sigma(a_{n-1})\cdot(\sigma\alpha)^{n-1} + \cdots + \sigma(a_1)\cdot(\sigma\alpha) + \sigma(a_0)$
$= (\sigma\alpha)^n + a_{n-1}(\sigma\alpha)^{n-1} + \cdots + a_1\cdot(\sigma\alpha) + a_0$
$= f(\sigma\alpha)$.

[Prop. Let $H \leq Aut(K)$ be a subgroup of the group of all automorphisms of K.
Then, the collection F of elements of K fixed by all the elements of H is a subfield.
Proof. First, given $\sigma \in Aut(K)$, the set $F_\sigma = \{a \in K \mid \sigma(a) = a\}$ is subfield, because:
Let $a, b \in F_\sigma$ be given. Then, $\sigma(a-b) = \sigma(a) - \sigma(b) = a - b$, and if $b \neq 0$,
then $\sigma(a \cdot b^{-1}) = \sigma(a)\cdot\sigma(b)^{-1} = ab^{-1}$, so the claim is proved.
And now, since the arbitrary intersection of fields is field, so Proved. ∎
Such a subfield is called the fixed field of $S \subseteq Aut(K)$.

[Prop. Let K be a field. Then,
If $F_1 \leq F_2 \leq K$, then $Aut(K/F_2) \leq Aut(K/F_1)$.
If $H_1 \leq H_2 \leq Aut(K)$, then $F_{H_2} \leq F_{H_1}$.
In other say, The inclusion relation of automorphisms and fixed field is reverse.

Lemma.

Def. A finite extension K/F is called a Galois Extension if:

$$|Aut(K/F)| = [K:F].$$

In such a case, denote $Gal(K/F) = Aut(K/F)$.

Def. A character χ of a group G with values in a field L is a group homomorphism

$$\chi : G \to L^\times$$

The characters $\chi_1, \cdots, \chi_n$ of G are said to be linearly independent over L if:

$a_1\chi_1 + \cdots + a_n\chi_n \equiv 0$ for some $a_i \in L$, then $a_1 = \cdots = a_n = 0$.

Thm. [Dedekind - Independence of Characters Theorem],
If $\chi_1, \cdots, \chi_n : G \to L^\times$ are distince characters of G with values in L,
then they are linearly independent over L.

Proof. Suppose that the characters are linearly dependent.
That is, there exists non-trivial representation; for some $a_1, \cdots, a_n \in L$,

$$a_1\chi_1 + \cdots + a_n\chi_n \equiv 0 \quad \text{where } a_1, \cdots, a_n \text{ are not all zero.}$$

So, the set $\{m \in \mathbb{N} \mid$ for some $b_1, \cdots, b_m \in L\setminus\{0\}$, and $\sigma \in S_n$, $b_1\chi_{\sigma(1)} + \cdots + b_m\chi_{\sigma(m)} \equiv 0\}$
is non-empty, thus it has a minimal element by the well-ordering principle,

$m \in \mathbb{N}$ with $b_1\chi_1 + \cdots + b_m\chi_m \equiv 0$, the indices are renumbered, $b_i \in L\setminus\{0\}$.

Since the characters are distinct, $\chi_1(g_0) \neq \chi_m(g_0)$, for some $g_0 \in G$.
Given $g \in G$, observe that

$$0 = b_1\chi_1(g_0g) + \cdots + b_m\chi_m(g_0g) = b_1\chi_1(g_0)\chi_1(g) + \cdots + b_m\chi_m(g_0)\chi_m(g)$$

Meanwhile,

$$0 = \chi_m(g_0)\cdot 0 = \chi_m(g_0)\cdot(b_1\chi_1(g) + \cdots + b_m\chi_m(g)) = b_1\chi_m(g_0)\chi_1(g) + \cdots + b_m\chi_m(g_0)\chi_m(g),$$

Substracting gives $[\chi_m(g_0) - \chi_1(g_0)]\cdot b_1\chi_1(g) + \cdots + [\chi_m(g_0) - \chi_{m-1}(g_0)]\cdot b_{m-1}\chi_{m-1}(g_0) = 0$,
because the the last term is terminated. But, since $[\chi_m(g_0) - \chi_1(g_0)] \neq 0$,
it is another representation, but that is contradiction to the minimality of m. ∎

Corollary. Distinct Field Automorphisms are linearly independent.

[Thm.
Let $G=\{\sigma_1(=id), \sigma_2, \cdots, \sigma_n\}$ be a subgroup of $Aut(K)$, and let $F=K^G$.
Then, $[K:F]=|G|=n$.

Proof. To show the equality, we will show that neither $|G|>[K:F]$ nor $|G|<[K:F]$.
Suppose that $n>[K:F]$ and let $w_1, \cdots, w_m$ be a basis for K over F. (So, $m=[K:F]$)
Consider a system

$$\begin{cases} \sigma_1(w_1)x_1+\sigma_2(w_1)x_2+\cdots+\sigma_n(w_1)x_n=0 \\ \vdots \\ \sigma_1(w_m)x_1+\sigma_2(w_m)x_2+\cdots+\sigma_n(w_m)x_n=0 \end{cases} \quad \left(\text{or } \left(\sigma_j(w_i)\right)_{i,j}\begin{pmatrix}x_1\\ \vdots \\ x_n\end{pmatrix}=0\right)$$

By the assumption, $n>m$, so the system has a non-trivial solution $\beta_1, \cdots, \beta_n$ in K.
(By the dimension theorem, nullity $A=n-\text{rank } A\geq n-m>0$.)
Given $a_1, \cdots, a_m \in F$, these are fixed by $\sigma_i \in G$, by the definition of F.
Multiplying a_i to the ith row of the equation, then

$$\sigma_1(a_1w_1)\beta_1+\cdots+\sigma_n(a_1w_1)\beta_n=0$$
$$\vdots$$
$$\sigma_1(a_nw_n)\beta_1+\cdots+\sigma_n(a_nw_n)\beta_n=0.$$

So, adding all the equations i for $\alpha=a_1w_1+\cdots+a_nw_n$,

$$\beta_1\sigma_1(\alpha)+\cdots+\beta_n\sigma_n(\alpha)=0.$$

Since $a_1, \cdots, a_m$ was arbitrary and $w_1, \cdots, w_m$ forms basis for K over F, the α is arbitrary element of K. Moreover, since $\beta_1, \cdots, \beta_n$ are not all zero, it contradicts the G is linearly independent, by the dedekind character Theorem.]
Now, suppose that $n<[K:F]$. That is, there are more than n F-linearly independent elements, say $\alpha_1, \cdots, \alpha_{n+1}$. Consider a system of $n+1$ unknowns,

$$\begin{cases} \sigma_1(\alpha_1)x_1+\cdots+\sigma_1(\alpha_{n+1})x_{n+1}=0 \\ \vdots \\ \sigma_n(\alpha_1)x_1+\cdots+\sigma_n(\alpha_{n+1})x_{n+1}=0 \end{cases}$$

. Then, it has n equation and $n+1$ unknowns
So there is non-trivial solution $\beta_1, \cdots, \beta_{n+1}$ in K.
Moreover, since $\sigma_1=id$ and $\alpha_1, \cdots, \alpha_{n+1}$ are F-linearly independent, not all β_i lies in F.

Corollary. Let K/F be any finite extension. Then,

$$|Aut(K/F)| \leq [K:F]$$

In Particular, the equality holds if and only if $F = K^{Aut(K/F)}$
if and only if K/F is Galois extension
if and only if K is splitting field of separable polynomial over F.

Proof.

[Corollary.
Let G be a finite subgroup of $Aut(K)$. Then

$$Aut(K/K^G) = G.$$

And, if $G_1 \neq G_2$, then $K^{G_1} \neq K^{G_2}$.

Fundamental Theorem of Galois Theory.

Let K/F be a Galois Extension. Denote $G = Gal(K/F)$.

1). A map

$$\Phi: \{H \le Gal(K/F) \mid H \text{ is subgroup}\} \to \{F \le E \le K \mid E \text{ is intermidate field}\}$$
$$: H \mapsto K^H$$


is bijection, with the inverse $\Phi^{-1}(E) = Gal(K/E)$.

2). Given $H_1, H_2 \le G$,

$H_1 \le H_2$ if and only if $K^{H_2} \le K^{H_1}$.

3). Given $H \le G$,

$$[K : K^H] = |H| \quad \text{and} \quad [K^H : F] = |Gal(K/F) : H|$$



4). Given $H \le G$, K/K^H is always Galois, with $Gal(K/K^H) = H$.

Moreover, K^H/F is Galois if and only if $H \trianglelefteq G$ is normal.

If K^H/F is Galois, then $Gal(K^H/F) \cong Gal(K/F)/H$.

Further, there exists one-to-one correspondence between

$$\{\varphi : K^H \to \overline{K} \mid \varphi \text{ is embedding which fix } F\} \leftrightarrow \{\sigma H \mid \sigma \in G\},$$

5). Given $H_1, H_2 \le G$, then

$$K^{\langle H_1, H_2 \rangle} = K^{H_1} \cap K^{H_2} \quad \text{and} \quad K^{H_1 \cap H_2} = K^{H_1} K^{H_2}$$

