Artin - linearly independence of characteristic theorem.

Def. Let G be a group and L be a field.
A group-Homomorphism $\chi : G \to L^\times$ is called a character of G in L.
The characters $\chi_1, \cdots, \chi_n$ are said to be linearly independent over L if
$\sum_{i=1}^{n} a_i \cdot \chi_i \equiv 0$ for any $a_i \in L$ implies $a_1 = \cdots = a_n = 0$.

Thm. If $\chi_1, \cdots, \chi_n : G \to L^\times$ are distinct characters of G in L,
then they are linearly independent over L.

Proof. Suppose that $\chi_1, \cdots, \chi_n$ are linearly dependent.
Choose $I \subseteq [n]$ with minimum cardinality among all non-empty sets satisfying
$\sum_{k \in I} a_k \cdot \chi_k \equiv 0$ and $\forall k \in I,\ a_k \neq 0$. $\cdots$ (*), with the coefficients a_i.
Since the characters are dependent, there are subsets of $[n]$ satisfying such conditions, so we can take this I. Moreover, since each Charac-ters have only non-zero valued, so I must have at least two elements.
Let $i, j \in I$ with $i \neq j$. Since $\chi_i \neq \chi_j$, for some $g_0 \in G$, we have

$$\chi_i(g_0) \neq \chi_j(g_0)$$

Given $g \in G$, $0 = \sum_{k \in I} a_k \cdot \chi_k(g_0 g) = \sum_{k \in I} a_k \cdot \chi_k(g_0) \cdot \chi_k(g)$. And, multiplying $\chi_j(g_0)$ to (*), $0 = \sum_{k \in I} a_k \cdot \chi_j(g_0) \cdot \chi_k(g)$. Now, substracting the two equations,
then we have $\sum_{k \in I} a_k \cdot (\chi_k(g_0) - \chi_j(g_0)) \cdot \chi_k(g) = 0$.
But, the coefficient of $\chi_j(g)$ is zero and of $\chi_i(g)$ is not zero,
so it is also non-trivial representation with the non-empty index set $J \subsetneq I$,
it contradicts to (*). Consequently, the characters are independent. □

Note: $\mathcal{P} := \left\{ (a_1, \cdots, a_n) \in L^n \setminus \{(0, \cdots, 0)\} \,\middle|\, \sum_{k \in [n]} a_k \cdot \chi_k \equiv 0 \right\}$ is non-empty iff
$\{\chi_1, \cdots, \chi_n\}$ is linearly dependent.

View of Representation theory

Let $F^G := \{f \mid f: G \to F\}$ where G is a group and F is a field.

Then, F^G is a vector space under the operations

$$(f_1 + f_2)(g) := f_1(g) + f_2(g) \quad \text{and} \quad (af)(g) := a \cdot f(g).$$

Let $h \in G$ be given. Define

$$T_h : F^G \to F^G : f \mapsto T_h f \quad \text{where} \quad T_h f : G \to F : g \mapsto f(hg).$$

Normal basis theorem.

Def. Let E/F be a Galois extension field of degree of n.
For fixed $\gamma \in E$, a set

$$\eta = \{\sigma_1(\gamma), \cdots, \sigma_n(\gamma)\} \quad \text{where } Gal(E/F) = \{\sigma_1, \cdots, \sigma_n\}.$$

is called a normal basis for E/F if η is a basis.

Lemma. Let E/F be Galois extension with $Gal(E/F) = \{\sigma_1, \cdots, \sigma_n\}$.
For a basis $\beta = \{e_1, \cdots, e_n\}$ for E over F,

$$A = \{(\sigma_1(e_i), \cdots, \sigma_n(e_i)) \mid 1 \le i \le n\}$$

forms an E-basis for E^n.

Proof. Let $W = Span_L(A) \le L^n$. Suppose that $W \neq L^n$.
Then, there exists a nonzero linear functional $\varphi : L^n \to L$ which vanishes on W. (In other words, $W \neq L^n$ implies $W^\circ \neq (L^n)^\circ = \{0\}$.)