

In this section, we study with the following steps:

1. Generalized Eigenspace.
2. Cycle of Generalized Eigenvector.
3. Necessary Condition for a given Space to be represented as a Direct sum of Generalized Eigenspaces.
4. Finite-Dimensional Generalized Eigenspace has a basis that represents the linear map as a Jordan block.

0.1 Jordan Canonical Form

Let $T : V \rightarrow V$ be a linear operator, and $f(t) \in F[t]$ be a characteristic polynomial of T .

For a linear map $L : V \rightarrow V$, and L -invariant subspace $W \leq V$, denote $L|_W$ as a restriction:

$$L|_W : W \rightarrow W : x \mapsto L(x)$$

0.1.1 Generalized Eigenspaces

Definition 0.1.1. Suppose that $\lambda \in F$ be an eigenvalue of T . Define a set:

$$K_\lambda \stackrel{\text{def}}{=} \{x \in V \mid (T - \lambda I)^p(x) = 0 \text{ for some } p \in \mathbb{N}\} = \bigcup_{n=1}^{\infty} \ker(T - \lambda I)^n$$

It is called a *generalized eigenspace* corresponding to λ .

For non-zero $x \in K_\lambda$, the smallest positive integer $p \in \mathbb{Z}$ s.t. $(T - \lambda I)^p(x) = 0$ is called *period* of x (or *length*). (If $x = 0$, the period of x is defined to be zero.)

Lemma 0.1.2. Let $\lambda \in F$ be an eigenvalue of T . Then, the following properties are satisfied:

1. K_λ is a T -invariant subspace of V containing E_λ .
2. If $\mu \in F$ is an eigenvalue of T such that $\mu \neq \lambda$, then $K_\mu \cap K_\lambda = \{0\}$.

Proof. 1) Let $x, y \in K_\lambda$ and $\alpha \in F$ be given. Denote $p, q \in \mathbb{Z}$ as period of x, y , respectively. Then,

$$(T - \lambda I)^{p+q}(\alpha x + y) = \alpha(T - \lambda I)^{p+q}(x) + (T - \lambda I)^{p+q}(y) = 0.$$

Clearly, $0 \in K_\lambda$. Therefore, subspace criterion gives $K_\lambda \leq V$.

To show T -invariant, let $x \in K_\lambda$. Then, for period $p \in \mathbb{Z}$ of x ,

$$(T - \lambda I)^p(T(x)) = ((T - \lambda I)^p \circ T)(x) = (T \circ (T - \lambda I)^p)(x) = T(0) = 0.$$

Thus $T(x) \in K_\lambda$. Including E_λ is clear.

Show 2). Let $w \in K_\mu \cap K_\lambda$ be given. Let $p, q \in \mathbb{Z}$ be period of w for μ and λ , respectively. Since $\mu \neq \lambda$, $(t - \mu)^p, (t - \lambda)^q \in F[t]$ are disjoint. Now, there exist $g_1(t), g_2(t) \in F[t]$ s.t.

$$g_1(t)(t - \mu)^p + g_2(t)(t - \lambda)^q = 1$$

because $F[t]$ is P.I.D. Now,

$$(g_1(T)(T - \mu I)^p + g_2(T)(T - \lambda I)^q)(w) = 0 + 0 = w$$

Thus $K_\mu \cap K_\lambda = \{0\}$. □

0.1.4 The Existence of Jordan-Canonical Form

Now, we know that: if the characteristic polynomial of T splits over F , then the space V can be represented by direct sum of generalized eigenspaces. And, the cycle of eigenvector generates some subspace of generalized eigenspace, moreover, it is Jordan-basis for the restriction. If we can show that some cycles form a partition for basis of the eigenspace, finally, we can conclude that there exists a Jordan Canonical Form.

Lemma 0.1.9. Let $T: V \rightarrow V$ be a linear operator, and $\lambda \in F$ be an eigenvalue of T . Consider that the generalized eigenvectors $x_1, \dots, x_q \in K_\lambda$ with period $p_i \in \mathbb{Z}$, respectively. If elements in $\{(T - \lambda I)^{p_i-1}(x_i) \mid 1 \leq i \leq q\}$ are mutually different and the set is linearly independent, then

$$\mathcal{C} \stackrel{\text{def}}{=} \bigcup_{i=1}^q \mathcal{C}_{x_i}$$

is linearly independent. (Disjointness is already shown by the Lemma.)

Proof. Using mathematical induction for number of elements of $\mathcal{C}$.

If $\mathcal{C}$ has only one element, then trivial. Suppose that the statement holds for any $\mathcal{C}$ having fewer than n elements. Now, suppose that $\mathcal{C}$ has exactly n elements. Denote that

$$W \stackrel{\text{def}}{=} \langle \mathcal{C} \rangle, \quad U \stackrel{\text{def}}{=} (T - \lambda I)|_W$$

Since each $\mathcal{C}_{x_i}$ generates T -invariant subspace and $\langle A \cup B \rangle = \langle A \rangle + \langle B \rangle$ and sum of T -invariant is T -invariant, W is T -invariant, also $(T - \lambda I)$ -invariant. Thus, the restriction U is well-defined.

For each $1 \leq i \leq q$, $\mathcal{C}_{x_i} \setminus \{x_i\}$ is either emptyset or a cycle having end vector as $(T - \lambda I)(x_i)$, having same initial vector.

$$\text{im } U = U[\langle \mathcal{C} \rangle] = U \left[\left\langle \bigcup_{i=1}^q \mathcal{C}_{x_i} \right\rangle \right] = U \left[\sum_{i=1}^q \langle \mathcal{C}_{x_i} \rangle \right] = \sum_{i=1}^q U[\langle \mathcal{C}_{x_i} \rangle] = \sum_{i=1}^q \langle U[\mathcal{C}_{x_i}] \rangle = \sum_{i=1}^q \langle \mathcal{C}_{x_i} \setminus \{x_i\} \rangle = \left\langle \underbrace{\bigcup_{i=1}^q \mathcal{C}_{x_i} \setminus \{x_i\}}_{\mathcal{C}'} \right\rangle$$

Since the set $\mathcal{C}'$ is consisted by $n - q$ elements, and the initial vectors are linearly independent, therefore $\mathcal{C}'$ is linearly independent, by hypothesis of the induction. Thus it is a basis for $\text{im } U$. Moreover, $\ker U$ has more than q elements, because each initial vector of $\mathcal{C}_{x_i}$ is contained in $\ker U$. Consequently,

$$n \geq \dim W = \dim \text{im } U + \dim \ker U \geq (n - q) + q = n$$

This implies $\dim W = n$, thus $\mathcal{C}$ is basis of W . Thus, linearly independent. □

Theorem 0.1.10. Let $T: V \rightarrow V$ be a linear operator, and $\lambda \in F$ be an eigenvalue of T . If the generalized eigenspace K_λ has finite dimension, then it has a basis consisting of a disjoint union of cycles of generalized eigenvectors.

Proof. Using mathematical induction for dimension of K_λ . If $\dim K_\lambda = 1$, trivial.

Suppose that for $n \in \mathbb{N}$, the statement holds for any generalized eigenspace K such that $\dim K < n$.

Now, assume that $\dim K_\lambda = n$. Denote $U \stackrel{\text{def}}{=} (T - \lambda I)|_{K_\lambda}$. Note that: $\text{im } U < K_\lambda$ because the Dimension Theorem gives

$$\dim K_\lambda = \dim \text{im } U + \dim \ker U$$

But $\ker U = E_\lambda \leq K_\lambda$ is not a zero space, $\dim \text{im } U < \dim K_\lambda$.

Moreover, for any $x \in \text{im } U$, $x = (T - \lambda I)(y)$ for some $y \in K_\lambda$, by definition. Now

$$T(x) = T((T - \lambda I)(y)) = (T - \lambda I)(T(y)) \in \text{im } U$$

That is, $\text{im } U$ is T -invariant, which means $\text{im } U$ is generalized eigenspace of $T|_{\text{im } U}$ corresponding to λ .

By the assumption of induction, there exist disjoint cycles C_i , $1 \leq i \leq q$ such that

$$\mathcal{C} = \bigcup_{i=1}^q C_i$$

is a basis of $\text{im } U$. For each $1 \leq i \leq q$, the end vector of C_i is contained in $\text{im } U$, thus we can extend the cycle C_i to

$$\bar{C}_i = C_i \cup \{v_i\}$$

And, denote w_i as an initial vector of a cycle C_i (It is also initial vector of $\bar{C}_i$.)

Since the initial vectors are contained in E_λ and linearly independent, we can obtain extended basis for E_λ ,

$$\{w_1, \dots, w_q, u_1, \dots, u_s\}$$

Finally, initial vectors in the following cycles are linearly independent:

$$\bar{C}_1, \dots, \bar{C}_q, \{u_1\}, \dots, \{u_s\}$$

by the Lemma, the union of these sets is linearly independent. The Dimension Theorem gives the result. $\square$