

## Commutator

Def. Let  $G$  be a group. Given  $x, y \in G$ , define

$$[x, y] \stackrel{\text{def}}{=} x^{-1}y^{-1}xy$$

is called the commutator of  $x$  and  $y$ . And, given two subset  $A$  and  $B$ ,

$$[A, B] \stackrel{\text{def}}{=} \langle [a, b] \mid a \in A, b \in B \rangle$$

Particularly,  $G' \stackrel{\text{def}}{=} [G, G]$  is called the derived set.

Prop.  $G'$  char  $G$ , (so  $G' \trianglelefteq G$ ).

Proof. Given  $\sigma \in \text{Aut}(G)$  and  $[x, y]$  for arbitrary  $x, y \in G$ ,

$$\sigma([x, y]) = \sigma(x^{-1}y^{-1}xy) = \sigma(x)^{-1} \cdot \sigma(y)^{-1} \cdot \sigma(x) \cdot \sigma(y) = [\sigma(x), \sigma(y)]$$

$$\text{So, } \sigma[[G, G]] = [G, G].$$

Prop. Given  $H \leq G$ ,  $H \trianglelefteq G$  if and only if  $[H, G] \leq H$

Proof.  $H \trianglelefteq G \Leftrightarrow$  for all  $g \in G$  and  $h \in H$ ,  $ghg^{-1} \in H$

$$[H, G] \leq H \Leftrightarrow \text{for all } g \in G \text{ and } h \in H, [h, g] = h^{-1}g^{-1}hg \in H.$$

So proved.

Prop. Given  $N \trianglelefteq G$ ,  $G/N$  is abelian if and only if  $G' \leq N$ .

Proof.  $G/N$  is abelian iff for any  $xN, yN \in G/N$ ,  $xyN = yxN$

iff for any  $x, y \in G$ ,  $x^{-1}y^{-1}xy \in N$

iff  $G' \leq N$ .

Note:  $G'$  is the smallest normal subgroup of  $G$  s.t. the factor is abelian.

Note: If  $G$  is abelian, then  $G' = 1$ .

Note:  $[x, y] \cdot [y, z] = x^{-1}y^{-1}xy \cdot y^{-1}z^{-1}yz = x^{-1}y^{-1}xz^{-1}yz$

$$= \begin{cases} [x, y]^t = y^{-1}x^{-1}yx = [y, x] \\ [xy, y^{-1}] = x^{-1}yx^{-1}y \\ [x^{-1}, y] = xy^{-1}x^{-1}y. \\ [x, y][y, x] = x^{-1}y^{-1}xy \cdot y^{-1}x^{-1}yx = \text{id}. \end{cases}$$

Commutator of Typical Subgroups.  
 $D_{2n}$ , the dihedral of order  $2n$ .

$S_n$ , the symmetric group on  $n$  letters

$$[S_3, S_3] = \langle [a, b] \mid a, b \in S_3 \rangle$$

$$[(1\ 2), (1\ 3)] = (1\ 2)(1\ 3)(1\ 2)(1\ 3) = (1\ 3\ 2)(1\ 3\ 2) = (1\ 2\ 3)$$

$$[(1\ 2), (2\ 3)] = (1\ 2)(2\ 3)(1\ 2)(2\ 3) = (1\ 2\ 3\ 1)(1\ 2\ 3\ 1) = (1\ 3\ 2)$$

$$[(1\ 3), (2\ 3)] = (1\ 3)(2\ 3)(1\ 3)(2\ 3) = (3\ 2\ 1)(3\ 2\ 1) = (1\ 2\ 3)$$

$$[(1\ 2), (1\ 2\ 3)] = (1\ 2)(1\ 2\ 3)^{-1}(1\ 2)(1\ 2\ 3) = (1\ 2)(3\ 2\ 1)(1\ 2)(1\ 2\ 3) = (1\ 3\ 2)$$

$$[(1\ 3), (1\ 2\ 3)] = (1\ 3)(1\ 2\ 3)^{-1}(1\ 3)(1\ 2\ 3) = (1\ 3)(3\ 2\ 1)(1\ 3)(1\ 2\ 3) = (1\ 3\ 2)$$

$$[(2\ 3), (1\ 2\ 3)] = (2\ 3)(1\ 2\ 3)^{-1}(2\ 3)(1\ 2\ 3) = (2\ 3)(3\ 2\ 1)(2\ 3)(1\ 2\ 3) = (1\ 3\ 2)$$

$$[(1\ 2\ 3), (1\ 3\ 2)] = \text{id}.$$

$$\text{Therefore, } [S_3, S_3] = \langle (1\ 2\ 3), (1\ 3\ 2) \rangle = \{\text{id}, (1\ 2\ 3), (1\ 3\ 2)\} = A_3.$$

Alternatively, since  $A_3$  is normal subgroup of  $S_3$  and  $S_3/A_3 \cong C_2$ , so  $[S_3, S_3] \subseteq A_3$ . Conversely, since  $(1\ 3\ 2) \in A_3$  and  $(1\ 3\ 2) = [(1\ 2), (2\ 3)] \in [S_3, S_3]$ . Therefore  $[S_3, S_3] \supseteq A_3 = \langle (1\ 3\ 2) \rangle$ .

$\Gamma A_n$ , the alternating group.

$\Gamma_8$ . The Quaternion.