

Lemma.

Let K/F be a field extension, and let $\alpha \in K$ be algebraic over F . Then, for any $\sigma \in \text{Aut}(K/F)$, $\sigma\alpha$ is a root of the minimal polynomial for α over F .

Proof. Let $f(x) = x^n + a_{n-1}x^{n-1} + \dots + a_1x + a_0 \in F[x]$ be the minimal polynomial for α . Then, $f(\alpha) = 0$, so $\sigma(f(\alpha)) = 0$. (The field homomorphism preserves zero).

Now, $0 = \sigma(f(\alpha)) = \sigma(\alpha^n + a_{n-1}\alpha^{n-1} + \dots + a_1\alpha + a_0)$

$$\begin{aligned} \sigma \text{ fixes } F &\hookrightarrow \begin{aligned} &= (\sigma\alpha)^n + \sigma(a_{n-1}) \cdot (\sigma\alpha)^{n-1} + \dots + \sigma(a_1) \cdot (\sigma\alpha) + \sigma(a_0) \\ &= (\sigma\alpha)^n + a_{n-1}(\sigma\alpha)^{n-1} + \dots + a_1(\sigma\alpha) + a_0 \\ &= f(\sigma\alpha). \end{aligned} \end{aligned}$$

Prop. Let $H \leq \text{Aut}(K)$ be a subgroup of the group of all automorphisms of K .

Then, the collection F of elements of K fixed by all the elements of H is a subfield.

Proof. First, given $\sigma \in \text{Aut}(K)$, the set $F_\sigma = \{\alpha \in K \mid \sigma(\alpha) = \alpha\}$ is subfield, because:

Let $a, b \in F_\sigma$ be given. Then, $\sigma(a-b) = \sigma(a) - \sigma(b) = a-b$, and if $b \neq 0$, then $\sigma(a \cdot b^{-1}) = \sigma(a) \cdot \sigma(b)^{-1} = a \cdot b^{-1}$, so the claim is proved.

And now, since the arbitrary intersection of fields is field, so proved. ~~QED~~
Such a subfield is called the fixed field of $S \subseteq \text{Aut}(K)$.

Prop. Let K be a field. Then,

If $F_1 \leq F_2 \leq K$, then $\text{Aut}(K/F_2) \leq \text{Aut}(K/F_1)$.

If $H_1 \leq H_2 \leq \text{Aut}(K)$, then $F_{H_2} \leq F_{H_1}$.

In other say, The inclusion relation of automorphisms and fixed field is reverse.

[Prop. Let E be the splitting field over F of the polynomial $f(x) \in F[x]$. Then, $|\text{Aut}(E/F)| \leq [E:F]$, and the equality holds if $f(x)$ is separable over F .

Proof. Using induction for $[E:F] = n$. If $n=1$, then $E=F$, so $\text{Aut}(E/F)$ is trivial. If $n>1$, then $f(x)$ has at least one irreducible factor $p(x)$ with $\deg p > 1$.

Let $\alpha \in E$ be a fixed root of $p(x)$. By the Conjugate isomorphism Theorem, there are at most $\deg p$ extensions of identity $\varphi: F \rightarrow F$ to $\overline{\varphi}: F(\alpha) \rightarrow E$, and the number of extensions is exactly $\deg p$ if $p(x)$ is separable.

By the tower Lemma, $[E:F(\alpha)] \cdot [F(\alpha):F] = [E:F]$ and since $[F(\alpha):F] > 1$, so $[E:F(\alpha)] < n$, we can use the assumption of induction (clearly, E is splitting field of f over $F(\alpha)$) we can deduce that $|\text{Aut}(E/F(\alpha))| \leq [E:F(\alpha)]$. Finally,

$$|\text{Aut}(E/F)| = \underset{\uparrow}{|\text{Aut}(E/F(\alpha))| \cdot |\text{Aut}(F(\alpha)/F)|} \leq [E:F(\alpha)] \cdot \underset{\uparrow}{[F(\alpha):F]} = [E:F]$$

the possibilities of composition

$[F(\alpha):F] = \deg p$.

Def. A finite extension K/F is called a Galois Extension if:

$$|Aut(K/F)| = [K:F].$$

In such a case, denote $G_K(K/F) = Aut(K/F)$.

Def. A character χ of a group G with values in a field L is a group Homomorphism.

$$\chi: G \rightarrow L^\times$$

The characters $\chi_1, \dots, \chi_n$ of G are said to be linearly independent over L if:

$a_1\chi_1 + \dots + a_n\chi_n = 0$ for some $a_i \in L$, then $a_1 = \dots = a_n = 0$.

