Topological Group.

Def. Let G be a Group under the binary operation $\cdot : G \times G \to G$.
A Topological Space $(G, \mathcal{U})$ satisfying T_1 condition, and

$$G \times G \to G : (x,y) \mapsto x \cdot y \quad \text{and} \quad G \to G : x \mapsto x^{-1} \quad \text{are continuous}$$

is called the Topological Group.

Lemma. Let G be a group and T_1-space. Then

G is Topological Group if and only if $G \times G \to G : (x,y) \mapsto xy^{-1}$ is continuous.

Proof. Suppose that G is Topological Group. Then

$\varphi(x,y) = xy$ and $\psi(y) = y^{-1}$ are continuous, therefore

$(x,y) \mapsto xy^{-1} = \varphi(x, \psi(y))$ is continuous, being composition of continuous maps

Conversely, suppose that $\mu : G \times G \to G : (x,y) \mapsto xy^{-1}$ is continuous.
Then, for fixed $x = 1$, $\mu(1,y) = y^{-1}$ is continuous, so $x \mapsto x^{-1}$ is continuous.
Now, $y \mapsto y^{-1}$ is continuous, so

$(x,y) \mapsto x \cdot (y^{-1})^{-1} = xy$ is continuous, being composition of continuous. ∎

$\Gamma GL_n(\mathbb{R})$

Consider $GL_n(\mathbb{R})$ as the subset of $\mathbb{R}^{n^2}$ equipped Euclidean Topology.

$\Gamma GL_n(\mathbb{R})$.

Consider $GL_n(\mathbb{R})$ as the subset of $\mathbb{R}^{n^2}$, with Euclidean space.

Define a metric for $M_{n\times n}(\mathbb{R})$ as:

$$d: M_{n\times n}(\mathbb{R}) \times M_{n\times n}(\mathbb{R}) \to \mathbb{R}: (A, B) \mapsto \sqrt{\sum_{i,j} (A_{ij} - B_{ij})^2}$$

Since the Matrix multiplication

$$\mu: M_{n\times n}(\mathbb{R}) \times M_{n\times n}(\mathbb{R}) \to M_{n\times n}(\mathbb{R}): (A, B) \mapsto \left(\sum_{k=1}^{n} A_{ik} \cdot B_{kj}\right)_{ij}$$

is continuous, because it is entry-wise continuous.

Since $GL_n(\mathbb{R})$ is closed under the multiplication, it has also continuous operation.

And, $GL_n(\mathbb{R}) \to GL_n(\mathbb{R}): A \mapsto A^{-1}$

is also continuous, because $A^{-1} = (\det A)^{-1} \cdot \operatorname{adj} A$

where $(\operatorname{adj} A)_{ij} = (-1)^{i+j} \cdot \det A(j|i)$.

is also multivariables polynomial, entrywise.

$SL_n(\mathbb{R})$

Consider $O_n(\mathbb{R}) = \{A \in M_{n\times n}(\mathbb{R}) \mid A^t = A^{-1}\}$, $SO_n(\mathbb{R}) = \{A \in O_n(\mathbb{R}) \mid \det A = 1\}$.

Then, O_n and SO_n are compact, and SO_n is connected.

proof. $A \in O_n(\mathbb{R}) \Leftrightarrow A^tA = I \Leftrightarrow \sum_{k=1}^{n} A_{ik} \cdot A_{jk} = \delta_{ij}$.

Define a map $f_{ij}: M_{n\times n}(\mathbb{R}) \to \mathbb{R} : A \mapsto \sum_{k=1}^{n} A_{ik} \cdot A_{jk}$.

Then, it is continuous therefore

$$O_n(\mathbb{R}) = \bigcap_{\substack{1\le i,j\le n \\ i\neq j}} f_{ij}^{-1}[\{0\}] \cap \bigcap_{1\le i=j\le n} f_{ij}^{-1}[\{1\}] \quad \text{is finite intersection of closed sets,}$$

thus closed. Moreover, $\dim O_n(\mathbb{R}) \le n$, so it is bounded. Therefore compact.

And, $\det : O_n(\mathbb{R}) \to \mathbb{R} : A \mapsto \det A$ is continuous, so,

$SO_n(\mathbb{R}) = \det^{-1}[O_n(\mathbb{R})]$ also closed, and bounded ~~or~~ subset of bounded set.

$\Gamma O_n(\mathbb{R})$

$\Gamma Son(IB)$

$\Gamma U_n(\mathbb{C})$