

Prop. Let $H \trianglelefteq G$. Then G acts by conjugation on H as automorphism of H . That is,
 $G \times H \rightarrow H : (g, h) \mapsto ghg^{-1}$.

The kernel of this action is $C_G(H)$. Particular, $G/C_G(H) \cong S \leq \text{Aut}(H)$.

Proof. By the cancellation, $h \mapsto ghg^{-1}$ is injective, and the surjective is given by

$$k \in H \Rightarrow g^{-1}k \in H \Rightarrow g(g^{-1}k)g^{-1} = k$$

Moreover, $ghkg^{-1} = ghg^{-1}gkg^{-1}$, thus Homomorphism. (well-defined is given by normality)

Let $\chi: G \rightarrow S_H : g \mapsto \chi_g$ where $\chi_g: H \rightarrow H: h \mapsto ghg^{-1}$ be the permutation represent of the action.

Now, $\text{Ker } \chi = \{g \in G \mid \chi_g = \text{id}\} = \{g \in G \mid ghg^{-1} = h \text{ for all } h \in H\} = C_G(H)$.

Now, $G/\text{Ker } \chi \cong S$ for some subgroup $S \leq \text{Aut}(H)$.

Corollary, $K \leq G$ and $g \in G$, then $K \cong gKg^{-1}$.

$K \rightarrow gKg^{-1} : k \mapsto gkg^{-1}$ is one-to-one, onto, Homomorphism.

Corollary. $H \leq G$, then $N_G(H)/C_G(H) \cong S \leq \text{Aut}(H)$.

And, $G/Z(G) \cong S \leq \text{Aut}(H)$.

Lemma. Let $P \in \text{Syl}_p(G)$. If Q is p -subgroup of G , then $Q \cap N_G(P) = Q \cap P$.

Proof. Since $P \leq N_G(P)$, $Q \cap P \leq Q \cap N_G(P)$ is trivial.

Since $Q \cap N_G(P) \leq N_G(P)$, denote $H = Q \cap N_G(P)$,

$$PH = \bigcup_{h \in H} Ph = \bigcup_{h \in H} hP = HP$$

Thus $PH \leq G$ and $|PH| = \frac{|P| \cdot |H|}{|P \cap H|}$. By the Lagrange's Thm, $H \leq Q$ means

order of H is power of p . Therefore, $P \leq PH$ and $|PH|$ is power of p , thus $P \cap H = P$, or $H \leq P$. Now, $H = Q \cap N_G(P) \leq Q$, $H \leq Q \cap P$. $\square$

Lemma. If G is finite abelian group and p is a prime number dividing $|G|$, then G has an element of order of p .

Proof. Using induction for $|G|$. If $|G|=1$, trivial.

Now, suppose that the statement hold for all groups whose order is smaller than $|G|$.

Since $|G|>1$, take $x \in G$ with $x \neq 1$. That is, $|x|>1$.

If $|G|=p$, then $|x|=p$, by the Lagrange's Theorem ($|x| \mid p$).

Now, suppose $|G| \neq p$ ($|G|>p$).

If p divides $|x|$, $|x|=p^n$ for some $n \in \mathbb{N}$, so $|x^p|=p$. So clear.

If p does not divide $|x|$, let $N = \langle x \rangle$. (Clearly, $p \nmid |N| = |x|$).

Since G is abelian, N is Normal, and $x \neq 1$, so it is non-trivial.

By the Lagrange Theorem, $|G| = |G/N| \cdot |N|$. But, p does not divide $|N|$,

p divides $|G/N|$. Now, using the hypothesis of the induction

G/N has an element of order p , denote this $\bar{y} = yN$. Since $\bar{y} \neq \bar{1}$, $y \notin N$.

But, $\bar{y}^p = \bar{1}$, so $y^p \in N$. This implies $\langle y^p \rangle \neq \langle y \rangle$. Since $|y^p|$ divides $|y|=p$, the only possibilities are $|y^p|=1$, this y is we desired.

The Sylow Theorem. Let G be a group of order $p^n \cdot m$, p is prime not dividing m .
First, $Syl_p(G) \neq \emptyset$.

Second. If $P \in Syl_p(G)$ and $Q \leq G$ is a p -subgroup, then $Q \leq P g^{-1}$ for some $g \in G$.

Third. $n_p \equiv 1 \pmod{p}$, $n_p = [G : N_G(P)]$ for any $P \in Syl_p(G)$, and $n_p | m$.

Proof. To show the first, we will use the induction for order of G .

If $|G|=1$, then trivial. Now, suppose that $Syl_p(H) \neq \emptyset$, for all group whose order is less than $|G|$.

In the case of p divides the order $Z(G)$, then by the Cauchy's Theorem for abelian group, there is an order p element γ in $Z(G)$. Put $N = \langle \gamma \rangle$. Then, N is Normal of G .

Now, G/N is group of order of $p^{n-1} \cdot m$, and by the hypothesis of induction,

G/N has a subgroup of order of p^{n-1} , denote this $\overline{P} = P/N < G/N$.

By the 4th Isomorphism Theorem, there is $P < G$ s.t. $|P/N| = p^{n-1}$, so $|P| = p^n$.

In the case of p does not divide the order of $Z(G)$,

let $g_1, \dots, g_r$ be the distinct non-central representatives of conjugacy classes of G .

Then, by the class equation,

$$p^n \cdot m = |G| = |Z(G)| + \sum_{i=1}^r [G : C_G(g_i)]$$

Then, for some $1 \leq i \leq r$, p does not divide $[G : C_G(g_i)]$, because $p \nmid |Z(G)|$.

For this i , let $H = C_G(g_i)$. Since $g_i \notin Z(G)$, $|H| < |G|$.

Now, since $[G : H]$ has no factor of p , so $|H| = p^\alpha \cdot k$, $p \nmid k$, $k < m$.

By the hypothesis of the induction H has a subgroup of order of p^α ,

so this is a subgroup we desired.

Corollary. Let $P \in Syl_p(G)$.

$n_p = 1 \iff P \trianglelefteq G \iff P \text{ char } G$

[Group of order of pg , $p < g$. (p, g are prime)

Let $P \in \text{Syl}_p(G)$ and $Q \in \text{Syl}_g(G)$.

Since n_g divides P , $n_g = 1$ or P , but $n_g \equiv 1 \pmod{g}$, thus $n_g = 1$. $\therefore Q \trianglelefteq G$.

1. Case of $P \nmid g-1$.

By the Sylow Thm, $n_p = 1$ or g , but if $n_p = g$, then $g \equiv 1 \pmod{p}$, $p \mid g-1$, contradiction.

Therefore, $n_p = 1$, $P \trianglelefteq G$. Now, G must be a cyclic, because:

Since $|P|$ and $|Q|$ are primes, both are cyclic. Let $P = \langle x \rangle$ and $Q = \langle y \rangle$.

Since $P \trianglelefteq G$, $N_G(P) = G$. Therefore,

$$G/C_G(P) = N_G(P)/C_G(P) \cong S \text{ for some } S \leq \text{Aut}(\mathbb{Z}_p)$$

But, $|\text{Aut}(\mathbb{Z}_p)| = p-1$, thus S is trivial. (because $|S|$ divides $|G| = pg$).

Now, $G = C_G(P)$, and $x \in P$, x and y are commute. This implies $|xy| = pg$. $G = \langle pg \rangle$.

2. Case of $P \mid g-1$

Let Q be a Sylow g -subgroup of S_g , the symmetric group of letters.

Since g is prime, $|N_{S_g}(Q)| = g \cdot (g-1)$, because

1). Since $|S_g| = g!$, $|Q| = g$.

2). S_g has $(g-1)!$ g -cycles.

3). Since order of gQg^{-1} is g , these are all cyclic.

4). $gQg^{-1} \cap Q = \{1\}$ for all $g \in G$, so

$$n_g = \# \{gQg^{-1} \mid g \in G\} = (g-1)! / (g-1) = (g-2)!$$

$$\text{Therefore } n_g = G_Q = [S_g : N_G(Q)] = (g-2)!, \text{ or } N_{S_g}(Q) = g(g-1). \quad \overset{p}{\uparrow}$$

Now, since $p \mid g-1$, p divides order of $N_{S_g}(Q)$, so it has a subgroup of order p .

And, since $p \leq N_{S_g}(Q)$, pQ is subgroup of order pg .

And, $C_{S_g}(Q) = Q$, pQ is non-abelian group.

$$|G : N_{S_g}(Q)| = G_Q$$

Group of order 48 is not simple.

$48 = 2^4 \cdot 3$. Therefore,

$$n_3 = 1 \text{ or } 4 \text{ or } 16 \quad \text{and} \quad n_2 = 1 \text{ or } 3.$$

If $n_2 = 1$, then $\text{PG Syl}_2(G)$ is the Normal, so there is nothing to prove.

Suppose $n_2 \neq 1$, or $n_2 = 3$. That is, there are three Sylow 2-subgroup,

$$P_1, P_2, P_3 \in G. \quad (\#P_i = 16).$$

For each $1 \leq i < j \leq 3$,

$$|P_i P_j| = \frac{|P_i| \cdot |P_j|}{|P_i \cap P_j|} = \frac{2^8}{|P_i \cap P_j|} \leq 48 \quad \begin{array}{l} \text{if } i, j, P_i = P_j \\ \downarrow \end{array}$$

So, $\frac{2^4}{3} \leq |P_i \cap P_j|$, $|P_i \cap P_j|$ cannot be 1, 2, 4, 16 therefore

$$|P_i \cap P_j| = 8.$$

Take $Q = P_1 \cap P_2$, as subgroup of order 8. Since $Q \subseteq P_1, P_2$ and index 2, $Q \trianglelefteq P_1$ and $Q \trianglelefteq P_2$. That is, for any $g \in P_1 \cup P_2$,

$$gQg^{-1} = Q.$$

therefore, $P_1 \cup P_2 \subseteq N_G(Q)$. Since $|P_1 \cap P_2| = 8$,

$$|P_1 \cup P_2| = |P_1| + |P_2| - |P_1 \cap P_2| = 24$$

Now, if $|N_G(Q)| = 24$, then it has index 2 to G ,

$$N_G(Q) \trianglelefteq G$$

And, if $|N_G(Q)| = 48$, then $N_G(Q) = G$, so $Q \trianglelefteq G$.