

Given a map  $f: \mathbb{R} \rightarrow \mathbb{R}$ , a map

$$g: x \mapsto \int_J k(x, t) f(t) dt$$

is called the 'integral transform'.

In this page, we cover typical int. forms.

Laplace transform.

Given a map  $f: \mathbb{R} \rightarrow \mathbb{C}$ , a Laplace transform of  $f$  is given by:

$$\mathcal{L}f: \mathbb{C} \rightarrow \mathbb{C}; x \mapsto \int_0^{\infty} e^{-xt} \cdot f(t) dt.$$

Prop. If  $f$  is differentiable and  $e^{-xR} \cdot f(R) \rightarrow 0$  as  $R \rightarrow \infty$ ,

then  $(\mathcal{L}f)(x) = \int_0^{\infty} e^{-xt} \cdot f(t) dt = \lim_{R \rightarrow \infty} \left[ e^{-xt} f(t) \Big|_0^R \right] + x \cdot \int_0^{\infty} e^{-xt} \cdot f(t) dt$

$$= x \cdot (\mathcal{L}f)(x) - f(0)$$

Fourier transform.

# Convolution

Given two maps  $f, g: \mathbb{R} \rightarrow \mathbb{C}$ , define

$$(f * g)(x) = \int_{-\infty}^{\infty} g(x-t) f(t) dt \quad x-t \in [-r, r] \Leftrightarrow -r+x$$

Prop. Let  $k_r(x) = \frac{1}{2r} \cdot \chi_{[-r, r]}(x)$ . Then,

$$(f * k_r)(x) = \int_{-\infty}^{\infty} k_r(x-t) \cdot f(t) dt = \frac{1}{2r} \cdot \int_{x-r}^{x+r} f(t) dt \quad (1)$$

Moreover,  $\int_{-\infty}^{\infty} |f| < \infty \Rightarrow f * k_r$  is uniformly continuous. (2)

and  $f$  is uniformly continuous  $\Rightarrow f * k_r \rightarrow f$  as  $r \rightarrow 0$  uniformly. (3)

Proof. (1) can be shown by:

$$g(x-t) \cdot f(t) = 0 \Leftrightarrow x-t \notin [-r, r] \Leftrightarrow x-r \leq t \leq x+r.$$

To show (2), let  $r > 0$  be fixed. Given  $\varepsilon > 0$ , take  $\delta > 0$  such that

$$0 < t-s < \delta \Rightarrow \int_s^t |f| < r\varepsilon.$$

Now, put  $\delta = \min\{2r, \delta_0\}$ . Then,  $|x-y| < \delta$  implies

$$\begin{aligned} |(f * k_r)(x) - (f * k_r)(y)| &= \frac{1}{2r} \cdot \left| \int_{x-r}^{x+r} f(t) dt - \int_{y-r}^{y+r} f(t) dt \right| \quad |x-y| < \delta < 2r. \\ &= \frac{1}{2r} \cdot \left| \int_{x-r}^{y-r} f(t) dt - \int_{x+r}^{y+r} f(t) dt \right| \\ &< \frac{1}{2r} (r\varepsilon + r\varepsilon) = \varepsilon. \end{aligned}$$

To show (3), let  $\varepsilon > 0$  be given. Then, there exists  $\delta > 0$  such that

$$|x-y| < \delta \Rightarrow |f(x) - f(y)| < \varepsilon.$$

Now, for  $r < \delta$ ,

$$\begin{aligned} |f(x) - (f * k_r)(x)| &= \left| \frac{1}{2r} \cdot \int_{x-r}^{x+r} f(t) dt - \frac{1}{2r} \cdot \int_{x-r}^{x+r} f(x) dt \right| \\ &\leq \frac{1}{2r} \cdot \int_{x-r}^{x+r} |f(x) - f(t)| dt < \varepsilon. \end{aligned}$$

Therefore,  $f * k_r \rightarrow f$  uniformly.  $\square$