

Thm. If  $f$  is a continuous complex-valued function on  $[a, b]$ , then there exists a sequence of polynomials  $P_n$  such that

$$P_n \rightarrow f \text{ uniformly on } [a, b].$$

Proof. WLOG, we assume  $[a, b] = [0, 1]$ , and  $f(0) = f(1) = 0$ .

(This is possible:  $g(x) = f(x) - f(0) - x(f(1) - f(0))$ )

And,  $f(x) = 0$  if  $x \notin [0, 1]$ .

Put  $Q_n(x) = C_n \cdot (1-x^2)^n \quad (n=1, 2, 3, \dots)$

where  $C_n$  is chosen such that  $\int_{-1}^1 Q_n(x) dx = 1$ .

To magnitade  $C_n$ ,

$$\begin{aligned} \int_{-1}^1 (1-x^2)^n dx &= 2 \cdot \int_0^1 (1-x^2)^n dx \geq 2 \cdot \int_0^{\frac{1}{\sqrt{n}}} (1-x^2)^n dx \\ &\geq 2 \cdot \int_0^{\frac{1}{\sqrt{n}}} (1-nx^2) dx = 2nx \cdot \left[ \frac{(1-x^2)^{n+1}}{-(n+1)} + 1 \right] \geq 0 \\ &= 2 \cdot \left( \frac{1}{\sqrt{n}} - \frac{n}{3} \cdot \frac{1}{(\sqrt{n})^3} \right) = \frac{4}{3} \frac{1}{\sqrt{n}} > \frac{1}{\sqrt{n}} \end{aligned}$$

so,  $C_n < \sqrt{n}$ . Now, for any  $\delta > 0$ ,

$$Q_n(x) \leq \sqrt{n} \cdot (1-\delta^2)^n \quad (\delta \leq |x| \leq 1), \text{ so that } Q_n \rightarrow 0 \text{ uniformly in } \delta \leq |x| \leq 1$$

Now, set  $P_n(x) = \int_{-1}^1 f(x+t) \cdot Q_n(t) dt \quad (0 \leq x \leq 1)$

Since  $f(s) = 0$  if  $s \notin [0, 1]$ , (Perised in the next page.)

$$\begin{aligned} &= \int_{-x}^{1-x} f(x+t) Q_n(t) dt \\ &= \int_0^1 f(s) Q_n(s-x) ds \quad (x+t=s \Rightarrow dt=ds) \end{aligned}$$

Given  $\epsilon > 0$ , choose  $\delta > 0$  s.t.  $|x-y| < \delta \Rightarrow |f(x) - f(y)| < \frac{\epsilon}{2}$ , Since  $f$  is continuous on compact

$$\begin{aligned} |P_n(x) - f(x)| &= \left| \int_0^1 [f(x+t) - f(x)] \cdot Q_n(t) dt \right| \leq \int_0^1 |f(x+t) - f(x)| \cdot Q_n(t) dt \\ &\leq 2M \cdot \left( \int_{-1}^{-\delta} Q_n(t) dt + \int_{\delta}^1 Q_n(t) dt \right) + \frac{\epsilon}{2} \cdot \int_{-\delta}^{\delta} Q_n(t) dt \quad (M = \sup |f|) \\ &\leq 4M \cdot \sqrt{n} (1-\delta^2)^n + \frac{\epsilon}{2} \rightarrow 0 \end{aligned}$$

