Given a map $f: \mathbb{R} \to \mathbb{R}$, a map

$$g: x \mapsto \int_J K(x,t) f(t)\, dt$$

is called the integral transform.

In this page, we cover typical int. forms.

[Laplace transform.
Given a map $f: \mathbb{R} \to \mathbb{C}$, a Laplace transform of f is given by:

$$\mathcal{L}f: \mathbb{C} \to \mathbb{C}; x \mapsto \int_0^\infty e^{-xt} \cdot f(t)\, dt.$$

[Prop. If f is differentiable and $e^{-x \cdot B} \cdot f(B) \to 0$ as $B \to \infty$,
then $(\mathcal{L}f')(x) = \int_0^\infty e^{-xt} \cdot f'(t)\, dt = \lim_{B \to \infty} \left[e^{-xt} f(t) \Big|_0^B \right] + x \cdot \int_0^\infty e^{-xt} \cdot f(t)\, dt$

$= x \cdot (\mathcal{L}f)(x) - f(0)$

Fourier transform.

Convolution

Given two maps $f, g : \mathbb{R} \to \mathbb{C}$, define

$$(f*g)(x) = \int_{-\infty}^{\infty} g(x-t) f(t)\, dt \qquad x-t \in [-r,r] \iff -r+x$$

Prop. Let $k_r(x) = \frac{1}{2r} \cdot \chi_{[-r,r]}(x)$. Then,

$$(f * k_r)(x) = \int_{-\infty}^{\infty} k_r(x-t) \cdot f(t)\, dt = \frac{1}{2r} \cdot \int_{x-r}^{x+r} f(t)\, dt \qquad (1)$$

Moreover, $\int_{-\infty}^{\infty} |f| < \infty \Rightarrow f * k_r$ is uniformly continuous. (2)

and f is uniformly continuous $\Rightarrow f * k_r \to f$ as $r \to 0$ uniformly. (3)

Proof. (1) Can be shown by:

$$g(x-t) \cdot f(t) = 0 \iff x-t \in [-r,r] \iff x-r \le t \le x+r.$$

To show (2), let $r > 0$ be fixed. Given $\varepsilon > 0$, take $\delta_0 > 0$ such that

$$0 < t-s < \delta_0 \Rightarrow \int_s^t |f| < r \cdot \varepsilon.$$

Now, put $\delta = \min\{2r, \delta_0\}$. Then, $|x-y| < \delta$ implies

$$\begin{aligned} |(f*k_r)(x) - (f*k_r)(y)| &= \frac{1}{2r} \cdot \left| \int_{x-r}^{x+r} f(t)\,dt - \int_{y-r}^{y+r} f(t)\,dt \right| \qquad |x-y| < \delta < 2r. \\ &= \frac{1}{2r} \cdot \left| \int_{x-r}^{y-r} f(t)\,dt - \int_{x+r}^{y+r} f(t)\,dt \right| \\ &< \frac{1}{2r} \cdot (r\varepsilon + r\varepsilon) = \varepsilon. \end{aligned}$$

To show (3), let $\varepsilon > 0$ be given. Then, there exists $\delta > 0$ such that

$$|x-y| < \delta \Rightarrow |f(x) - f(y)| < \varepsilon.$$

Now, for $r < \delta$,

$$\begin{aligned} |f(x) - (f*k_r)(x)| &= \left| \frac{1}{2r} \cdot \int_{x-r}^{x+r} f(x)\,dt - \frac{1}{2r} \cdot \int_{x-r}^{x+r} f(t)\,dt \right| \\ &\le \frac{1}{2r} \cdot \int_{x-r}^{x+r} |f(x) - f(t)|\,dt < \varepsilon. \end{aligned}$$

Therefore, $f * k_r \to f$ uniformly. ∎