



$T: V \rightarrow V$  is diagonalizable  $\Leftrightarrow m_T$  splits and separable

" triangulizable  $\Leftrightarrow m_T$  is separable

$E/F$  is separable  $\Leftrightarrow \# \text{Emb}_F(E, \bar{F}) = [E:F]$

$E/F$  is splitting  $\Leftrightarrow \text{Emb}_F(E, \bar{F}) = \text{Aut}(E/F)$

relation  $\{ \}$

$$\alpha, \beta \in \mathbb{R}^3, \quad r \in \mathbb{R}^3,$$



$$1. |\alpha \times \beta| = |\alpha| \cdot |\beta| \cdot \sin \theta.$$

$$2. \alpha \cdot \beta = |\alpha| \cdot |\beta| \cdot \cos \theta.$$


$$3. V = A \cdot h = \underbrace{|\alpha| \cdot |\beta| \cdot \sin \theta}_{|\alpha \times \beta|} \cdot \underbrace{h}_{|r| \cdot \cos \phi}$$

$$= |\alpha \times \beta| \cdot |r| \cdot \cos \phi$$

$$= |r \cdot (\alpha \times \beta)|.$$



$$d = \frac{|\alpha \times \beta|}{|\alpha|}.$$



$$d = \frac{|\alpha \cdot (\beta \times r)|}{|\alpha \times \beta|}$$

$$L: \mathbb{R} \rightarrow \mathbb{R}^3: t \mapsto (a \ b \ c) \cdot (t) = \begin{pmatrix} at \\ bt \\ ct \end{pmatrix}$$

$$T: \mathbb{R}^3 \rightarrow \mathbb{R}: (x, y, z) \mapsto (a \ b \ c) \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$

$$n = (a, b, c), \text{ and } P_0 = (x_0, y_0, z_0).$$

$$n \cdot (P_0 - P) = 0 \Leftrightarrow a(x - x_0) + b(y - y_0) + c(z - z_0) = 0.$$