

Integrating factor.

Consider an equation  $y' + g(x)y = f(x)$ .

Let  $h(x)$  be a unknown. observe that

$$h(x) \cdot y' + h(x) g(x) y = h(x) f(x).$$

suppose that  $(h(x)y)' = h(x)y' + h(x)g(x)y$ .

Then, since  $(h(x)y)' = h(x)y' + h'(x)y$ , so  $(h'(x) - h(x)g(x))y = 0$ .

Therefore,  $h'(x) = h(x)g(x)$ , so

$$\frac{h'(x)}{h(x)} = g(x) \Rightarrow \ln|h(x)| = \int g(x) dx + C \Rightarrow |h(x)| = C_0 \cdot e^{\int g(x) dx}.$$

Example.

Solve  $y' + 2y = e^{2x}$ .

Proof. Multiplying to the both side  $e^{2x} = e^{2x}$ , then

$$e^{2x} y' + e^{2x} \cdot 2y = e^{3x}$$

Since  $e^{2x} y' + e^{2x} \cdot 2y = (e^{2x} y)'$ , so

$$e^{2x} y = \int e^{3x} dx = \frac{1}{3} e^{3x} + C.$$

$$\Rightarrow y = \frac{1}{3} e^x + C \cdot e^{-2x}.$$

Solve  $y' + \frac{1}{x} y = x$ ,  $x > 0$ .

Multiplying  $e^{\int \frac{1}{x} dx} = e^{\ln x} = x$  to the both side, then

$$xy' + y = x^2 \Rightarrow (xy)' = x^2 \Rightarrow xy = \frac{1}{3} x^3 + C \Rightarrow y = \frac{1}{3} x^2 + \frac{C}{x}.$$

