

City Medical Center — Patient Information

Guide

Discharge Instructions, Medications & Post-Care Advice

About This Document

This guide has been prepared by the City Medical Center to help patients understand their discharge instructions, prescribed medications, dietary guidelines, and follow-up requirements. Please read this document carefully before leaving the hospital. If you have any questions, speak with your nurse or doctor before discharge.

Important: This document is for informational purposes only. Always consult your doctor before making any changes to your treatment.

SECTION 1 — PRESCRIBED MEDICATIONS

Your Medication Schedule

You have been prescribed the following medications upon discharge. Take all medications exactly as directed. Do not stop any medication without consulting your doctor, even if you feel better.

Medication	Dose	Frequency	Duration	Take With
Metformin 500mg	1 tablet	Twice daily (morning & night)	3 months	With meals
Amlodipine 5mg	1 tablet	Once daily (morning)	Ongoing	With or without food
Omeprazole 20mg	1 capsule	Once daily (before breakfast)	4 weeks	30 min before eating
Paracetamol 500mg	1–2 tablets	Every 6–8 hours (if needed for pain)	5 days max	With water
Vitamin D3 1000IU	1 tablet	Once daily	3 months	With a meal

■■ IMPORTANT MEDICATION NOTES:

- Metformin should always be taken WITH food to avoid stomach upset.
- Do NOT take Paracetamol if you are already using other products containing paracetamol (e.g. cold medicine).
- Amlodipine may cause mild ankle swelling in the first few weeks — this is normal.
- If you miss a dose, take it as soon as you remember — but SKIP it if your next dose is within 2 hours.
- Store all medications at room temperature, away from direct sunlight and moisture.

SECTION 2 — DIETARY GUIDELINES & RESTRICTIONS

Foods to Eat & Foods to Avoid

Your diet plays a critical role in your recovery. Following the guidelines below will help manage your blood sugar and blood pressure levels effectively.

RECOMMENDED FOODS ■	FOODS TO AVOID ■
Brown rice, whole-wheat bread, oats	White rice, white bread, sugary cereals
Grilled chicken, fish, lentils, eggs	Fried foods, processed meats (sausages, salami)
Fresh vegetables (spinach, broccoli, cucumber)	Pickles, canned vegetables with added salt
Low-fat dairy (yogurt, skimmed milk)	Full-fat cream, butter, cheese
Fresh fruits (apple, pear, berries)	Fruit juices, canned fruits in syrup
Water, herbal teas, coconut water	Soft drinks, energy drinks, alcohol
Nuts and seeds (unsalted)	Salted snacks, chips, biscuits

- Limit salt intake to less than 5 grams (1 teaspoon) per day.
- Eat small meals every 3–4 hours rather than 2–3 large meals.
- Drink at least 8 glasses (2 litres) of water daily.
- Do not skip breakfast — especially important when taking Metformin.

SECTION 3 — PHYSICAL ACTIVITY & REST

Activity Guidelines After Discharge

Rest is essential for recovery, but light activity also helps circulation and prevents complications. Follow this activity plan for the first 4 weeks after discharge:

Week	Allowed Activities	Activities to Avoid
Week 1	Light walking indoors (10–15 min), sitting up regularly	Climbing stairs, lifting anything over 2 kg
Week 2	Short outdoor walks (15–20 min), gentle stretching	Driving, strenuous housework, heavy lifting
Week 3	Increase walking to 30 min daily, light household tasks	Running, sports, prolonged standing
Week 4	Resume most daily activities at a comfortable pace	High-intensity exercise without doctor approval

SECTION 4 — FOLLOW-UP APPOINTMENTS & TESTS

Scheduled Follow-Up

Appointment	When	Doctor / Department	Instructions
Blood sugar check (HbA1c)	After 4 weeks	Endocrinology — Dr. Khan	Fasting 8 hrs before test
Blood pressure review	After 2 weeks	Cardiology — Dr. Raza	Bring medication list

General post-discharge review	After 1 week	General Medicine — OPD	Bring this discharge letter
Kidney function test (Creatinine)	After 6 weeks	Pathology Lab	Early morning, fasting

SECTION 5 — WHEN TO SEEK IMMEDIATE MEDICAL HELP

Go to the Emergency Department or Call 115 if you experience:

- Chest pain or pressure that does not go away
- Sudden difficulty breathing or shortness of breath at rest
- Blood sugar reading below 70 mg/dL (sweating, shaking, confusion)
- Blood pressure above 180/110 mmHg on home monitor
- Severe headache, blurred vision, or sudden weakness in arms/legs
- High fever (above 38.5°C / 101.3°F) lasting more than 24 hours
- Unusual swelling in legs, feet, or face
- Vomiting that prevents you from taking your medications

SECTION 6 — HOSPITAL CONTACT INFORMATION

Department	Phone Number	Hours
Emergency Department	115 / (021) 111-911-911	24 hours, 7 days
General OPD	(021) 3456-7890 Ext. 200	Mon–Sat, 8:00 AM – 6:00 PM
Cardiology Clinic	(021) 3456-7890 Ext. 310	Mon–Fri, 9:00 AM – 2:00 PM
Endocrinology Clinic	(021) 3456-7890 Ext. 325	Tue & Thu, 10:00 AM – 1:00 PM
Pharmacy	(021) 3456-7890 Ext. 105	Mon–Sat, 8:00 AM – 8:00 PM
Patient Relations / Billing	(021) 3456-7890 Ext. 150	Mon–Fri, 9:00 AM – 5:00 PM

Visiting Hours & Hospital Policies

Policy	Details
General Visiting Hours	12:00 PM – 2:00 PM and 5:00 PM – 7:00 PM daily
ICU Visiting Hours	11:00 AM – 12:00 PM and 4:00 PM – 5:00 PM (immediate family only)
Max Visitors Per Patient	2 visitors at a time in general wards; 1 in ICU
Children Policy	Children under 12 not permitted in ICU or post-surgical wards
Re-admission Policy	Call OPD within 48 hours of discharge if condition worsens — do not wait

Medical Records	Request copies from Medical Records office within 7 days of discharge
-----------------	---

City Medical Center | 123 Healthcare Avenue, Karachi | www.citymedical.pk

This document is for the patient named on the discharge form. It does not replace personalised medical advice.

Prepared by: Discharge Planning Unit | Version 2.4 | July 2026