

Masterarbeit

Titel der Bachelorarbeit

Jan Kohlmüller

Datum der Abgabe

Betreuung: Roman Sauer

Fakultät für Mathematik

Karlsruher Institut für Technologie

Contents

1	Introduction	3
2	l^2-Betti numbers	3
2.1	Von Neumann Dimension	3
2.2	l^2 -Betti numbers of G -CW-complexes	4
2.3	l^2 -Betti numbers arising from a group	5
3	Turing dynamical systems	6
4	Computing l^2-Betti numbers	8
4.1	Groupoids	9
4.2	Groupoid ring	10
4.3	Pontryagin duality	11
4.4	l^2 -Betti numbers of groupoids	12
4.5	l^2 -Betti numbers of Turing dynamical systems	12
5	The lamplighter group	12

1 Introduction

2 l^2 -Betti numbers

In this whole chapter G is always a discrete countable group with multiplication \cdot_G .

2.1 Von Neumann Dimension

Definition 2.1. *The group ring RG over a Ring R is the free module over G with the additional multiplication*

$$\sum_{h \in H} \alpha_h \cdot h \cdot_{RG} \sum_{k \in K} \beta_k \cdot k = \sum_{h \in H} \sum_{k \in K} \alpha_h \beta_k \cdot h \cdot_G k$$

with $\alpha_h, \beta_k \in R$ and $H, K \subseteq G$ finite subsets.

Definition 2.2. $l^2(G)$ is the Hilbert space consisting of formal sums $\sum_{g \in G} \lambda_g \cdot g$ such that $\lambda_g \in \mathbb{C}$ and $\sum_{g \in G} |\lambda_g|^2 < \infty$. The scalar product is defined through

$$\left\langle \sum_{g \in G} \alpha_g \cdot g, \sum_{g \in G} \beta_g \cdot g \right\rangle := \sum_{g \in G} \alpha_g \overline{\beta_g}.$$

hier hab ich nicht gezeigt, dass es ein Skalarprodukt ist

Definition 2.3. *The group von Neumann algebra $\mathcal{N}(G)$ is defined as the algebra of G -equivariant bounded operators from $l^2(G)$ to $l^2(G)$ where G acts on $l^2(G)$ by left multiplication.*

We will denote the bounded operators of a Hilbert space H_1 with $\mathcal{B}(H_1)$. If G acts on a Hilbert Space H_2 we will call the subspace of G -equivariant elements of H_2 by H_2^G . Therefore we can also define the group von Neumann algebra through $\mathcal{N}(G) := \mathcal{B}(l^2(G))^G$.

Definition 2.4. *Let $e_G \in G$ be the unit element of G . Then*

$$tr_{\mathcal{N}(G)} : \mathcal{N}(G) \rightarrow \mathbb{C}, A \mapsto \langle A(1 \cdot e_G), 1 \cdot e_G \rangle$$

is called the von Neumann trace on $\mathcal{N}(G)$.

Remark 2.5. *Right multiplication with an element of the Group ring $\mathbb{C}G$ is a G -equivariant operator on $l^2(G)$ and therefore has a well defined von Neumann trace.*

Definition 2.6. *A Hilbert $\mathcal{N}(G)$ -module is a Hilbert space H with a linear isometric G -action such that there exists an isometric linear G -embedding from H to $l^2(G)^n$ for some $n \in \mathbb{N}$.*

Definition 2.7. *Let H be a Hilbert $\mathcal{N}(G)$ -module with corresponding embedding $\iota : H \rightarrow l^2(G)^n$ and let $pr_{\iota(H)} : l^2(G)^n \rightarrow H$ be the corresponding G -equivariant projection. Let*

2 l^2 -Betti numbers

$\iota_i : l^2(G) \rightarrow l^2(G)^n, y \mapsto (x_1, \dots, x_n)$ with $x_j = \delta_{i,j} \cdot y$ the inclusion of the i -th $l^2(G)$ and $pr_i : l^2(G)^n \rightarrow l^2(G), (x_1, \dots, x_n) \mapsto x_i$ the corresponding projection. Then

$$tr_{\mathcal{N}(G)} : \mathcal{B}(H)^G \rightarrow \mathbb{C}, A \mapsto \sum_{i=1}^n tr_{\mathcal{N}(G)}(pr_i \circ \iota \circ A \circ pr_{\iota(H)} \circ \iota_i)$$

is called the von Neumann trace on H .

One can show that the trace is independent of the chosen embedding and therefore well-defined. **eine oder mehrere Projektionen?**

Definition 2.8. Let H be a Hilbert $\mathcal{N}(G)$ -module. We define the von Neumann dimension of H as

$$\dim_{\mathcal{N}(G)}(H) = tr_{\mathcal{N}(G)}(id_H).$$

2.2 l^2 -Betti numbers of G -CW-complexes

Quelle für CW-Komplexe und für Homologie

Definition 2.9. Let X be a CW-complex and $\rho : G \rightarrow \text{Homeo}(X)$ an action of G on X by homeomorphism such that for every open cell $E \subset X$ and every $g \in G$ it holds that $\rho(g)(E)$ is again an open cell and $\rho(g)|_E = id_E$ if $\rho(g)(E) \cap E \neq \emptyset$. Then ρ is called a cellular action.

Definition 2.10. A G -CW-complex is a CW-complex together with a cellular G -action.

Definition 2.11. Let X be a G -CW complex. X is called

- proper if all stabilizer groups are finite,
- free if all stabilizer groups are trivial,
- finite type if for every $n \in \mathbb{N}$ the set of all n -cells consists of only finitely many orbits

Remark 2.12. One can show that the chain complex of a G -CW complex X is a chain complex $C_*(X)$ of left $\mathbb{Z}G$ -modules, i.e. each $C_n(X)$ is a left $\mathbb{Z}G$ -module and the differentials are homomorphisms of these modules.

ZG definieren

Definition 2.13. The l^2 -chain complex of a G -CW complex X is

$$C_*^{(2)}(X) = l^2G \otimes_{\mathbb{Z}G} C_*(X).$$

Hier muss man die Definition evtl noch an das richtige Cstar anpassen, l^2G ist rechtes $\mathbb{Z}G$ modul, was heißt es an chain complex zu tensoren, der Abschluss?

Definition 2.14. Let X be a proper, finite type G -CW complex and $C_*^{(2)}(X)$ the corresponding l^2 -chain complex with its differential $d_*^{(2)}$. Then

$$H_n^{(2)}(X) := \ker(d_n^{(2)}) / \overline{\text{im}(d_{n+1}^{(2)})}$$

is called the n -th l^2 -homology and

$$b_n^{(2)}(X) = \dim_{\mathcal{N}(G)} H_n^{(2)}(X)$$

the n -th l^2 -Betti number.

Remark 2.15. One can show that for a proper, finite type G -CW complex $H_n^{(2)}(X)$ is a Hilbert $\mathcal{N}(G)$ -module and therefore the l^2 -Betti number is in fact well-defined.

2.3 l^2 -Betti numbers arising from a group

Remark 2.16. A Matrix $A \in \mathbb{C}G^{n \times m}$ can be seen as a map from $l^2 G^n \rightarrow l^2 G^m$ via $x \mapsto x^\top \cdot A$. Therefore $\ker(A)$ is a Hilbert $\mathcal{N}(G)$ -module and has a well-defined von-Neumann-dimension.

Theorem 2.17. Let $x \in \mathbb{R}$ and G be a discrete, countable, finitely generated group. The following are equivalent:

- i) There exists a cocompact free finite type G -CW-complex X and a $n \in \mathbb{N}$ such that $b_n^{(2)}(X) = x$
- ii) There exists a Matrix $A \in \mathbb{Q}G^{n \times n}$ for a natural numbers $n \in \mathbb{N}$ such that $\dim_{\mathcal{N}(G)}(\ker(A)) = x$

Proof. "i) \Rightarrow ii)" Let X be a cocompact free finite type G -CW-complex and $n \in \mathbb{N}$ such that $b_n^{(2)}(X) = x$. We define

$$\Delta_n^{(2)} = d_n^{(2)*} d_n^{(2)} + d_{n+1}^{(2)} d_{n+1}^{(2)*},$$

which is a map from $C_n^{(2)}(X)$ to $C_n^{(2)}(X)$ and for which $\ker(\Delta_n^{(2)}) = H_n^{(2)}$ holds. With 2.15 ii) follows. **Warum ist das ne Matrix mit Eintraegen aus \mathbb{Q} ?**

"ii) \Rightarrow i)" Let $n \in \mathbb{N}$ and $A \in \mathbb{Q}G^{n \times n}$. Because $k \cdot A$ has the same kernel as A for every $k \in \mathbb{Z}$ we can assume that $A \in \mathbb{Z}G^{n \times n}$. The corresponding map will be also called A . We will construct a G -CW-complex X with $d_3^{(2)} = A$ and $d_4^{(2)}$ trivial. Therefore $H_3^{(2)}(X) = \ker(A)$ and i) follows.

We denote by $g \in \mathbb{N}$ the number of generators of G . Let Y be the G -covering of $\bigvee_{i=1}^g S^1$ corresponding to an epimorphism $F_g \rightarrow G$, where F_g denotes the free group on g generators. **dazu noch mehr erklaren? stimmt das alles?** Now Y has G as deck transformation group and therefore a natural free G -action. In addition it has exactly g many orbits of 0 and 1-cells and no other cells. We then glue n many 2 on each 0-cell, i.e. we attach them via the attaching map which sends all of S^1 to a single point. Now let $a = \sum_{g \in G} z_g g \in \mathbb{Z}G$ denote the first entry of A . We fix one 2-cell and call it e . We now glue in $G \times D^3$ where $h \times D^3$ is attached via the map which sends S^2 to z_g times $g \cdot h \cdot e$ for each $g \in G$. **Formulierung** We repeat this for every a in the first row of A . At last we repeat the whole process for every row of A where we take a different 2-cell e

for each new row which is glued to the same 0-cell as the first e . The resulting space X is still a free, finite type G -CW-complex. Because we have no 4-cells $d_4^{(2)}$ is trivial and because of the construction $d_3^{(2)} = A$ holds. \square

ker(delta) = homology zeigen? cocompact? Man braucht nur n Zellen, Caleygraph, erwaehnen dass G darauf operiert

Remark 2.18. *One can show that the Theorem also holds if G is not finitely generated. See e.g. [2, p. 371].*

Definition 2.19. *We say that a number $x \in \mathbb{R}$ arises from a discrete countable group G if it fulfills one of the above conditions.*

3 Turing dynamical systems

Let us first give a definition of a Turing machine. For convenience we will only allow 1 and 0 as symbols on our tape.

Definition 3.1. *A Turing machine is a x -tuple $T = (S, \delta, A, R, I)$ where S is a finite set of states and $\delta : S \times \{0, 1\} \rightarrow S \times \mathbb{Z}/2\mathbb{Z} \times \{-1, 1\}$ is called the transition function. $A \subset S$ is called the set of accepting states, $R \subset S$ is called the set of rejecting states and $I \in S$ is called the initial state.*

In addition each Turing machine uses a tape, e.g. a set $\mathbb{Z}/2\mathbb{Z}^{\mathbb{Z}}$ with a head on the element corresponding to the index 0. A Turing machine can operate on an element of $Y \in \mathbb{Z}/2\mathbb{Z}^{\mathbb{Z}}$. We start with the initial state and use the transition function δ to get $\delta(I, Y_0)$. The first coordinate corresponds to the new element Y_0 , the second to the new state and the third corresponds to shifting the tape to the left or the right and therefore getting a new Y_0 . We repeat this step until we get a state in $A \cup R$. We say, that the Turing machine accepts Y if we get a state in A after finitely many steps and it rejects Y if we get a state in R . We say that the Turing machine holds for Y if it accepts or rejects it. We will only look at Turing machines which hold for every input. **mach das mit den Y0 mal ordentlich, du kacknoob! input = Y**

For the purpose of calculating l^2 -Betti numbers of groups we need to extend this definition. For the rest of this chapter let (X, μ) be a probability measure space divided into finitely many disjoint measurable subsets X_i . Let Γ be a countable discrete group and ρ be a right measure preserving action of Γ on X . We now choose 3 disjoint subsets A, R, I , where each of them is a union of certain X_i . They will be called the accepting set, the rejecting set and the initial set. In addition we choose a $\gamma_i \in \Gamma$ for each $X_i \subset X$ such that for each i with $X_i \subset A$ or $X_i \subset R$ it holds that $\gamma_i = e$ where e is the neutral element in Γ . Let $Ind : X \rightarrow \mathbb{N}$ be the map which assigns to each $x \in X$ the corresponding index of the X_i it is contained in.

Definition 3.2. *The map*

$$T_X : X \rightarrow X, x \mapsto \rho(\gamma_{Ind(x)})(x)$$

is called the Turing map. All the data required for a Turing map together with the map itself will be called a Turing dynamical system and will be denoted by (T_X)

all data spezifizieren Mostly we don't look at the map T_X but the map T_X^∞ . We say that the Turing dynamical system accepts an input $y \in I$ if $T_X^\infty(y) \in A$ and it rejects it if $T_X^\infty(y) \in R$. **Tinfy definiert-hält**

Example 3.3. Now let $T = (S, \delta, \tilde{A}, \tilde{R}, \tilde{I})$ be a Turing machine. We will show how to emulate it with a Turing dynamical system. Lets first fix notation. For a subset M of a set $N^\mathbb{Z}$ with $M = \{(n_i)_{i \in \mathbb{Z}} \in N^\mathbb{Z} | n_{-k} = m_{-k}, \dots, n_l = m_l\}$ **erklären, was es mit k und l auf sich hat** we simply write $[m_{-k} \dots \underline{m_0} \dots m_l]$ and $[m_{-k} \dots \underline{m_0} \dots m_l][\sigma]$ for $M \times \{\sigma\}$. We define $X = \mathbb{Z}/2\mathbb{Z}^\mathbb{Z} \times S$ and $\Gamma = \text{Bij}(S) \times \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}$. **in unserem fall ist sigma usw., S Menge, Bij Grupper erwähnen Stimmt das?** The action of Γ on X is defined by the following rules:

- $\text{Bij}(S)$ acts on S in the natural way.
- The generator $\bar{1}$ of the group $\mathbb{Z}/2\mathbb{Z}$ acts on the $\mathbb{Z}/2\mathbb{Z}^\mathbb{Z}$ part of X by adding $\bar{1}$ to the element with index 0.
- The generator 1 of the group \mathbb{Z} acts also on the $\mathbb{Z}/2\mathbb{Z}^\mathbb{Z}$ part by shifting every element 1 to the right, e.g. increasing the index of every element by 1. **nochmal ausführlich**

We will now use the transition function to construct the Turing map. We choose the following devision of X :

$$X = \bigcup_{x \in \mathbb{Z}/2\mathbb{Z}, \sigma \in S} [x][\sigma]$$

For $X_i = [x][\sigma]$ we get $\delta(\sigma, x) = (\tilde{\sigma}, \alpha, \beta)$. **sigma und x reihenfolge Satz neu** Let $\tau \in \text{Bij}(S)$ map σ to $\tilde{\sigma}$ and $y \in \mathbb{Z}/2\mathbb{Z}$ be $y = x - \alpha$, i.e. an Automorphism **Quatsch Operation** which maps x to α . Our element γ_i corresponding to X_i is given by

$$\gamma_i = \begin{cases} (id, \bar{0}, 0) & \text{if } X_i \subset A \cup R \\ (\tau, y, \beta) & \text{else} \end{cases}$$

tau nicht eindeutig We only need to define A , R and I . A is given by

$$A = \bigcup_{x \in \mathbb{Z}/2\mathbb{Z}, \sigma \in \tilde{A}} [x][\sigma]$$

which is of course a union of some X_i . R and I can be defined analogous.

Abschlusssatz, reihenfolge vertauschen

To conclude this chapter we will give some definition which we will use later on.

Definition 3.4. The first fundamental set $\mathcal{F}_1(T_X)$ is the subset of I consisting of those points x with $T_X^\infty(x) \in A$ and there is no point y with $T_X(y) = x$. The second fundamental set is the subset of A defined as $\mathcal{F}_2(T_X) = T_X^\infty(\mathcal{F}_1(T_X))$. The first (second) fundamental value $\Omega_1(T_X)(\Omega_2(T_X))$ is the measure of the corresponding fundamental set.

Definition 3.5. We say that a Turing dynamical system (T_X)

- stops at any configuration if $T_X^\infty(x) \in A \cup R$ for almost all $x \in X$
- has disjoint accepting chains if $T_X^\infty(x) \neq T_X^\infty(y)$ for almost all $x, y \in I$ with $x \neq y$
- does not restart if the set $T_X(X) \cap I$ has measure 0

Definition 3.6. A Turing dynamical system (T_X) where X is a infinite product of $\mathbb{Z}/2\mathbb{Z}$, each X_i is of the form $X_i = \{(x_j) \in X \mid x_k = 0 \ \forall k \in I_1, x_j = 1 \ \forall j \in I_2 \mid I_1, I_2 \subset J \text{ finite}\}$, the action of Γ is by continuous group automorphisms and (T_X) stops an any configuration, has disjoint accepting chains and does not restart is called a computing Touring dynamical system.

Will ich das wirklich so nennen? Ist X richtig definiert? Warum cont group autos? Indexmenge J beim Produkt, andere Xi defintion maß

Before we give an example of a computing Turing dynamical system we fix the notation of the shifting operation, because we will need it later on.

Definition 3.7. An element $z \in \mathbb{Z}$ acts on $\mathbb{Z}/2\mathbb{Z}^{\mathbb{Z}} = \prod_{i \in \mathbb{Z}} \mathbb{Z}/2\mathbb{Z}$ by decreasing every index by z . We call this action the shift action and denote it by ζ . \mathbb{Z} acts on $\bigoplus_{i \in \mathbb{Z}} \mathbb{Z}/2\mathbb{Z}$ in the same way which we will also call ζ .

Example 3.8. Let T be a read-only Turing machine which holds for every input. Let (T_X) be the corresponding Turing dynamical system as constructed in 3.3. We assume that T is in such a way that (T_X) has disjoint accepting chains (which is not always the case). We will construct a computable Turing dynamical system from T . We can easily assure that (T_X) does not restart by adding a new state I' to T with $\delta(I', x) = \delta(I, x) \ \forall x \in \{0, 1\}$ and setting I' as the initial state of T . Because we do not change the symbols of the tape it suffices to use $\Gamma = \mathbb{Z} \times \text{Bij}(S)$ where \mathbb{Z} operates on $X = \mathbb{Z}/2\mathbb{Z}^{\mathbb{Z}} \times S$ via shifting and $\text{Bij}(S)$ acts on S in the natural way. Because a Turing machine has only finitely many states there exists a number $n \in \mathbb{N}$ such that $|S| < 2^n$. We identify every state with a different element $z \in \mathbb{Z}/2\mathbb{Z}^n$ with $z \neq 0$. We can then assume that $X = \mathbb{Z}/2\mathbb{Z}^{\mathbb{Z}} \times \mathbb{Z}/2\mathbb{Z}^n$ by filling $\mathbb{Z}/2\mathbb{Z}^n$ with "dummy states" which will never be used. Because $\text{Aut}(\mathbb{Z}/2\mathbb{Z}^n)$ acts transitively on all of $\mathbb{Z}/2\mathbb{Z}^n \setminus \{0\}$ we can assume that $\Gamma = \mathbb{Z} \times \text{Aut}(S)$. Because in the sets X_i in 3.3 there are only $n+1$ fixed components they are of the desired form and therefore the resulting Turing dynamical system is computing.

4 Computing l^2 -Betti numbers

In this chapter we want to prove the following central theorem which connects l^2 -Betti numbers to the concept of Turing dynamical systems.

Theorem 4.1. Let (T_X) be a computing Touring dynamical system. Then $\mu(I) - \Omega_1(T_X)$ is a l^2 -Betti number arising from $\hat{X} \rtimes_{\hat{\rho}} \Gamma$ where \hat{X} and $\hat{\rho}$ are the Pontryagin duals of the corresponding group or map.

We will give the definition of Pontryagin duals and the semidirect product later in this

chapter. The rest of this chapter is used to prove this theorem and give the needed definitions.

4.1 Groupoids

We begin by giving some algebraic definitions, namely the construct of groupoids which extends the definition of groups **sicher bin ich mir dabei jetzt net**. Therefore we require some basic knowledge about categories. **Quelle zu Kategorientheorie**

Definition 4.2. A groupoid is a small category whose morphisms are all invertible.

Example 4.3. Let G be a group. We want to see how we can express G as a groupoid. Let $\mathcal{G}_0 = \{\bullet\}$ be the set with only one element and \mathcal{G} be the category with objects G_0 and morphisms G from \bullet to \bullet . The composition of morphisms in \mathcal{G} is the same as the multiplication in G . Then \mathcal{G} is a small category and every morphism is invertible because every group element has an inverse. Therefore \mathcal{G} is a groupoid.

We will always denote the set of objects of a groupoid \mathcal{G} by \mathcal{G}_0 and we can identify it with a subset of the set of morphisms of \mathcal{G} by the embedding $\mathbf{1} : \mathcal{G}_0 \rightarrow \text{mor}(\mathcal{G}), x \mapsto [\text{id} : x \rightarrow x]$ which sends every object to the corresponding identity morphism. Therefore we will only look at $\text{mor}(\mathcal{G})$ and also call it \mathcal{G} .

Definition 4.4. For every Groupoid \mathcal{G} we define the maps $s : \mathcal{G} \rightarrow \mathcal{G}_0, [f : X \rightarrow Y] \mapsto X$ and $r : \mathcal{G} \rightarrow \mathcal{G}_0, [f : X \rightarrow Y] \mapsto Y$ which will be called source and range map.

das mit den Pfeilen ist doof

Definition 4.5. A discrete measurable groupoid is a groupoid where \mathcal{G} is also a measurable space, s, r and the maps gained through inverting or composition are all measurable and the fibers of s and r are countable. If in addition we have a measure μ such that

$$\int_{\mathcal{G}_0} |r^{-1}(x) \cap U| d\mu(x) = \int_{\mathcal{G}_0} |s^{-1}(x) \cap U| d\mu(x) \quad \forall U \subset \mathcal{G}$$

holds we call \mathcal{G} discrete measured

Warum braucht man das eigentlich nochmal, was bedeutet das genau und stimmt das eigentlich? inverting map

Definition 4.6. Let Γ be a discrete, countable group, X a probability measure space and $\rho : \Gamma \rightarrow \text{Bij}(X)$ a right measure preserving action. The action groupoid $\mathcal{G}(\rho)$ is the groupoid with objects X and morphisms $X \times \Gamma$ such that $s(x, \gamma) = x$ and $r(x, \gamma) = \rho(\gamma)(x)$. The composition of morphisms is defined through $(x, \gamma_1) \cdot_{\mathcal{G}(\rho)} (\rho(\gamma_1)(x), \gamma_2) = (x, \gamma_1 \cdot \gamma_2)$ and the inverse of (x, γ) is $(\rho(\gamma)(x), \gamma^{-1})$.

Beispiel oder Erklärung und wie das bei TDS ist

4.2 Groupoid ring

A fundamental concept used for the calculation of l^2 -Betti numbers of groups was the group ring. We will transfer this to the notion of groupoids.

Definition 4.7. Let U be a subset of \mathcal{G}_0 . A measurable edge is a map $\Phi : U \rightarrow \mathcal{G}$ such that $s \circ \Phi = \text{id}$ and $r \circ \Phi$ is injective.

Warum heißt das eigentlich messbar? From the definition we see, that defining a measurable edge means taking a subset of \mathcal{G}_0 and associating a morphism to every object such that the morphism starts in this object and no two such morphisms end in the same object. We want to define the inverse of a measurable edge in such a way, that it is also a measurable edge. Therefore it does not suffice to take the inverse of the map Φ . Instead we invert every morphism in the image of Φ .

Definition 4.8. Let $\Phi : U \rightarrow \mathcal{G}$ be a measurable edge. The inverse of Φ will be called Φ^{-1} and is defined as $\Phi^{-1} : r(\text{Im}(\Phi)) \rightarrow \mathcal{G}$ such that Φ^{-1} is a measurable edge and $\Phi^{-1} \circ r \circ \Phi = \text{id}$.

am ende muss noch invertierung ans phi

We now come to the definition of the groupoid ring. We want it to be ring of operators of $L^2(\mathcal{G})$ **Das sollte ich vorher mal irgendwann definiert haben**, such that it is in a way generated by measurable edges. For a measurable edge Φ we define a operator $\tilde{\Phi}$ through

$$\tilde{\Phi} : L^2(\mathcal{G}) \rightarrow L^2(\mathcal{G}), F \mapsto \left[\tilde{\Phi}(F) : \mathcal{G} \rightarrow \mathbb{C}, \gamma \mapsto \begin{cases} F(\gamma \cdot_{\mathcal{G}} \Phi^{-1}(r(\gamma))) & \text{if } r(\gamma) \in \text{Dom}(\Phi^{-1}) \\ 0 & \text{otherwise} \end{cases} \right].$$

da sollte man ein Bild zu malen In addition for a map $f \in L^\infty(\mathcal{G}_0)$ we also define a operator \tilde{f} on $L^2(\mathcal{G})$ through

$$\tilde{f} : L^2(\mathcal{G}) \rightarrow L^2(\mathcal{G}), F \mapsto [\tilde{f}(F) : \mathcal{G} \rightarrow \mathbb{C}, \gamma \mapsto F(\gamma) \cdot f(r(\gamma))].$$

Definition 4.9. For a groupoid \mathcal{G} the groupoid ring $\mathbb{C}\mathcal{G}$ is the ring of bounded operators on $L^2(\mathcal{G})$ generated by all measurable edges and all elements of $L^\infty(\mathcal{G})$.

We always denote the elements of $\mathbb{C}\mathcal{G}$ by a linear combination $\sum_{i \in I} \tilde{\Phi}_i \cdot_{\mathbb{C}\mathcal{G}} \tilde{f}_i$ where Φ is a measurable edge, $f \in L^\infty(\mathcal{G}_0)$ and I is a finite set. One can show that each element of $\mathbb{C}\mathcal{G}$ can be (although non-uniquely) represented in such a way. **Evtl. könnte ich dazu noch ein bisschen was schreiben**

Example 4.10. Let G be a group and \mathcal{G} the corresponding groupoid as in Example 4.3. We want to show that the groupoid ring $\mathbb{C}\mathcal{G}$ is isomorphic to the g $\mathbb{C}G$. Because we have only one element in \mathcal{G}_0 the only measurable edges we have are the maps from \bullet to a specific group element of G . The elements of $L^\infty(\mathcal{G}_0)$ are just maps from \bullet to a specific element in \mathbb{C} . Therefore $L^\infty(\mathcal{G}_0)$ is isomorphic to \mathbb{C} . Now let $\Phi_1 : \bullet \mapsto g_1$ and $\Phi_2 : \bullet \mapsto g_2$ be measurable edges.

Das muss ich noch zuende machen

Remark 4.11. Let $\mathcal{G}(\rho)$ be an action Groupoid of $\rho : \Gamma \rightarrow \text{Bij}(X)$. For each element $\gamma \in \Gamma$ we get a measurable edge $\bar{\gamma} : X \rightarrow x \times \Gamma$ which maps $x \in X$ to (x, γ) .

4.3 Pontryagin duality

We will now create a link between the groupoid ring and the group ring. For this we need the Pontryagin duals. For the rest of this chapter X is a locally compact abelian group, Γ is another group acting on X by continuous group automorphisms. We will call this action $\rho : \Gamma \rightarrow \text{Aut}(X)$.

Definition 4.12. A character of X is a homomorphism $\hat{x} : X \rightarrow S$, where S is the multiplicative group of complex numbers of absolute value 1. The group of all characters of X is called the character group or Pontryagin dual group of X and is denoted by \hat{X} .

Beweis, dass das duale eine Gruppe ist?

Definition 4.13. The Pontryagin dual $\hat{\rho} : \Gamma \rightarrow \text{Aut}(\hat{X})$ of the action ρ is an action on \hat{X} and defined as $\hat{\rho}(\gamma)(f)(x) = f(\rho(\gamma^{-1})(x))$ for $f \in \hat{X}$ and $x \in X$.

Um das ganze Auto-Zeug muss man sich mal noch Gedanken machen.

Remark 4.14. An element of the Pontryagin dual $\hat{x} \in \hat{X}$ can also be seen as an element of $L^\infty(X)$. We call this element $P(x)$

Definition 4.15. Let N, H be groups and $\Theta : H \rightarrow \text{Aut}(N)$ a homomorphism. The semidirect product $N \rtimes_\Theta H$ of H and N is defined on the Set $N \times H$ with multiplication

$$(n_1, h_1) \cdot_{N \rtimes_\Theta H} (n_2, h_2) = (n_1 \cdot_N \Theta(h_1)(n_2), h_1 \cdot_H h_2).$$

For more information about semidirect products see e.g. [4].

Theorem 4.16. The map

$$P \otimes 1 : \mathbb{C}(\hat{X} \rtimes_{\hat{\rho}} \Gamma) \rightarrow \mathbb{C}\mathcal{G}(\rho), \sum_I c_i \cdot (\hat{x}_i, \gamma_i) \mapsto \sum_I c_i \cdot \widetilde{P(\hat{x}_i)} \cdot_{\mathbb{C}\mathcal{G}(\rho)} \tilde{\gamma}_i$$

with $c_i \in \mathbb{C}$, $(\hat{x}_i) \in \hat{X}$ and $\gamma_i \in \Gamma$ is

i) a ringhomomorphism

ii) trace-preserving

Proof. i) Because of the definition it is clear that $P \otimes 1$ preserves the addition. So let $c_i \in \mathbb{C}$, $(\hat{x}_i) \in \hat{X}$ and $\gamma_i \in \Gamma$ with $i \in \{1, 2\}$. It follows that

$$\begin{aligned} & P \otimes 1(c_1 \cdot (\hat{x}_1, \gamma_1) \cdot_{\mathbb{C}(\hat{X} \rtimes_{\hat{\rho}} \Gamma)} c_2 \cdot (\hat{x}_2, \gamma_2)) \\ &= P \otimes 1(c_1 \cdot c_2 \cdot (\hat{x}_1, \gamma_1) \cdot_{N \rtimes_{\hat{\rho}} H} (\hat{x}_2, \gamma_2)) \\ &= P \otimes 1(c_1 \cdot c_2 \cdot (\hat{x}_1 \cdot_{\hat{X}} \hat{\rho}(\gamma_1)(\hat{x}_2), \gamma_1 \cdot_\Gamma \gamma_2)) \\ &= c_1 \cdot c_2 \cdot \widetilde{P(\hat{x}_1 \cdot_{\hat{X}} \hat{\rho}(\gamma_1)(\hat{x}_2))} \cdot_{\mathbb{C}\mathcal{G}(\rho)} \widetilde{\gamma_1 \cdot_\Gamma \gamma_2}. \end{aligned}$$

5 The lamplighter group

From the definition of the groupoid ring it follows directly that the last term equals

$$c_1 \cdot c_2 \cdot \widetilde{P(\hat{x}_1)} \cdot_{\mathbb{C}\mathcal{G}(\rho)} P(\widetilde{\hat{\rho}(\gamma_1)(\hat{x}_2)}) \cdot_{\mathbb{C}\mathcal{G}(\rho)} \widetilde{\gamma}_1 \cdot_{\mathbb{C}\mathcal{G}(\rho)} \widetilde{\gamma}_2.$$

Because elements of \mathbb{C} commute with all elements in the groupoid ring it remains to show that

$$P(\widetilde{\hat{\rho}(\gamma_1)(\hat{x}_2)}) \cdot_{\mathbb{C}\mathcal{G}(\rho)} \widetilde{\gamma}_1 = \widetilde{\gamma}_1 \cdot_{\mathbb{C}\mathcal{G}(\rho)} \hat{x}_2$$

in the groupoid ring. Let $F \in L^2\mathcal{G}(\rho)$ and $\alpha \in \mathcal{G}$. Then the left side equals

$$P(\widetilde{\hat{\rho}(\gamma_1)(\hat{x}_2)}) \cdot_{\mathbb{C}\mathcal{G}(\rho)} \widetilde{\gamma}_1(F)(\alpha) = \hat{x}_2(\rho(\gamma_1^{-1})(r(\alpha))) \cdot F(\alpha \cdot_{\mathcal{G}(\rho)} \bar{\gamma}_1^{-1}(r(\alpha)))$$

and the right side equals

$$\widetilde{\gamma}_1 \cdot_{\mathbb{C}\mathcal{G}(\rho)} \hat{x}_2(F)(\alpha) = \widetilde{\gamma}_1(\hat{x}_2(r(\alpha)) \cdot F(\alpha)) = \hat{x}_2(r(\alpha \cdot_{\mathcal{G}(\rho)} \bar{\gamma}_1^{-1}(r(\alpha)))) \cdot F(\alpha \cdot_{\mathcal{G}(\rho)} \bar{\gamma}_1^{-1}(r(\alpha)))$$

which is the same.

ii)

□

braucht man *?, Elemente aus C an GR multiplizieren , mehr erklaeren?

4.4 l^2 -Betti numbers of groupoids

4.5 l^2 -Betti numbers of Turing dynamical systems

5 The lamplighter group

Definition 5.1. *The lamplighter group L is defined as*

$$L = \left(\bigoplus_{i \in \mathbb{Z}} \mathbb{Z}/2\mathbb{Z} \right) \rtimes_{\zeta} \mathbb{Z}.$$

where ζ is the shift operation defined in 3.7.

Lets first look at the Turing dynamical system corresponding to a read-only Turing machine as constructed in 3.8. By using 4.1 we see that $\mu(I) - \Omega_1(T_X)$ is a l^2 -Betti number arising from $\hat{X} \rtimes_{\hat{\rho}} \Gamma$.

Theorem 5.2. *Every rational number arises from the lamplighter group.*

References

- [1] Grabowski, Łukasz: *On Turing dynamical systems and the Atiyah problem*, Springer, 2014
- [2] Lück, Wolfgang *L^2 -Invariants: Theory and Applications to Geometry and K-Theory*, Springer, 2002
- [3] Hatcher, Allen *Algebraic topology*, Cambridge University Press, 2015
- [4] Aluffi, Paolo *Algebra: Chapter 0*, in: Graduate Studies in Mathematics Volume 104, American Mathematical Society

Erklärung

Hiermit versichere ich, dass ich diese Arbeit selbständig verfasst und keine anderen, als die angegebenen Quellen und Hilfsmittel benutzt, die wörtlich oder inhaltlich übernommenen Stellen als solche kenntlich gemacht und die Satzung des Karlsruher Instituts für Technologie zur Sicherung guter wissenschaftlicher Praxis in der jeweils gültigen Fassung beachtet habe.

Ort, den Datum