

TWO-DIMENSIONAL “HAUL-OUT” MODEL

Brian M. Brost

23 APR 2015

Implementation

The file `haulout.2d.sim.R` simulates data according to the model statement presented below, and `haulout.2d.mcmc.R` contains the MCMC algorithm for model fitting.

Model statement

Let \mathcal{S} be the support of the movement process and $\tilde{\mathcal{S}}$ be the support of haul-out sites. Here, both are defined in 2-dimensional space. Note that \mathcal{S} and $\tilde{\mathcal{S}}$ overlap, and that $\mathcal{S} \cap \tilde{\mathcal{S}} = \tilde{\mathcal{S}}$. The domain defined by \mathcal{S} therefore represents at-sea locations or locations of the individual while milling adjacent to the haul-out site. Also note that $\bar{\mathcal{S}}$, the complement of \mathcal{S} , represents inaccessible locations (i.e., terrestrial sites that are not haul-outs). Let $\mathbf{s}_t = (s_{1,t}, s_{2,t})'$ and $\boldsymbol{\mu}_t = (\mu_{1,t}, \mu_{2,t})'$, for $t = 1, \dots, T$, be observed and true locations respectively.

$$\begin{aligned} \mathbf{s}_t &\sim \text{N}(\boldsymbol{\mu}_t, \sigma^2 \mathbf{I}) \\ \boldsymbol{\mu}_t &\sim \begin{cases} \text{Unif}(\tilde{\mathcal{S}}), & z_t = 1 \\ \text{Unif}(\mathcal{S}), & z_t = 0 \end{cases} \\ z_t &\sim \text{Bern}(p) \\ p &\sim \text{Beta}(\alpha, \beta) \\ \sigma^2 &\sim \text{IG}(r, q) \end{aligned}$$

Full conditional distributions

True locations ($\boldsymbol{\mu}_t$):

$$\begin{aligned} [\boldsymbol{\mu}_t | \cdot] &\propto [\mathbf{s}_t | \boldsymbol{\mu}_t, \sigma^2] [\boldsymbol{\mu}_t | z_t, \mathcal{S}, \tilde{\mathcal{S}}] \\ &\propto [\mathbf{s}_t | \boldsymbol{\mu}_t, \sigma^2] [\boldsymbol{\mu}_t | \tilde{\mathcal{S}}]^{z_t} [\boldsymbol{\mu}_t | \mathcal{S}]^{1-z_t}. \end{aligned}$$

For $z_t = 1$,

$$\begin{aligned} [\boldsymbol{\mu}_t | \cdot] &\propto [\mathbf{s}_t | \boldsymbol{\mu}_t, \sigma^2] [\boldsymbol{\mu}_t | \tilde{\mathcal{S}}]^{z_t} \\ &\propto [\mathbf{s}_t | \boldsymbol{\mu}_t, \sigma^2] \\ &\propto \exp \left\{ -\frac{1}{2} \left((\mathbf{s}_t - \boldsymbol{\mu}_t)' (\sigma^2 \mathbf{I})^{-1} (\mathbf{s}_t - \boldsymbol{\mu}_t) \right) \right\} \\ &\propto \exp \left\{ -\frac{1}{2} \left(\mathbf{s}_t' (\sigma^2 \mathbf{I})^{-1} \mathbf{s}_t - 2 \mathbf{s}_t' (\sigma^2 \mathbf{I})^{-1} \boldsymbol{\mu}_t + \boldsymbol{\mu}_t' (\sigma^2 \mathbf{I})^{-1} \boldsymbol{\mu}_t \right) \right\} \\ &\propto \exp \left\{ -\frac{1}{2} \left(-2 \mathbf{s}_t' (\sigma^2 \mathbf{I})^{-1} \boldsymbol{\mu}_t + \boldsymbol{\mu}_t' (\sigma^2 \mathbf{I})^{-1} \boldsymbol{\mu}_t \right) \right\} \\ &\propto \text{N}(\mathbf{A}^{-1} \mathbf{b}, \mathbf{A}^{-1}) \end{aligned}$$

where $\mathbf{A} = (\sigma^2 \mathbf{I})^{-1}$ and $\mathbf{b} = \mathbf{s}_t' (\sigma^2 \mathbf{I})^{-1}$; therefore, $[\boldsymbol{\mu}_t | \cdot] = \text{N}(\mathbf{s}_t, \sigma^2 \mathbf{I})$ for $z_t = 1$. Note that proposed values for $\boldsymbol{\mu}_t$ not in $\tilde{\mathcal{S}}$ are rejected. For $z_t = 0$,

$$[\boldsymbol{\mu}_t | \cdot] = \text{N}(\mathbf{s}_t, \sigma^2 \mathbf{I}).$$

In this case, note that proposed values for $\boldsymbol{\mu}_t$ not in \mathcal{S} are rejected.

Haul-out indicator variable (z_t):

$$[z_t|\cdot] \propto [\boldsymbol{\mu}_t|\tilde{\mathcal{S}}]^{z_t}[\boldsymbol{\mu}_t|\mathcal{S}]^{1-z_t}[z_t|p].$$

For all $\boldsymbol{\mu}_t \notin \tilde{\mathcal{S}}$, let $z_t = 0$. For all $\boldsymbol{\mu}_t \in \tilde{\mathcal{S}}$, sample z_t from

$$\begin{aligned} [z_t|\cdot] &\propto [\boldsymbol{\mu}_t|\tilde{\mathcal{S}}]^{z_t}[\boldsymbol{\mu}_t|\mathcal{S}]^{1-z_t}[z_t|p] \\ &= \text{Bern}(\tilde{p}), \end{aligned}$$

where

$$\begin{aligned} \tilde{p} &= \frac{p[\boldsymbol{\mu}_t|\tilde{\mathcal{S}}]}{p[\boldsymbol{\mu}_t|\tilde{\mathcal{S}}] + (1-p)[\boldsymbol{\mu}_t|\mathcal{S}]} \\ &= \frac{p(|\tilde{\mathcal{S}}|^{-1})}{p(|\tilde{\mathcal{S}}|^{-1}) + (1-p)(|\mathcal{S}|^{-1})}. \end{aligned}$$

The notation $|\cdot|$ denotes the area of the respective domain.

Probability of being hauled-out (p):

$$\begin{aligned} [p|\cdot] &\propto \prod_{t=1}^T [z_t|p][p] \\ &\propto \prod_{t=1}^T p^{z_t}(1-p)^{1-z_t} p^{\alpha-1}(1-p)^{\beta-1} \\ &\propto p^{\sum_{t=1}^T z_t} (1-p)^{\sum_{t=1}^T (1-z_t)} p^{\alpha-1}(1-p)^{\beta-1} \\ &= \text{Beta}\left(\sum_{t=1}^T z_t + \alpha, \sum_{t=1}^T (1-z_t) + \beta\right) \end{aligned}$$

Error in the observation process (σ^2):

$$\begin{aligned} [\sigma^2|\cdot] &\propto \prod_{t=1}^T [\mathbf{s}_t|\boldsymbol{\mu}_t, \sigma^2][\sigma^2] \\ &\propto \prod_{t=1}^T |\sigma^2 \mathbf{I}|^{-1/2} \exp\left\{-\frac{1}{2} \left((\mathbf{s}_t - \boldsymbol{\mu}_t)' (\sigma^2 \mathbf{I})^{-1} (\mathbf{s}_t - \boldsymbol{\mu}_t)\right)\right\} (\sigma^2)^{-(q+1)} \exp\left\{-\frac{1}{\sigma^2 r}\right\} \\ &\propto \prod_{t=1}^T (\sigma^2)^{-1} \exp\left\{-\frac{1}{2\sigma^2} ((\mathbf{s}_t - \boldsymbol{\mu}_t)' (\mathbf{s}_t - \boldsymbol{\mu}_t))\right\} (\sigma^2)^{-(q+1)} \exp\left\{-\frac{1}{\sigma^2 r}\right\} \\ &\propto (\sigma^2)^{-(T+q+1)} \exp\left\{-\frac{1}{\sigma^2} \left(\frac{\sum_{t=1}^T (\mathbf{s}_t - \boldsymbol{\mu}_t)' (\mathbf{s}_t - \boldsymbol{\mu}_t)}{2} + \frac{1}{r}\right)\right\} \\ &= \text{IG}\left(\left(\frac{\sum_{t=1}^T (\mathbf{s}_t - \boldsymbol{\mu}_t)' (\mathbf{s}_t - \boldsymbol{\mu}_t)}{2} + \frac{1}{r}\right)^{-1}, T+q\right). \end{aligned}$$

Note that the current version of haulout.2d.mcmc.R contains code for the conjugate update of σ^2 presented above, but this code is currently 'commented' out. Instead, error is modeled as $[\sigma|\cdot] \sim \text{Unif}(a, b)$, and the update for σ proceeds using Metropolis-Hastings.