



Mathematical
Institute

Finite element methods

YUJI NAKATSUKASA

Many slides by Ricardo Ruiz Baier

Some pictures from Andy Wathen

Mathematical Institute, Oxford

Computational Techniques

InFoMM Centre for Doctoral Training

Michaelmas Term 2019, Week 6

November 21, 2019



Oxford
Mathematics




- References (among many):
 - H. Elman, D. Silvester, A. Wathen, Finite Elements and Fast Iterative Solvers, OUP, 2014
 - P. E. Farrell, Finite Element Methods for PDEs, C6.4 lecture notes
 - E. Süli, Lecture Notes for FEM for PDEs
people.maths.ox.ac.uk/suli/fem.pdf
 - Brenner and Scott, The Mathematical Theory of Finite Element Methods, Springer, 2007 (advanced, mathematical analysis)
- For software, we suggest FEniCS (demo session later by Fede Danieli) or IFISS (Elman-Silvester-Wathen book)
- For advanced questions, we suggest asking our amazing local experts!
 - P. E. Farrell (theory, programming & applications)
 - E. Süli (analysis, theory)
 - A. Wathen (preconditioning, LA aspects)

$$au_{xx} + bu_{xy} + cu_{yy} + du_x + eu_y + fu = g$$

- elliptic: $b^2 - 4ac < 0$
e.g. Poisson problem $\nabla^2 u = f$
- parabolic: $b^2 - 4ac = 0$
e.g. heat equation $u_t = \nabla^2 u - f$
- hyperbolic $b^2 - 4ac > 0$
e.g. wave equation $u_{tt} = c^2 u_{xx}$

e.g. consider Poisson problem $\nabla^2 u = f$

- Approximate solution u with **piecewise polynomial** $\hat{u}(x) = \sum_{i=1}^n c_i \phi_i(x)$
e.g. $\phi_i(x)$: hat function
- **Integration by parts** + divergence thm to 'move' one derivative, 'relax' smoothness requirement
- Find c_i via '**weak solution**' by requiring **Galerkin** condition: 'residual is orthogonal to test functions', in LA terms,
 $Q^T(Ax - b) = 0$ (recall least-squares $\min_x \|Ax - b\|$, and CG $Q^T(AQy - b) = 0$)



First, we stick to the Poisson problem on a bounded domain (in **strong** form)

$$-\nabla^2 u = f \quad \text{in } \Omega \subset \mathbb{R}^d, \quad u = 0 \text{ on } \partial\Omega.$$

Weak formulation

- multiply by a test function $v \in V$
- integrate by parts

$$\int_{\Omega} \nabla u \cdot \nabla v \, dx - \int_{\partial\Omega} (\nabla u \cdot \mathbf{n}) v \, ds = \int_{\Omega} f v \, dx$$

- BCs $\rightarrow V = \{w \in H^1(\Omega) : w = 0 \text{ on } \partial\Omega\}$
(roughly, $H^k(\Omega)$: k -times differentiable with k th der $\in L^2(\Omega)$)
- find $u \in V$ such that

$$\int_{\Omega} \nabla u \cdot \nabla v \, dx = \int_{\Omega} f v \, dx, \quad \forall v \in V$$

$$-\nabla^2 u = f \quad \text{in } \Omega \subset \mathbb{R}^d, \quad u = 0 \text{ on } \partial\Omega.$$

1. Multiply by a test function $v \in V$ and integrate:

$$-\int_{\Omega} v \nabla^2 u dx = \int_{\Omega} f v dx$$

2. Integrate by parts: first recall product rule

$$v \nabla^2 u + \nabla v \cdot \nabla u = \nabla \cdot (v \nabla u)$$

Integrate $-\int_{\Omega} v \nabla^2 u dx = \int_{\Omega} \nabla v \cdot \nabla u dx - \int_{\Omega} \nabla \cdot (v \nabla u) dx$. By diver. thm. $\int_{\Omega} \nabla \cdot (v \nabla u) dx = \int_{\partial\Omega} (\nabla u \cdot \mathbf{n}) v ds$ (\mathbf{n} : outward normal vec)

$$\int_{\Omega} \nabla u \cdot \nabla v dx - \int_{\partial\Omega} (\nabla u \cdot \mathbf{n}) v ds = \int_{\Omega} f v dx \quad (1)$$

→ reduced regularity: **before** ($u \in C^2(\bar{\Omega})$), **after** ($u \in C^1(\bar{\Omega})$)


3. Take test functions $v = \xi_1(x), \dots, \xi_n(x)$ in (1) (simplest case: $\phi_i = \xi_i$ = hat func) to find $\hat{u}(x) = \sum_{i=1}^n c_i \phi_i(x)$ via $n \times n$ linear system $A\mathbf{c} = \mathbf{f}$, where

$$A_{ij} = \int_{\Omega} \nabla \phi_j \cdot \nabla \xi_i \, dx - \int_{\partial\Omega} (\nabla \phi_j \cdot \mathbf{n}) \xi_i \, ds, \quad f_i = \int_{\Omega} f \xi_i \, dx$$

- When $\phi_i = \xi_i$ = hat functions, reduced regularity significant—why?
- We'll take $\xi_i = 0$ on boundary $\partial\Omega$


Linear system is sparse and positive definite

$$\text{Recall } A_{ij} = \int_{\Omega} \nabla \phi_j \cdot \nabla \xi_j \, dx$$




If support of ϕ_j , ξ_j do not overlap, $A_{ij} = 0$
 $\Rightarrow A$ highly sparse! exploit in solving $A\mathbf{c} = \mathbf{f}$

For nonzero entries, compute A_{ij} via splitting into



What is the sparsity structure of A here?



- Taking $\phi_i = \xi_i = 0$ on $\partial\Omega$, $A_{ij} = \int_{\Omega} \nabla \phi_j \cdot \nabla \xi_j \, dx$
- Then A symmetric positive definite:

$$\begin{aligned} v^T A v &= \sum_{j=1}^n \sum_{i=1}^n v_j A_{ji} v_i = \sum_{j=1}^n \sum_{i=1}^n v_j \left(\int_{\Omega} \nabla \phi_j \cdot \nabla \phi_j \, dx \right) v_i \\ &= \int_{\Omega} \left(\sum_{j=1}^n v_j \nabla \phi_j \right) \cdot \left(\sum_{i=1}^n v_i \nabla \phi_j \right) \, dx \geq 0. \end{aligned}$$

- (preconditioned) conjugate gradient applicable/effective for $A\mathbf{c} = \mathbf{f}$

- In this special case, $V = H_0^1(\Omega)$
- If $u = g$ on $\partial\Omega$ we rewrite the weak problem: find $u \in V_g$ such that

$$a(u, v) = F(v), \quad \forall v \in V_0$$

- **Trial space:** $V_g = \{w \in H^1(\Omega) : w = g \text{ on } \partial\Omega\}$, **test space:** $V_0 = H_0^1(\Omega)$
- Alternatively: **lifting strategy** (solve the homogeneous weak form for $u - u_g$ where u_g is a function st. $u_g = g$ on $\partial\Omega$)
- common notation: $(u, v) := \int_{\Omega} uv \, dx$, $\|v\|_{0,\Omega}^2 = (v, v)$
- u solution of the weak formulation need not belong to $C^2(\bar{\Omega})$, but if it does, then it is a *strong solution*

Theorem

(Lax-Milgram) Let $(V, \|\cdot\|_V)$ be a Hilbert space and V_0 a closed subspace and consider the problem: find $u \in V$ st

$$a(u, v) = F(v), \quad \forall v \in V_0.$$

Assume

- *$a(\cdot, \cdot)$ is bounded:* $|a(v, w)| \leq C_1 \|v\|_V \|w\|_V, v, w \in V$
- *$a(\cdot, \cdot)$ is V -elliptic (or coercive):* $a(v, v) \geq C_2 \|v\|_V^2, v \in V$
- *$F(\cdot)$ is bounded:* $|F(v)| \leq C_3 \|v\|_V, v \in V$

Then the problem is uniquely solvable and $\|u\|_V \leq C_2^{-1} \|F\|_{V'}$.

(But this is not a solution *method*!)

Let's check it: find $u \in H^1(\Omega)$ such that

$$u = 0 \quad \text{on } \partial\Omega, \quad \text{and } a(u, v) = F(v) \quad \forall v \in H_0^1(\Omega).$$

- $H^1(\Omega)$ with the norm $\|v\|_{1,\Omega}^2 := \|v\|_{0,\Omega}^2 + \|\nabla v\|_{0,\Omega}^2$ is a Hilbert space
- the bilinear form is **bounded** (C-S and norm def.)
- the linear functional is **bounded** (C-S and norm def.)
- the bilinear form is $H^1(\Omega)$ –elliptic (established using Poincaré ineq.)

Cauchy-Schwarz inequality: $|(v, w)| \leq \|v\| \|w\|$, for $v, w \in V$

Poincaré inequality: $\|v\|_{0,\Omega} \leq C \|\nabla v\|_{0,\Omega}$, for $v \in H_0^1(\Omega)$

Let's now consider V_h **subspace** of V , with $\dim V_h = n < \infty$

- Replace V by V_h in the weak form. We get: find $u_h \in V_h$ (an approximation of u) st.

$$a(\mathbf{u}_h, v_h) = F(v_h) \quad \forall v_h \in V_h.$$

- Done. This was Galerkin's method
- It can be reduced to a set of n linear eqns. and n unknowns
- Comparing the "continuous" and "discrete" problems gives the **Galerkin orthogonality** ("strong" consistency)

$$a(u - u_h, v_h) = 0 \quad \forall v_h \in V_h$$

(using that a, F are unchanged and that a is linear)

$V_h \subset V \Rightarrow$ Lax-Milgram also applicable for the Galerkin problem \Rightarrow

- The solution of the Galerkin problem **exists and is unique**
- The method is **uniformly stable** wrt h since $\|u_h\|_V \leq C_2^{-1} \|F\|_{V'}$

Céa's estimate: $a(\cdot, \cdot)$ bilinear, continuous and V -elliptic. Then

$$\|u - u_h\|_V \leq C_1 C_2^{-1} \inf_{v_h \in V_h} \|u - v_h\|_V$$

Convergence:

$$\lim_{h \rightarrow 0} \|u_h - u\|_V = 0,$$

valid if V_h is chosen adequately

Theorem: $\|\nabla u - \nabla u_h\| = \min\{\|\nabla u - \nabla v_h\| : v_h = \sum_{i=1}^n c_i \phi_i\},$

where $\|\nabla u\|^2 := \int_{\Omega} (\nabla u \cdot \nabla u) dx (= a(u, u))$, energy norm

Proof:

$$\begin{aligned}\|\nabla u - \nabla u_h\|^2 &= a(u - u_h, u - u_h) = a(u - u_h, u - v_h + v_h - u_h) \\ &= a(u - u_h, u - v_h) + a(u - u_h, v_h - u_h) \\ &= a(u - u_h, u - v_h)\end{aligned}$$

due to Galerkin orthogonality, since

$a(u - u_h, v_h - u_h) = \int_{\Omega} (\nabla(u - u_h) \cdot \nabla(v_h - u_h)) dx = (r, v_h - u_h)$. By Cauchy-Schwarz,

$$a(u - u_h, u - v_h) \leq \|\nabla(u - u_h)\| \cdot \|\nabla(u - v_h)\|$$

- Let $\{\phi_j\}$ be a basis of V_h
- \Rightarrow we have only to guarantee that the Galerkin problem holds for all functions of the basis

$$a(u_h, \phi_i) = F(\phi_i), \quad i = 1, \dots, n.$$

- Since $u_h \in V_h$, then $u_h(x) = \sum_{j=1}^n c_j \phi_j(x)$, (with unknown coeffs)
- Then $\sum_{j=1}^n c_j a(\phi_j, \phi_i) = F(\phi_i), \quad i = 1, \dots, n$
- A : stiffness matrix ($a_{ij} = a(\phi_j, \phi_i)$), \mathbf{f} : load vector $f_i = F(\phi_i)$
- $A\mathbf{c} = \mathbf{f}$. If associated to a coercive problem, then A is positive definite



But V_h still not revealed! (which will actually dictate the form of A)

Galerkin method II

Bases and the FEM

Let's "discretize" the remainder of the problem (spaces, weak form, domain)

- Polygonal domain $\Omega \subset \mathbb{R}^2$, partition it into triangles
- If two triangles have some intersection, it is either on common vertex or a common full edge. In particular, two different triangles do not overlap
- h : length of the longest edge of all K in the "regular mesh" \mathcal{T}_h



- \mathbb{P}_r : polynomials of degree r or less. E.g.
 $\mathbb{P}_1 = \{g(\mathbf{x}) = a + bx_1 + cx_2, \text{ with } a, b, c \in \mathbb{R}\}$
- $\dim \mathbb{P}_r = (r+1)(r+2)/2$
- On each $K \in \mathcal{T}_h$, v_h is well-defined knowing its value in $\dim \mathbb{P}_r$ points

Finite element space

$$X_h^r = \{v_h \in C^0(\bar{\Omega}) : v_h|_K \in \mathbb{P}_r, \forall K \in \mathcal{T}_h\}$$

and the one accounting for the BC

$$\overset{\circ}{X}_h^r = \{v_h \in X_h^r : v_h|_{\partial\Omega} = 0\}$$

Lemma


If $v \in C^0(\bar{\Omega})$ and $v \in H^1(K)$ for all $K \in \mathcal{T}_h$, then $v \in H^1(\Omega)$.

For our Poisson problem (with the given BC) we set $V_h = \overset{\circ}{X}_h^r$

OK. V_h more or less clear, but what about $\{\phi_j\}$?

Since (in this particular case) $V_h = \overset{\circ}{X}_h^r$, each v_h is characterized by values in the "nodes" \mathbf{N}_j , $i = 1, \dots, n$. Thus, a basis **can** be

$$\phi_j(\mathbf{N}_i) = \delta_{ij} = \begin{cases} 0 & i \neq j, \\ 1 & i = j \end{cases}$$



If $r = 1$, the nodes coincide with the triangle vertices (in the interior). [a.k.a. Lagrangian Finite Elements]

- $v_h \in V_h$ is then a linear combination of ϕ_i 's:

$$v_h(x) = \sum_{i=1}^n v_i \phi_i(x) \quad \forall x \in \Omega,$$

- v_i **can be** evaluations at the nodes $v_i = v_h(\mathbf{N}_i)$

- Back to Poisson

$$\int_{\Omega} \nabla u_h \cdot \nabla v_h \, dx = \int_{\Omega} f v_h \, dx \quad \forall v_h \in V_h$$



- Expanding also the discrete solution, the Galerkin method gives

$$\sum_{j=1}^n u_j \int_{\Omega} \nabla \phi_j \cdot \nabla \phi_i \, dx = \int_{\Omega} f \phi_i \, dx, \quad i = 1, \dots, n$$



- Stiffness matrix ($n \times n$) A with $a_{ij} = \int_{\Omega} \nabla \phi_j \cdot \nabla \phi_i \, dx$

- $A\mathbf{u} = \mathbf{f}$

Higher-order



3-d



Inhomogeneous Dirichlet b.c.

$$-\nabla^2 u = f \quad \text{in } \Omega \subset \mathbb{R}^d, \quad u = g \text{ on } \partial\Omega.$$

- Take $u_h(x) = \sum_{j=1}^n c_j \phi_j(x) + \sum_{j=n+1}^{n+n_d} g(x_j) \phi_j(x)$
 - red term prescribed s.t. b.c. satisfied
 - e.g. $\phi_{n+\ell}(x)$ hat func at $x_{n+\ell} \in \partial\Omega$
- The rest remain same; note test space does not include $\phi_{n+\ell}$

Neumann b.c.

$$-\nabla^2 u = f \quad \text{in } \Omega \subset \mathbb{R}^d, \quad \nabla u \cdot \mathbf{n} = g \text{ on } \partial\Omega.$$

Recall weak form

$$\int_{\Omega} \nabla u \cdot \nabla v \, dx = \int_{\partial\Omega} (\nabla u \cdot \mathbf{n}) v \, ds + \int_{\Omega} f v \, dx = \int_{\partial\Omega} g v \, ds + \int_{\Omega} f v \, dx$$

- Take $u_h(x) = \sum_{j=1}^{n+n_e} c_j \phi_j(x)$ ($\phi_{n+\ell}(x)$ nonzero on $\partial\Omega$)
- test space $\xi_j = \phi_j$, $j = 1, \dots, n + n_e$
- Note $\int_{\partial\Omega} g v \, ds$ influences right-hand side in $A\mathbf{c} = \mathbf{f}$
- Robin ($u + \nabla u \cdot \mathbf{n} = g$ on $\partial\Omega$) or mixed ($u = g_1$ on $\partial\Omega$, $\nabla u \cdot \mathbf{n} = g_2$ on $\partial\Omega_2$) b.c. possible

1. Estimate the local interpolation error $v - \Pi_K^r v$, where

$$\Pi_K^r : C^0(K) \rightarrow \mathbb{P}_r(K), \quad v \mapsto \Pi_K^r v$$

2. Extension of the estimate to the whole mesh

$$|v - \Pi_K^r v|_{m,\Omega} \leq Ch^{r+1-m} |v|_{r+1,K}, \quad m = 0, 1$$

3. Error estimate in the "energy norm" (C indep. of h and u)

$$\|u - u_h\|_{1,\Omega} \leq C_1 C_2^{-1} h^r |u|_{r+1,\Omega}$$

Evidently, 2 ways of increase accuracy (reduce h or increase r). The latter effective only if u is smooth enough...

If $u \in H^{p+1}(\Omega)$ for some $p > 0$, then

$$\|u - u_h\|_{1,\Omega} \leq Ch^s |u|_{s+1,\Omega}, \quad s = \min\{r, p\}$$

Then, if e.g. $u \in H^2(\Omega)$ (i.e. $p = 1$), then going for polynomials of degree ≥ 2 **won't get you more accuracy**

Summary:

r	$u \in H^1(I)$ ($p = 0$)	$u \in H^2(I)$ ($p = 1$)	$u \in H^3(I)$ ($p = 2$)	$u \in H^4(I)$ ($p = 3$)	$u \in H^5(I)$ ($p = 4$)
1	converges	h^1	h^1	h^1	h^1
2	converges	h^1	h^2	h^2	h^2
3	converges	h^1	h^2	h^3	h^3
4	converges	h^1	h^2	h^3	h^4

Sometimes we're also interested in L^2 -norm estimates. For Poisson one can prove that if $u \in H^{p+1}(\Omega)$ for some $p > 0$, then

$$\|u - u_h\|_{0,\Omega} \leq Ch^{s+1} |u|_{s+1,\Omega}, \quad s = \min\{r, p\}$$

Generalised Stokes equations

Strong form

We study the *generalised Stokes* problem with homogeneous Dirichlet boundary conditions

$$\begin{aligned}\mathbf{u} - \nu \Delta \mathbf{u} + \nabla p &= \mathbf{f} \quad \text{in } \Omega, \\ \nabla \cdot \mathbf{u} &= 0 \quad \text{in } \Omega, \\ \mathbf{u} &= \mathbf{0} \quad \text{on } \partial\Omega,\end{aligned}$$

- \mathbf{u} vector field (in \mathbb{R}^2 or \mathbb{R}^3), p : pressure (scalar func.) the medium)
- describe the steady motion of an incompressible viscous fluid in a porous domain
- the model is valid for $Re \ll 1$

Generalised Stokes equations

Weak form

- Testing against \mathbf{v}, q , integrate over Ω , and apply IBP on the momentum equation: find $\mathbf{u} \in \mathbf{V}$ and $p \in Q_0$ (mixed FEM) st

$$\int_{\Omega} (\mathbf{u} \cdot \mathbf{v} + \nu \nabla \mathbf{u} : \nabla \mathbf{v}) - \int_{\Omega} p \nabla \cdot \mathbf{v} = \int_{\Omega} \mathbf{f} \cdot \mathbf{v} \quad \forall \mathbf{v} \in \mathbf{V},$$

$$\int_{\Omega} q \nabla \cdot \mathbf{u} = 0 \quad \forall q \in Q_0,$$

where $\mathbf{V} = [H_0^1(\Omega)]^d$ and $Q_0 = L_0^2(\Omega) = \{q \in L^2(\Omega) : q = 0 \text{ on } \partial\Omega\}$,
 $\nabla \mathbf{u} : \nabla \mathbf{v} = \nabla u_x \cdot \nabla v_x + \nabla u_y \cdot \nabla v_y$ (in 2d)

- bilinear forms $a : \mathbf{V} \times \mathbf{V} \rightarrow \mathbb{R}$ and $b : \mathbf{V} \times Q \rightarrow \mathbb{R}$, and functional $\mathcal{F}(\mathbf{v}) = \int_{\Omega} \mathbf{f} \cdot \mathbf{v}$:

$$a(\mathbf{u}, \mathbf{v}) = \int_{\Omega} (\mathbf{u} \cdot \mathbf{v} + \nu \nabla \mathbf{u} \cdot \nabla \mathbf{v}), \quad b(\mathbf{u}, q) = - \int_{\Omega} q \nabla \cdot \mathbf{u}.$$

- Find $(\mathbf{u}, p) \in \mathbf{V} \times Q_0$ such that

$$\begin{aligned} a(\mathbf{u}, \mathbf{v}) + b(\mathbf{v}, p) &= \mathcal{F}(\mathbf{v}) & \forall \mathbf{v} \in \mathbf{V}, \\ b(\mathbf{u}, q) &= 0 & \forall q \in Q_0, \end{aligned}$$

- For Stokes eqn: find $(\mathbf{u}_h, p_h) \in \mathbf{V}_h \times Q_h$ such that

$$\begin{aligned}a(\mathbf{u}_h, \mathbf{v}_h) + b(\mathbf{v}_h, p_h) &= \mathcal{F}(\mathbf{v}_h) & \forall \mathbf{v}_h \in \mathbf{V}_h, \\b(\mathbf{u}_h, q_h) &= 0 & \forall q_h \in Q_h,\end{aligned}$$

- $\{\mathbf{V}_h \subset \mathbf{V}\}$ and $\{Q_h \subset Q_0\}$ are families of **finite dimensional subspaces**

Galerkin (conforming) finite element method II

Find $\mathbf{u} \in \mathbf{V}$ and $p \in Q_0$ (**mixed FEM**) st

$$\begin{aligned}\int_{\Omega} (\mathbf{u} \cdot \mathbf{v} + \mathbf{v} \nabla \mathbf{u} : \nabla \mathbf{v}) - \int_{\Omega} p \nabla \cdot \mathbf{v} &= \int_{\Omega} \mathbf{f} \cdot \mathbf{v} \quad \forall \mathbf{v} \in \mathbf{V}, \\ \int_{\Omega} q \nabla \cdot \mathbf{u} &= 0 \quad \forall q \in Q_0,\end{aligned}$$

Associated linear system.

- $\{\varphi_j\}_{j=1}^N$ and $\{\phi_k\}_{k=1}^M$, basis functions for \mathbf{V}_h and Q_h
- $\mathbf{u}_h = \sum_{j=1}^N u_j \varphi_j(x)$, $p_h = \sum_{k=1}^M p_k \phi_k(x)$, with $N = \dim(\mathbf{V}_h)$, $M = \dim(Q_h)$
- Choosing the basis functions as tests:

$$\begin{aligned} \mathbf{A} \mathbf{U} + \mathbf{B}^T \mathbf{P} &= \mathbf{F}, \\ \mathbf{B} \mathbf{U} &= \mathbf{0}, \end{aligned} \quad \Leftrightarrow \quad \begin{pmatrix} \mathbf{A} & \mathbf{B}^T \\ \mathbf{B} & \mathbf{0} \end{pmatrix} \begin{pmatrix} \mathbf{U} \\ \mathbf{P} \end{pmatrix} = \begin{pmatrix} \mathbf{F} \\ \mathbf{0} \end{pmatrix},$$

Galerkin (conforming) finite element method III

- $\mathbf{A} \in \mathbb{R}^{N \times N}$ and $\mathbf{B} \in \mathbb{R}^{M \times N}$ are associated to $a(\cdot, \cdot)$ and $b(\cdot, \cdot)$

$$(\mathbf{A})_{ij} = a(\varphi_j, \varphi_i), \quad \mathbf{B}_{kj} = b(\varphi_j, \phi_k), \quad i, j = 1, \dots, N, \quad k = 1, \dots, M.$$

- Unknowns: $\mathbf{U} = (u_1, \dots, u_N)^T$, $\mathbf{P} = (p_1, \dots, p_M)^T$
- Datum: $\mathbf{F} = (f_1, \dots, f_N)^T$ with $f_i = \int_{\Omega} \mathbf{f} \cdot \varphi_i$
- The (generalised) Stokes matrix

$$\mathbf{S} = \begin{pmatrix} \mathbf{A} & \mathbf{B}^T \\ \mathbf{B} & \mathbf{0} \end{pmatrix} \in \mathbb{R}^{(N+M) \times (N+M)}$$

is **block-symmetric** (since \mathbf{A} is symmetric) and **indefinite** (positive and negative eigenvalues)

- A stable solver is MINRES (symmetric variant of GMRES); preconditioning of course important

CG optimality from FEM arguments

$Ax_* = b$, x_k : CG solution after k steps, i.e., $Q^T(Ax_k - b) = 0$. Then

$$x_k = \operatorname{argmin}_{x \in \operatorname{span}(Q)} \|x - x_*\|_A$$

Since $\|y\|_A^2 = (y, y)_A = y^T A y = \|A^{1/2} y\|^2$, statement equivalent to

$$\|A^{1/2}(x_k - x_*)\| = \min_x \{ \|A^{1/2}(x - x_*)\| : x = \sum_{i=1}^k y_i q_i \}.$$

FEM-type proof: (recall Poisson) for any $y \in Q$,

$$\begin{aligned} \|A^{1/2}(x_k - x_*)\|^2 &= (x_k - x_*, x_k - x_*)_A = (x_k - x_*, x_k - y + y - x_*) \\ &= (x_k - x_*, y - x_*)_A + (x_k - x_*, x_k - y)_A \\ &= (x_k - x_*, y - x_*)_A \end{aligned}$$

due to Galerkin orthogonality:

$$(x_k - x_*, x_k - y)_A = (A(x_k - x_*), x_k - y) = (Ax_k - b, x_k - y) = (r, x_k - y) = 0.$$

By Cauchy-Schwarz,

$$\|A^{1/2}(x_k - x_*)\|^2 \leq \|A^{1/2}(x_k - x_*)\| \|A^{1/2}(x_k - y)\|.$$

CG convergence

$A^T = A$ positive definite. Let $e_k := x_* - x_k$. $e_0 = x_*$, and

$$\begin{aligned}\frac{\|e_k\|_A}{\|e_0\|_A} &= \min_{x \in \mathcal{K}_{k-1}(A, b)} \|x_k - x_*\|_A / \|x_*\|_A \\ &= \min_{p_{k-1} \in \mathcal{P}_{k-1}} \|p_{k-1}(A)b - A^{-1}b\|_A / \|e_0\|_A \\ &= \min_{p_{k-1} \in \mathcal{P}_{k-1}} \|(p_{k-1}(A)A - I)e_0\|_A / \|e_0\|_A \\ &= \min_{p \in \mathcal{P}_k, p(0)=1} \|p(A)e_0\|_A / \|e_0\|_A \\ &= \min_{p \in \mathcal{P}_k, p(0)=1} \|Q \begin{bmatrix} p(\lambda_1) & & \\ & \ddots & \\ & & p(\lambda_n) \end{bmatrix} Q^T e_0\|_A / \|e_0\|_A\end{aligned}$$

$$\begin{aligned}\text{Now } \|Q \begin{bmatrix} p(\lambda_1) & & \\ & \ddots & \\ & & p(\lambda_n) \end{bmatrix} Q^T e_0\|_A^2 &= \sum_i \lambda_i p(\lambda_i)^2 (Q^T e_0)_i^2 \leq \\ \max_j p(\lambda_j)^2 \sum_i \lambda_i (Q^T e_0)_i^2 &= \max_j p(\lambda_j)^2 \|e_0\|_A^2\end{aligned}$$

$$\frac{\|e_k\|_A}{\|e_0\|_A} \leq \|Q\|_A \|Q^T\|_A \min_{p \in \mathcal{P}_k, p(0)=1} \max |p(\lambda_i)|$$

Now

$$\min_{p \in \mathcal{P}_k, p(0)=1} \max |p(\lambda_i)| \leq \left(\frac{\sqrt{\kappa_2(A)} - 1}{\sqrt{\kappa_2(A)} + 1} \right)^k$$

- note $\kappa_2(A) = \frac{\sigma_{\max}(A)}{\sigma_{\min}(A)} = \frac{\lambda_{\max}(A)}{\lambda_{\min}(A)}$
- obtained by Chebyshev polynomial on $[\lambda_{\min}(A), \lambda_{\max}(A)]$

(special case of GMRES) $A^T = A$ Recall that
 $x \in \mathcal{K}_k(A, b) \Rightarrow x = p_{k-1}(A)b$. Hence MINRES solution is

$$\begin{aligned}\min_{x \in \mathcal{K}_k(A, b)} \|Ax - b\|_2 &= \min_{p_{k-1} \in \mathcal{P}_{k-1}} \|Ap_{k-1}(A)b - b\|_2 \\ &= \min_{\tilde{p} \in \mathcal{P}_k, \tilde{p}(0)=0} \|(\tilde{p}(A) - I)b\|_2 \\ &= \min_{p \in \mathcal{P}_k, p(0)=1} \|p(A)b\|_2\end{aligned}$$

A is diagonalizable $A = Q\Lambda Q^T$, so

$$\begin{aligned}\|p(A)\|_2 &= \|Qp(\Lambda)Q^T\|_2 \leq \|Q\|_2 \|Q^T\|_2 \|p(\Lambda)\|_2 \\ &= \max_{z \in \lambda(A)} |p(z)|\end{aligned}$$

Interpretation: (again) find polynomial s.t. $p(0) = 1$ and $|p(\lambda_i)|$ small

$$\frac{\|Ax - b\|_2}{\|b\|_2} \leq \min_{p \in \mathcal{P}_k, p(0)=1} \max |p(\lambda_i)|$$

Now

$$\min_{p \in \mathcal{P}_k, p(0)=1} \max |p(\lambda_i)| \leq \left(2 \frac{\kappa_2(A) - 1}{\kappa_2(A) + 1} \right)^{k/2}$$

- minimization needed on positive **and** negative sides, hence slower convergence when A indefinite (same bound as CG when $A \succ 0$)
- obtained by Chebyshev+change of variables [A. Greenbaum's book]

Steady-state Navier-Stokes equation

$$-\nu \nabla^2 \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p = \mathbf{f}$$

$$\nabla \cdot \mathbf{u} = 0$$

- **Nonlinear** in \mathbf{u} : iterative solution of linearized problems necessary (Picard, Newton)
- Multiple stable solutions can exist
- See e.g. Elman-Silvester-Wathen Ch.8

(from Ricardo)

Generalised Stokes equations

Choosing the spaces wisely, **we can eliminate p**

- Subspaces of $[H^1(\Omega)]^d$:

$$\mathbf{V}_{\text{div}} = \{\mathbf{v} \in [H^1(\Omega)]^d : \nabla \cdot \mathbf{v} = 0 \text{ in } \Omega\}, \quad \mathbf{V}_{\text{div}}^0 = \{\mathbf{v} \in \mathbf{V}_{\text{div}} : \mathbf{v} = \mathbf{0} \text{ on } \Gamma_D\}.$$

- Take $\mathbf{v} \in \mathbf{V}_{\text{div}}$ in the momentum equation and the term involving the pressure p vanishes
- Equation only for the velocity:

$$\text{find } \mathbf{u} \in \mathbf{V}_{\text{div}}^0 : \quad a(\mathbf{u}, \mathbf{v}) = \int_{\Omega} \mathbf{f} \cdot \mathbf{v} \quad \forall \mathbf{v} \in \mathbf{V}_{\text{div}}^0.$$

- Well-posedness via **Lax & Milgram**
- Result: if we can solve the reduced problem in \mathbf{u} , then there exists a unique p st (\mathbf{u}, p) is solution of the complete problem
- But! not practical since it requires to construct a FE space $\mathbf{V}_{\text{div},h}$ of **divergence-free functions** (up to date, only 1 paper on that)
- Plus, how do I **compute p** ?

- Conditions for well-posedness:

Abstract theory of saddle-point problems by Brezzi (1974)

Theorem: Let $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ be Hilbert spaces. Consider $\mathcal{A}(\cdot, \cdot) : X \times X \rightarrow \mathbb{R}$, $\mathcal{B}(\cdot, \cdot) : X \times Y \rightarrow \mathbb{R}$, $\ell \in X'$, $\sigma \in Y'$, and the **saddle-point** problem: find $(u, \eta) \in X \times Y$ such that

$$\mathcal{A}(u, v) + \mathcal{B}(v, \eta) = x' \langle \ell, v \rangle_X \quad \forall v \in X, \quad (2)$$

$$\mathcal{B}(u, \mu) = y' \langle \sigma, \mu \rangle_Y \quad \forall \mu \in Y. \quad (3)$$

If the following hypotheses are satisfied:

1. $\mathcal{A}(\cdot, \cdot)$ is **continuous**: $|\mathcal{A}(u, v)| \leq \gamma \|u\|_X \|v\|_X \quad \forall u, v \in X$
2. \mathcal{A} is **X^0 -elliptic**, with $X^0 = \{v \in X : \mathcal{B}(v, \mu) = 0 \forall \mu \in Y\}$,

$$|\mathcal{A}(v, v)| \geq \|v\|_X^2 \quad \forall v \in X^0;$$

3. $\mathcal{B}(\cdot, \cdot)$ is **continuous**: $|\mathcal{B}(u, \mu)| \leq \delta \|u\|_X \|\mu\|_Y \quad \forall u \in X, \forall \mu \in Y$

4. **inf-sup condition**: $\exists \beta^* > 0$ st. $\inf_{\mu \in Y, \mu \neq 0} \sup_{v \in X, v \neq 0} \frac{\mathcal{B}(v, \mu)}{\|v\|_X \|\mu\|_Y} \geq \beta^*$

Then, (2)-(3) has a unique solution $(u, \eta) \in X \times Y$ and

$$\begin{aligned} \|u\|_X &\leq \left[\|\ell\|_{X'} + \frac{1+\gamma}{\beta^*} \|\sigma\|_{Y'} \right] \\ \|\eta\|_Y &\leq \frac{1}{\beta^*} \left[\left(1 + \frac{\gamma}{\bar{\alpha}}\right) \|\ell\|_{X'} + \frac{\gamma(\bar{\alpha} + \gamma)}{\bar{\alpha}\beta^*} \|\sigma\|_{Y'} \right]. \end{aligned}$$

The Stokes equation falls in this framework with $X = \mathbf{V}$, $X^0 = \mathbf{V}_{\text{div}}$, knowing that $H_0^1(\Omega)$ and $L^2(\Omega)$ satisfy the inf-sup condition

- For the Brinkman problem: find $(\mathbf{u}_h, p_h) \in \mathbf{V}_h \times Q_h$ such that

$$\begin{aligned}a(\mathbf{u}_h, \mathbf{v}_h) + b(\mathbf{v}_h, p_h) &= \mathcal{F}(\mathbf{v}_h) & \forall \mathbf{v}_h \in \mathbf{V}_h, \\b(\mathbf{u}_h, q_h) &= 0 & \forall q_h \in Q_h,\end{aligned}$$

- $\{\mathbf{V}_h \subset \mathbf{V}\}$ and $\{Q_h \subset Q_0\}$ are families of **finite dimensional subspaces**

- Solvability also falls into the Brezzi theory with $X = \mathbf{V}_h$ and $X^0 = \mathbf{V}_h^0 = \{\mathbf{v}_h \in \mathbf{V}_h : b(\mathbf{v}_h, q_h) = 0 \forall q_h \in Q_h\}$
- $\beta^* > 0$ appearing in the inf-sup condition may depend on h !

$$\exists \beta^* > 0 : \inf_{q_h \in Q_h, q_h \neq 0} \sup_{\mathbf{v}_h \in \mathbf{V}_h, \mathbf{v}_h \neq 0} \frac{b(\mathbf{v}_h, q_h)}{\|\mathbf{v}_h\|_{H^1(\Omega)} \|q_h\|_{L^2(\Omega)}} \geq \beta^*$$

- A-priori estimates

$$\|\mathbf{u}_h\|_{\mathbf{V}} \leq \frac{1}{\bar{\alpha}} \|\mathbf{f}\|_{\mathbf{V}'}, \quad \|p_h\|_Q \leq \frac{1}{\beta} \left(1 + \frac{\gamma}{\bar{\alpha}}\right) \|\mathbf{f}\|_{\mathbf{V}'},$$

- Céa's lemma

$$\begin{aligned} \|\mathbf{u} - \mathbf{u}_h\|_{\mathbf{V}} &\leq \left(1 + \frac{\gamma}{\beta^*}\right) \left(1 + \frac{\gamma}{\bar{\alpha}}\right) \inf_{\mathbf{v}_h \in \mathbf{V}_h} \|\mathbf{u} - \mathbf{v}_h\|_{\mathbf{V}} + \frac{\delta}{\bar{\alpha}} \inf_{q_h \in Q_h} \|p - q_h\|_Q, \\ \|p - p_h\|_Q &\leq \frac{\gamma}{\beta^*} \left(1 + \frac{\gamma}{\bar{\alpha}}\right) \inf_{\mathbf{v}_h \in \mathbf{V}_h} \|\mathbf{u} - \mathbf{v}_h\|_{\mathbf{V}} + \left(1 + \frac{\delta}{\beta^*} + \frac{\delta\gamma}{\bar{\alpha}\beta^*}\right) \inf_{q_h \in Q_h} \|p - q_h\|_Q. \end{aligned}$$

Associated linear system.

- $\{\varphi_j\}_{j=1}^N$ and $\{\phi_k\}_{k=1}^M$, basis functions for \mathbf{V}_h and Q_h
- $\mathbf{u}_h = \sum_{j=1}^N u_j \varphi_j(x)$, $p_h = \sum_{k=1}^M p_k \phi_k(x)$, with $N = \dim(\mathbf{V}_h)$, $M = \dim(Q_h)$
- Choosing the basis functions as tests:

$$\begin{aligned} \mathbf{A}\mathbf{U} + \mathbf{B}^T \mathbf{P} &= \mathbf{F}, \\ \mathbf{B}\mathbf{U} &= \mathbf{0}, \end{aligned} \quad \Leftrightarrow \quad \begin{pmatrix} \mathbf{A} & \mathbf{B}^T \\ \mathbf{B} & \mathbf{0} \end{pmatrix} \begin{pmatrix} \mathbf{U} \\ \mathbf{P} \end{pmatrix} = \begin{pmatrix} \mathbf{F} \\ \mathbf{0} \end{pmatrix},$$

- $\mathbf{A} \in \mathbb{R}^{N \times N}$ and $\mathbf{B} \in \mathbb{R}^{M \times N}$ are associated to $a(\cdot, \cdot)$ and $b(\cdot, \cdot)$

$$(\mathbf{A})_{ij} = a(\varphi_j, \varphi_i), \quad \mathbf{B}_{kj} = b(\varphi_j, \phi_k), \quad i, j = 1, \dots, N, \quad k = 1, \dots, M.$$

- Unknowns: $\mathbf{U} = (u_1, \dots, u_N)^T$, $\mathbf{P} = (p_1, \dots, p_M)^T$
- Datum: $\mathbf{F} = (f_1, \dots, f_N)^T$ with $f_i = \int_{\Omega} \mathbf{f} \cdot \varphi_i$
- The (generalised) Stokes matrix

$$\mathbf{S} = \begin{pmatrix} \mathbf{A} & \mathbf{B}^T \\ \mathbf{B} & \mathbf{0} \end{pmatrix} \in \mathbb{R}^{(N+M) \times (N+M)}$$

is **block-symmetric** (since \mathbf{A} is symmetric) and **non-definite** (real eigenvalues of variable sign)

- The algebraic problem has a unique solution iff $\det(S) \neq 0$ (true if the discrete inf-sup condition holds)
- If the inf-sup condition is not satisfied


$$\exists q_h^* \in Q_h : \quad b(\mathbf{v}_h, q_h^*) = 0 \quad \forall \mathbf{v}_h \in \mathbf{V}_h.$$

- Thus, if (\mathbf{u}_h, p_h) is a solution, then also $(\mathbf{u}_h, p_h + q_h^*)$, because
$$a(\mathbf{u}_h, \mathbf{v}_h) + b(\mathbf{v}_h, p_h + q_h^*) = a(\mathbf{u}_h, \mathbf{v}_h) + b(\mathbf{v}_h, p_h) + \underbrace{b(\mathbf{v}_h, q_h^*)}_{=0} = \mathcal{F}(\mathbf{v}_h), \quad \forall \mathbf{v}_h \in \mathbf{V}_h.$$
- Non-uniqueness!!

- p_h^* breaking the inf-sup condition are called **spurious pressure modes**
- Who's fault is this?!! Q_h **and** V_h ...
- Pairs (V_h, Q_h) violating the inf-sup condition are called **inf-sup unstable**
- The weak form does not require the pressure to be continuous
- Possible choices (degrees of freedom of the velocity “•” and those of the pressure are “□”)
- See a list in Girault-Raviart or Brezzi-Fortin books


More on the discrete inf-sup condition III

Elements with Discontinuous Pressure



More on the discrete inf-sup condition IV

Elements with Continuous Pressure



- Hope in the horizon: you can still use unstable pairs (why would you want to do that?)
- Some remedies available (cf Exercises of week 4)
- General stabilisation technique: find $\mathbf{u}_h \in \mathbf{V}_h$, $q_h \in Q_h$ such that

$$\begin{aligned}a(\mathbf{u}_h, \mathbf{v}_h) + b(\mathbf{v}_h, p_h) &= \mathcal{F}(\mathbf{v}_h) - \Psi_h^{(\rho)}(\mathbf{v}_h) & \forall \mathbf{v}_h \in \mathbf{V}_h \\ b(\mathbf{u}_h, q_h) &= \Phi_h(q_h) & \forall q_h \in Q_h,\end{aligned}$$

where

$$\Psi_h^{(\rho)}(\mathbf{v}_h) = \bar{\delta} \sum_{K \in \mathcal{T}_h} h_K^2 \int_K (\alpha \mathbf{u}_h - \mathbf{v} \Delta \mathbf{u}_h + \nabla p_h - \mathbf{f}) \cdot (\rho \alpha \mathbf{v}_h - \rho \mathbf{v} \Delta \mathbf{v}_h)$$

$$\Phi_h(q_h) = \bar{\delta} \sum_{K \in \mathcal{T}_h} h_K^2 \int_K (\alpha \mathbf{u}_h - \mathbf{v} \Delta \mathbf{u}_h + \nabla p_h - \mathbf{f}) \cdot \nabla q_h.$$

with $\bar{\delta} > 0, \rho$ stabilisation parameters to be set

- $\rho = 0 \Rightarrow \psi_h^{(0)} = 0 \leftrightarrow$ Streamline Upwind/Petrov-Galerkin (SUPG) method
- $\rho = -1 \leftrightarrow$ Galerkin/Least-Squares (GLS or GaLS) method
- These methods are **strongly consistent** (other versions may not)

Stokes flow ($\alpha = 0$).

- Notice that if using $\mathbb{P}_1 - \mathbb{P}_1$ elements, then $\Delta \mathbf{v}_h = \Delta \mathbf{u}_h = \mathbf{0}$ for all $K \in \mathcal{T}_h$
- The stabilised method is well-posed for adequate stabilisation parameters (see e.g. Quarteroni-Valli, section 9.4)
- Stability and convergence also follow

- Matrix form

$$\begin{pmatrix} \mathbf{A} & \mathbf{B}^T \\ \mathbf{B} & -\mathbf{C} \end{pmatrix} \begin{pmatrix} \mathbf{U} \\ \mathbf{P} \end{pmatrix} = \begin{pmatrix} \mathbf{F} \\ \mathbf{G} \end{pmatrix}$$

$$\text{with } \mathbf{C}_{km} = \bar{\delta} \sum_{K \in \mathcal{T}_h} h_K^2 \int_K \nabla \phi_m \cdot \nabla \phi_k, \quad k, m = 1, \dots, M$$

$$\mathbf{g}_k = -\bar{\delta} \sum_{K \in \mathcal{T}_h} h_K^2 \int_K \mathbf{f} \cdot \nabla \phi_k, \quad k = 1, \dots, M.$$

- Similar method (also with a “name”): **Brezzi-Pitkaranta** (uses $\mathbb{P}_1 - \mathbb{P}_1$)

$$\begin{aligned} a_0(\mathbf{u}_h, \mathbf{v}_h) + b(\mathbf{v}_h, p_h) &= \mathcal{F}(\mathbf{v}_h) & \forall \mathbf{v}_h \in \mathbf{V}_h \\ b(\mathbf{u}_h, q_h) &= \sum_{K \in \mathcal{T}_h} \delta_K (\nabla p_h, \nabla q_h)_{0,K} & \forall q_h \in Q_h, \end{aligned}$$

$$\text{with } \delta_K = \frac{|K|^2}{5(c_1^2 + c_2^2 + c_3^2)}, \quad |K|: \text{ area of } K, \quad c_i: \text{ length of edges}$$