

# The Effects of a Primordial Black Hole on the Surface of a Neutron Star

Brady Metherall

April 13, 2017

# Black Holes

Primordial  
Black Holes

Brady  
Metherall

Introduction

Model

Analytic  
Solution

Simulation  
Solution

- A black hole is a region of space where enough mass is concentrated such that nothing, including light, can escape its gravitational pull.
- Typical black holes form from the collapse of a massive star, and have a minimum mass on the order of a few solar masses.
- In the early universe while it was very hot and dense, small perturbations could be enough to create small black holes called primordial black holes.
- The mass of a primordial black hole is probably much less than one solar mass ( $10^{-18} - 10^{-6} M_{\odot}$ ), and microscopic in size.

# Primordial Black Holes as Dark Matter

Primordial  
Black Holes

Brady  
Metherall

Introduction

Model

Analytic  
Solution

Simulation  
Solution

Dark matter is matter that does not emit/absorb light, and is chargeless. Dark matter accounts for about a quarter of the energy in the universe, but we don't know what it is.

Primordial black holes may be a good candidate for dark matter since: they are chargeless, do not emit light, have a very small radius, and are non-relativistic. Also, most other proposed explanations of dark matter involve creating a new particle not part of the Standard Model.

# Flat Star Model

Primordial  
Black Holes

Brady  
Metherall

Introduction

Model

Analytic  
Solution

Simulation  
Solution

Investigate the collision of a primordial black hole with a neutron star with the following assumptions:

- Neutron stars are flat and infinite
- Primordial black holes are point masses
- Neutron stars are incompressible fluids
- Gravitational interactions are Newtonian
- Primordial black hole has a constant velocity

# Eigenfunctions of Laplacian

Primordial  
Black Holes

Brady  
Metherall

Introduction

Model

Analytic  
Solution

Velocity  
Potential

Deformation of  
Surface  
Energy

Simulation  
Solution

To find the velocity potential we need the eigenfunctions of the Laplacian. First, assume a product solution and solve the Laplacian.

$$\varphi = R(r)Z(z)\Theta(\theta)T(t)$$

$$\nabla^2 \varphi = 0$$

$$\Rightarrow \varphi \propto \left\{ \begin{matrix} J_\mu(kr) \\ Y_\mu(kr) \end{matrix} \right\} \left\{ \begin{matrix} e^{-kz} \\ e^{kz} \end{matrix} \right\} \left\{ \begin{matrix} \sin(\mu\theta) \\ \cos(\mu\theta) \end{matrix} \right\} T(t)$$

$$\Rightarrow \varphi \propto J_0(kr)e^{kz}T(t)$$

$T(t)$  comes from boundary conditions.

# Hankel Transform

Primordial  
Black Holes

Brady  
Metherall

Introduction

Model

Analytic  
Solution

Velocity  
Potential

Deformation of  
Surface

Energy

Simulation  
Solution

The Hankel transform is needed to solve the time component of the velocity potential, and the calculation of the energy.

## Definition

The Hankel transform of a function  $f(s)$  is given by

$$(\mathcal{H}_\nu f)(\sigma) = \int_0^\infty f(s) J_\nu(s\sigma) s ds,$$

where  $J_\nu$  is the Bessel function of the first kind, of order  $\nu \geq -\frac{1}{2}$ , and  $\sigma$  is a non-negative real variable.

## Corollary

*The Hankel transform is self-reciprocal, that is, it is its own inverse.*

# Solving for $\varphi$

Primordial  
Black Holes

Brady  
Metherall

Introduction

Model

Analytic  
Solution

Velocity  
Potential  
Deformation of  
Surface  
Energy

Simulation  
Solution

From the pressure condition

$$\left( \frac{\partial^2 \varphi}{\partial t^2} + g \frac{\partial \varphi}{\partial z} + \frac{\partial \Phi}{\partial t} \right) \Big|_{z=0} = 0.$$

We can write the potential as a Hankel transform

$$\frac{\partial \Phi}{\partial t} \Big|_{z=0} = \int_0^\infty \left( \mathcal{H} \frac{\partial \Phi}{\partial t} \Big|_{z=0} \right) (k) J_0(kr) k \, dk.$$

Then,

$$\varphi = \frac{Gmv}{g} \int_0^\infty \frac{J_0(kr) e^{kz}}{1 + kv^2/g} \left( -\operatorname{sgn}(t) e^{-kv|t|} + 2 H(t) \cos(\omega_k t) \right) dk,$$

with  $\omega_k^2 = gk$

# Deformation of Surface

Primordial  
Black Holes

Brady  
Metherall

Introduction

Model

Analytic  
Solution

Velocity  
Potential

Deformation of  
Surface

Energy

Simulation  
Solution

We can easily solve for the shape of the surface now that we have the velocity potential, using

$$\left. \frac{\partial \varphi}{\partial z} \right|_{z=0} = \frac{\partial \eta}{\partial t}.$$
$$\eta = \frac{Gm}{g} \int_0^\infty \frac{J_0(kr)}{1 + kv^2/g} \left( e^{-kv|t|} + 2H(t)v\sqrt{\frac{k}{g}} \sin(\omega_k t) \right) dk$$



# Analytic Surface Waves

Primordial  
Black Holes

Brady  
Metherall

Introduction

Model

Analytic  
Solution

Velocity  
Potential

Deformation of  
Surface

Energy

Simulation  
Solution

Figure: Waves!

# Energy Transferred

Primordial  
Black Holes

Brady  
Metherall

Introduction

Model

Analytic  
Solution

Velocity  
Potential

Deformation of  
Surface

Energy

Simulation  
Solution

We can calculate the energy by taking the sum of the kinetic, and potential energies,

$$\begin{aligned} E(t) &= \frac{1}{2}\rho \int_{-\infty}^0 \int_0^{\infty} |\nabla\varphi|^2 r dr dz \int_0^{2\pi} d\theta \\ &\quad + \rho g \int_0^{\infty} \int_0^{\eta} z dz r dr \int_0^{2\pi} d\theta \\ \lim_{t \rightarrow \infty} E(t) &= 4\pi\rho \frac{G^2 m^2}{g}. \end{aligned}$$

We take the limit as  $t$  approaches infinity, because we are interested in the total energy transferred.

# Analytic Energy

Primordial  
Black Holes

Brady  
Metherall

Introduction


Model

Analytic  
Solution

Velocity  
Potential  
Deformation of  
Surface

Energy

Simulation  
Solution



# Analytic Surface Waves

Primordial  
Black Holes

Brady  
Metherall

Introduction

Model

Analytic  
Solution

Simulation  
Solution

Smooth Particle  
Hydrodynamics

Figure: Waves!

Primordial  
Black Holes

Brady  
Metherall


Introduction

Model

Analytic  
Solution

Simulation  
Solution

Smooth Particle  
Hydrodynamics



Primordial  
Black Holes

Brady  
Metherall

Introduction

Model

Analytic  
Solution

Simulation  
Solution

Smooth Particle  
Hydrodynamics

