

The Effects of a Primordial Black Hole on the Surface of a Neutron Star

Brady Metherall

April 13, 2017

Primordial Black Holes

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Simulation
Solution

- Normal black holes have a minimum mass on the order of a few solar masses.
- In the early universe while it was very hot and dense, small perturbations could be enough to create a small black hole.
- The mass of a primordial black hole is much less than one solar mass ($10^{-18} - 10^{-6} M_{\odot}$), and microscopic in size.

Primordial Black Holes as Dark Matter

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Simulation
Solution

Dark matter is matter that does not emit/absorb light, and is chargeless. Dark matter accounts for about a quarter of the energy in the universe, but we don't know what it is.

Primordial black holes may be a good candidate for dark matter since: they are chargeless, do not emit light, have a very small radius, and are non-relativistic. Also, most other proposed explanations of dark matter involve creating a new particle not part of the standard model.

Flat Star Model

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Simulation
Solution

- Neutron stars are flat and infinite
- Primordial black holes are point masses
- Neutron stars are incompressible fluids
- Gravitational interactions are Newtonian
- Constant velocity

Eigenfunctions of Laplacian

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Solving for φ
Solving for η
Plotting η
Solving for E

Simulation
Solution

Assume a product solution for the velocity potential and solve the Laplacian.

$$\varphi = R(r)Z(z)\Theta(\theta)T(t)$$

$$\nabla^2 \varphi = 0$$

$$\Rightarrow \varphi \propto \begin{Bmatrix} J_\mu(kr) \\ Y_\mu(kr) \end{Bmatrix} \begin{Bmatrix} e^{-kz} \\ e^{kz} \end{Bmatrix} \begin{Bmatrix} \sin(\mu\theta) \\ \cos(\mu\theta) \end{Bmatrix} T(t)$$

$$\Rightarrow \varphi \propto J_0(kr)e^{kz}T(t)$$

$T(t)$ comes from boundary conditions.

Hankel Transform

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Solving for φ
Solving for η
Plotting η
Solving for E

Simulation
Solution

Definition

The Hankel transform of a function $f(s)$ is given by

$$\mathcal{H}_\nu\{f\}(\sigma) = \int_0^\infty f(s) J_\nu(s\sigma) s \, ds,$$

where J_ν is the Bessel function of the first kind, of order $\nu \geq -\frac{1}{2}$, and σ is a non-negative real variable.

Corollary

The Hankel transform is self-reciprocal, that is, the inverse Hankel transform is also given by Definition 1.

Solving for φ

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Solving for φ
Solving for η
Plotting η
Solving for E

Simulation
Solution

From the pressure condition:

$$\left(\frac{\partial^2 \varphi}{\partial t^2} + g \frac{\partial \varphi}{\partial z} + \frac{\partial \Phi}{\partial t} \right) \Big|_{z=0} = 0$$

$$\text{where } \Phi = \frac{-Gm}{\sqrt{r^2 + (z + vt)^2}} = \int_0^\infty a(k) J_0(kr) k \, dk$$

$$\varphi = \frac{Gmv}{g} \int_0^\infty \frac{J_0(kr) e^{kz}}{1 + kv^2/g} \left(-\text{sgn}(t) e^{-kv|t|} + 2 H(t) \cos(\omega_k t) \right) dk$$

with $\omega_k^2 = gk$

Deformation of Surface

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Solving for φ

Solving for η

Plotting η

Solving for E

Simulation
Solution

We can easily solve for the shape of the surface now that we have the velocity potential,

$$\left. \frac{\partial \varphi}{\partial z} \right|_{z=0} = \frac{\partial \eta}{\partial t}$$
$$\eta = \frac{Gm}{g} \int_0^\infty \frac{J_0(kr)}{1 + kv^2/g} \left(e^{-kv|t|} + 2H(t)v\sqrt{\frac{k}{g}} \sin(\omega_k t) \right) dk.$$

Maple was used to perform numerical integration and plot the surface waves.

Analytic Surface Waves

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Solving for φ

Solving for η

Plotting η

Solving for E

Simulation
Solution

Figure: Waves!

Energy Transferred

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Solving for φ

Solving for η

Plotting η

Solving for E

Simulation

Solution

We can calculate the energy by taking the sum of the kinetic, and potential energies,

$$\begin{aligned} E &= \lim_{t \rightarrow \infty} \frac{1}{2} \rho \int_{-\infty}^0 \int_0^{\infty} |\nabla \varphi|^2 r dr dz \int_0^{2\pi} d\theta \\ &\quad + \rho g \int_0^{\infty} \int_0^{\eta} z dz r dr \int_0^{2\pi} d\theta \\ &= 4\pi \rho \frac{G^2 m^2}{g}. \end{aligned}$$

We take the limit as t approaches infinity, because we are interested in the total energy transferred.

Future Work

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Simulation
Solution

Smooth Particle
Hydrodynamics

My focus next semester will be on simulating the collision with smooth particle hydrodynamics. Smooth particle hydrodynamics is a computational fluid dynamics method that uses N-Body methods, and is typically used for free surface simulations.