

The Effects of a Primordial Black Hole on the Surface of a Neutron Star

Brady Metherall

April 13, 2017

Black Holes

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Simulation
Solution

Conclusion

- A black hole is a region of space where enough mass is concentrated such that nothing, including light, can escape its gravitational pull.
- Typical black holes form from the collapse of a massive star, and have a minimum mass on the order of a few solar masses.
- In the early universe, while it was very hot and dense, small perturbations could be enough to create small black holes called primordial black holes.
- The mass of a primordial black hole is probably much less than one solar mass ($10^{-18} - 10^{-6} M_{\odot}$), and microscopic in size.

Primordial Black Holes as Dark Matter

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Simulation
Solution

Conclusion

Dark matter is matter that does not emit/absorb light, and is chargeless. Dark matter accounts for about a quarter of the energy in the universe, but we don't know what it is.

Primordial black holes may be a good candidate for dark matter since: they are chargeless, do not emit light, have a very small radius, and are non-relativistic. Also, most other proposed explanations of dark matter involve creating a new particle not part of the Standard Model.

Flat Star Model

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Simulation
Solution

Conclusion

Investigate the collision of a primordial black hole with a neutron star with the following assumptions:

- Neutron stars are flat and infinite
- Primordial black holes are point masses
- Neutron stars are incompressible fluids
- Gravitational interactions are Newtonian
- Primordial black hole has a constant velocity

Eigenfunctions of Laplacian

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Velocity
Potential

Deformation of
Surface
Energy

Simulation
Solution

Conclusion

To find the velocity potential we need the eigenfunctions of the Laplacian. First, assume a product solution and solve the Laplacian.

$$\varphi = R(r)Z(z)\Theta(\theta)T(t)$$

$$\nabla^2 \varphi = 0$$

$$\Rightarrow \varphi \propto \begin{Bmatrix} J_\mu(kr) \\ Y_\mu(kr) \end{Bmatrix} \begin{Bmatrix} e^{-kz} \\ e^{kz} \end{Bmatrix} \begin{Bmatrix} \sin(\mu\theta) \\ \cos(\mu\theta) \end{Bmatrix} T(t)$$

$$\Rightarrow \varphi \propto J_0(kr)e^{kz}T(t)$$

$T(t)$ comes from boundary conditions.

Hankel Transform

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Velocity
Potential
Deformation of
Surface
Energy

Simulation
Solution

Conclusion

The Hankel transform is needed to solve the time component of the velocity potential, and the calculation of the energy.

Definition

The Hankel transform of a function $f(s)$ is given by

$$(\mathcal{H}_\nu f)(\sigma) = \int_0^\infty f(s) J_\nu(s\sigma) s ds,$$

where J_ν is the Bessel function of the first kind, of order $\nu \geq -\frac{1}{2}$, and σ is a non-negative real variable.

Corollary

The Hankel transform is self-reciprocal, that is, it is its own inverse.

Solving for φ

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Velocity
Potential
Deformation of
Surface
Energy

Simulation
Solution

Conclusion

From the pressure condition

$$\left(\frac{\partial^2 \varphi}{\partial t^2} + g \frac{\partial \varphi}{\partial z} + \frac{\partial \Phi}{\partial t} \right) \Big|_{z=0} = 0.$$

We can write the potential as a Hankel transform

$$\frac{\partial \Phi}{\partial t} \Big|_{z=0} = \int_0^\infty \left(\mathcal{H} \frac{\partial \Phi}{\partial t} \Big|_{z=0} \right) (k) J_0(kr) k \, dk.$$

Then,

$$\varphi = \frac{Gmv}{g} \int_0^\infty \frac{J_0(kr) e^{kz}}{1 + kv^2/g} \left(-\operatorname{sgn}(t) e^{-kv|t|} + 2 H(t) \cos(\omega_k t) \right) dk,$$

with $\omega_k^2 = gk$.

Deformation of Surface

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Velocity
Potential

Deformation of
Surface

Energy

Simulation
Solution

Conclusion

We can easily solve for the shape of the surface now that we have the velocity potential, using

$$\left. \frac{\partial \varphi}{\partial z} \right|_{z=0} = \frac{\partial \eta}{\partial t}.$$
$$\eta = \frac{Gm}{g} \int_0^\infty \frac{J_0(kr)}{1 + kv^2/g} \left(e^{-kv|t|} + 2H(t)v\sqrt{\frac{k}{g}} \sin(\omega_k t) \right) dk.$$

Analytic Surface Waves

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Velocity
Potential

Deformation of
Surface

Energy

Simulation
Solution

Conclusion

Figure: Analytic waves

Energy Transferred

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Velocity
Potential

Deformation of
Surface

Energy

Simulation
Solution

Conclusion

We can calculate the energy by taking the sum of the kinetic, and potential energies,

$$\begin{aligned} E(t) &= \frac{1}{2}\rho \int_{-\infty}^0 \int_0^{\infty} |\nabla\varphi|^2 r dr dz \int_0^{2\pi} d\theta \\ &\quad + \rho g \int_0^{\infty} \int_0^{\eta} z dz r dr \int_0^{2\pi} d\theta \\ \lim_{t \rightarrow \infty} E(t) &= 4\pi\rho \frac{G^2 m^2}{g}. \end{aligned}$$

We take the limit as t approaches infinity because we are interested in the total energy transferred.

Analytic Energy

Primordial
Black Holes

Brady
Metherall

Introduction

Model


Analytic
Solution

Velocity
Potential
Deformation of
Surface

Energy

Simulation
Solution

Conclusion



Smoothed Particle Hydrodynamics

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Simulation
Solution

Smoothed
Particle
Hydrodynamics
Energy

Conclusion

In order to simulate the collision smoothed particle hydrodynamics (SPH) was used.

- Developed in 1977 for astrophysics simulations
- Mesh-free, N -body methods
- Uses nearest neighbours

The neutron star is treated as a 120×15 tank with a particle spacing of 0.075. The simulation was run in parallel over three days with PySPH, a SPH framework.

Analytic Surface Waves

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Simulation
Solution

Smoothed
Particle
Hydrodynamics
Energy

Conclusion

Figure: Simulation waves

Energy Calculation

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Simulation
Solution

Smoothed
Particle
Hydrodynamics

Energy

Conclusion

The energy can be calculated in the same fashion as before:

$$E = \sum_i \frac{1}{2} m_i (u_i^2 + v_i^2) + m_i g y_i.$$

However, this must be transformed to three dimensions:

$$E = \frac{\pi}{dx} \sum_i \left(\frac{1}{2} (u_i^2 + v_i^2) + g y_i \right) m_i |x_i|.$$

Raw Noisy Energy

Primordial
Black Holes

Brady
Metherall

Introduction


Model

Analytic
Solution

Simulation
Solution

Smoothed
Particle
Hydrodynamics
Energy

Conclusion



Cleaned Energy

Primordial
Black Holes

Brady
Metherall

Introduction


Model

Analytic
Solution

Simulation
Solution

Smoothed
Particle
Hydrodynamics
Energy

Conclusion



Conclusion

Primordial
Black Holes

Brady
Metherall

Introduction

Model

Analytic
Solution

Simulation
Solution

Conclusion

If this project were continued some modifications would be

- Better initial conditions for the simulation
- Use a spherical model instead of planar
- Use general relativity instead of Newtonian gravity