

Collisions of Primordial Black Holes and Neutron Stars

Brady Metherall 100516905

Monday November 28, 2016

Primordial Black Holes

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Future Work

- Normal black holes have a minimum mass on the order of a few solar masses.
- In the early universe while it was very hot and dense, small perturbations could be enough to create a small black hole.
- The mass of a primordial black hole is much less than one solar mass ($10^{-18} - 10^{-6} M_{\odot}$), and microscopic in size.

Primordial Black Holes as Dark Matter

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Future Work

Dark matter is matter that does not emit/absorb light, and is chargeless. Dark matter accounts for about a quarter of the matter in the universe, but we don't know what it is.

Primordial black holes may be a good candidate for dark matter since: they are chargeless, do not emit light, have a very small radius, and are non-relativistic. Also, most other proposed explanations of dark matter involve creating a new particle not part of the standard model.

Flat Star Model

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ
Solving for η
Plotting η
Solving for E

Future Work

- Neutron stars are flat and infinite
- Primordial black holes are point masses
- Neutron stars are incompressible fluids
- Gravitational interactions are Newtonian
- Constant velocity

Eigenfunctions of Laplacian

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ

Solving for η

Plotting η

Solving for E

Future Work

Assume a product solution for the velocity potential and solve the Laplacian.

$$\varphi = R(r)Z(z)\Theta(\theta)T(t)$$

$$\nabla^2 \varphi = 0$$

$$\Rightarrow \varphi \propto \begin{Bmatrix} J_\mu(kr) \\ Y_\mu(kr) \end{Bmatrix} \begin{Bmatrix} e^{-kz} \\ e^{kz} \end{Bmatrix} \begin{Bmatrix} \sin(\mu\theta) \\ \cos(\mu\theta) \end{Bmatrix} T(t)$$

$$\Rightarrow \varphi \propto J_0(kr)e^{kz}T(t)$$

$T(t)$ comes from boundary conditions.

Hankel Transform

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ
Solving for η
Plotting η
Solving for E

Future Work

Definition

The Hankel transform of a function $f(s)$ is given by

$$\mathcal{H}_\nu\{f\}(\sigma) = \int_0^\infty f(s) J_\nu(s\sigma) s \, ds,$$

where J_ν is the Bessel function of the first kind, of order $\nu \geq -\frac{1}{2}$, and σ is a non-negative real variable.

Corollary

The Hankel transform is self-reciprocal, that is, the inverse Hankel transform is also given by Definition 1.

Solving for φ

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ

Solving for η

Plotting η

Solving for E

Future Work

From the pressure condition:

$$\left(\frac{\partial^2 \varphi}{\partial t^2} + g \frac{\partial \varphi}{\partial z} + \frac{\partial \Phi}{\partial t} \right) \Big|_{z=0} = 0$$

$$\text{where } \Phi = \frac{-Gm}{\sqrt{r^2 + (z + vt)^2}} = \int_0^\infty a(k) J_0(kr) k dk$$

$$\varphi = \frac{Gmv}{g} \int_0^\infty \frac{J_0(kr) e^{kz}}{1 + kv^2/g} \left(-\text{sgn}(t) e^{-kv|t|} + 2 H(t) \cos(\omega_k t) \right) dk$$
$$\text{with } \omega_k^2 = gk$$

Deformation of Surface

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ
Solving for η
Plotting η
Solving for E

Future Work

We can easily solve for the shape of the surface now that we have the velocity potential,

$$\left. \frac{\partial \varphi}{\partial z} \right|_{z=0} = \frac{\partial \eta}{\partial t}$$
$$\eta = \frac{Gm}{g} \int_0^\infty \frac{J_0(kr)}{1 + kv^2/g} \left(e^{-kv|t|} + 2H(t)v\sqrt{\frac{k}{g}} \sin(\omega_k t) \right) dk.$$

Maple was used to perform numerical integration and plot the surface waves.

Plotting η

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ

Solving for η

Plotting η

Solving for E

Future Work




Figure : $t = -1$ s.

Plotting η

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ

Solving for η

Plotting η

Solving for E

Future Work




Figure : $t = 0$ s.

Plotting η

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ

Solving for η

Plotting η

Solving for E

Future Work




Figure : $t = 1$ s.

Plotting η

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ

Solving for η

Plotting η

Solving for E

Future Work




Figure : $t = 2$ s.

Plotting η

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ

Solving for η

Plotting η

Solving for E

Future Work




Figure : $t = 3$ s.

Plotting η

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ

Solving for η

Plotting η

Solving for E

Future Work




Figure : $t = 4$ s.

Energy Transferred

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ
Solving for η
Plotting η
Solving for E

Future Work

We can calculate the energy by taking the sum of the kinetic, and potential energies,

$$\begin{aligned} E &= \lim_{t \rightarrow \infty} \frac{1}{2} \rho \int_{-\infty}^0 \int_0^{\infty} |\nabla \varphi|^2 r dr dz \int_0^{2\pi} d\theta \\ &\quad + \rho g \int_0^{\infty} \int_0^{\eta} z dz r dr \int_0^{2\pi} d\theta \\ &= 4\pi \rho \frac{G^2 m^2}{g}. \end{aligned}$$

We take the limit as t approaches infinity, because we are interested in the total energy transferred.

Future Work

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Future Work

Smooth Particle
Hydrodynamics

My focus next semester will be on simulating the collision with smooth particle hydrodynamics. Smooth particle hydrodynamics is a computational fluid dynamics method that uses N-Body methods, and is typically used for free surface simulations.