

Collisions of Primordial Black Holes and Neutron Stars

Brady Metherall 100516905

Monday November 28, 2016

Introduction

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Numerical
Approach

A primordial black hole is a black hole that was hypothetically created in the early universe.

Flat Star Model

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ
Solving for η
Plotting η
Solving for E

Numerical
Approach

- Neutron stars are flat and infinite
- Primordial black holes are point masses
- Neutron stars are incompressible fluids
- Gravitational interactions are Newtonian
- Constant velocity

Eigenfunctions of Laplacian

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ

Solving for η

Plotting η

Solving for E

Numerical
Approach

Assume a product solution for the velocity potential and solve the Laplacian.

$$\varphi = R(r)Z(z)\Theta(\theta)T(t)$$

$$\nabla^2 \varphi = 0$$

$$\Rightarrow \varphi \propto \begin{Bmatrix} J_\mu(kr) \\ Y_\mu(kr) \end{Bmatrix} \begin{Bmatrix} e^{-kz} \\ e^{kz} \end{Bmatrix} \begin{Bmatrix} \sin(\mu\theta) \\ \cos(\mu\theta) \end{Bmatrix} T(t)$$

$$\Rightarrow \varphi \propto J_0(kr)e^{kz}T(t)$$

$T(t)$ comes from boundary conditions.

Solving for φ

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ

Solving for η

Plotting η

Solving for E

Numerical
Approach

$$\Phi = \frac{-Gm}{\sqrt{r^2 + (z + vt)^2}}$$

$$\left(\frac{\partial^2 \varphi}{\partial t^2} + g \frac{\partial \varphi}{\partial z} \right) \Big|_{z=0} = - \frac{\partial \Phi}{\partial t} \Big|_{z=0}$$

$$\varphi = \frac{Gmv}{g} \int_0^\infty \frac{J_0(kr)e^{kz}}{1 + kv^2/g} \left(-\operatorname{sgn}(t)e^{-kv|t|} + 2H(t) \cos(\omega_k t) \right) dk$$

with $\omega_k^2 = gk$

Deformation of Surface

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ
Solving for η
Plotting η
Solving for E

Numerical
Approach

$$\left. \frac{\partial \varphi}{\partial z} \right|_{z=0} = \frac{\partial \eta}{\partial t}$$

$$\eta = \frac{Gm}{g} \int_0^\infty \frac{J_0(kr)}{1 + kv^2/g} \left(e^{-kv|t|} + 2H(t)v\sqrt{\frac{k}{g}} \sin(\omega_k t) \right) dk$$

Plotting η

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ

Solving for η

Plotting η

Solving for E

Numerical
Approach




Figure : $t = -1$ s.

Plotting η

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ

Solving for η

Plotting η

Solving for E

Numerical
Approach




Figure : $t = 0$ s.

Plotting η

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ

Solving for η

Plotting η

Solving for E

Numerical
Approach




Figure : $t = 1$ s.

Plotting η

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ

Solving for η

Plotting η

Solving for E

Numerical
Approach




Figure : $t = 2$ s.

Plotting η

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ

Solving for η

Plotting η

Solving for E

Numerical
Approach




Figure : $t = 3$ s.

Plotting η

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ

Solving for η

Plotting η

Solving for E

Numerical
Approach




Figure : $t = 4$ s.

Energy Transferred

Collisions of
Primordial
Black Holes
and Neutron
Stars

Brady
Metherall

Introduction

Flat Star
Model

Solving for φ

Solving for η

Plotting η

Solving for E

Numerical
Approach

$$\begin{aligned} E &= \lim_{t \rightarrow \infty} \frac{1}{2} \rho \int_{-\infty}^0 \int_0^{\infty} |\nabla \varphi|^2 r dr dz \int_0^{2\pi} d\theta \\ &\quad + \rho g \int_0^{\infty} \int_0^{\eta} z dz r dr \int_0^{2\pi} d\theta \\ &= 4\pi \rho \frac{G^2 m^2}{g} \end{aligned}$$

Collisions of Primordial Black Holes and Neutron Stars

Brady
Metherall

Introduction

Flat Star
Model

Numerical
Approach

Smooth Particle
Hydrodynamics