

Makana Health Oncology Portfolio

Choose a product to proceed:

 **Oncura**[®]
(oncurimab) 80mg

EGFR-mutated NSCLC
(launched)

 **Immunexa**[™]
(immuneximab) 100mg

PD-1 immunotherapy
(PHASE III)

 **Makuparib**
(makulunparib) 50mg

PARP inhibitor
(PHASE II pipeline)

PIPELINE +





Asset	Mechanism	Indication	Phase
ONCURA®	EGFR inhibitor	EGFR-mutated metastatic NSCLC	LAUNCHED
IMMUNEXA™	PD-1 immunotherapy	Advanced solid tumours	PHASE III
MAKUPARIB	PARP inhibitor	DNA repair-deficient cancers	PHASE II PIPELINE

EXPLORING COMBINATION AND SEQUENTIAL TREATMENT STRATEGIES ACROSS THE PORTFOLIO





 **Oncura**[®]
(oncurimab) 80mg

EGFR-mutated NSCLC
(launched)


 **Immunexa**[™]
(immuneximab) 100mg

PD-1 immunotherapy
(PHASE III)

 **Makuparib**
(makulunparib) 50mg

PARP inhibitor
(PHASE II pipeline)

Please Select your flow to begin the presentation



MEDICAL ONCOLOGIST



SHORT CALL




ONCOLOGY NURSE



SWIPE TO THEN TAP SELECT.

ANIMATION ☐

Please Select your flow to begin the presentation



MEDICAL ONCOLOGIST



SHORT CALL




ONCOLOGY NURSE



SWIPE TO THEN TAP SELECT.

ANIMATION



TRANSFORMING LIVES WITH ONCE-WEEKLY TARGETED THERAPY

For appropriate patients with confirmed EGFR-mutated metastatic non-small cell lung cancer (NSCLC).
Please note: All patient profiles, names, and scenarios shown are fictional and are used for illustrative purposes only.
Individual patient characteristics, treatment decisions, and outcomes may vary.

US-ONCU-0541 | February 2026



1. Makana Health. Oncura (oncurimab). Summary of Product Characteristics (SmPC). 2026.
2. Patel R, Nguyen T, Chen L, et al. ASCEND-1: Phase III study of oncurimab vs comparator in EGFR-mutated metastatic NSCLC. *J Clin Oncol*. 2026;44(suppl).
3. Smith J, Adeyemi K, Rossi G, et al. Once-weekly EGFR inhibition: pharmacokinetics and exposure–response of oncurimab. *Lung Cancer*. 2025.
4. Brown P, Li S, Hernandez M, et al. Management of EGFR inhibitor–associated adverse events in NSCLC. *Oncologist*. 2024.

Maria's story

Patient profile: Maria* 62 years old


Lifestyle: Never-smoker, lives with her husband, wants treatment that fits into her routine

Condition: Recently diagnosed with metastatic EGFR-mutated NSCLC



*Maria is a fictional patient used for illustrative purposes.

Maria's treatment journey




The impact on Maria



EGFR-mutated NSCLC prevalence by region


Estimated prevalence of EGFR-mutated metastatic NSCLC varies across geographic regions.



*Data sourced from Salesforce epidemiology object

EGFR-mutated NSCLC prevalence by region


Estimated prevalence of EGFR-mutated metastatic NSCLC varies across geographic regions.



*Data sourced from Salesforce epidemiology object

EGFR-mutated NSCLC prevalence by region


Estimated prevalence of EGFR-mutated metastatic NSCLC varies across geographic regions.



*Data sourced from Salesforce epidemiology object

EGFR-mutated NSCLC prevalence by region


Estimated prevalence of EGFR-mutated metastatic NSCLC varies across geographic regions.



*Data sourced from Salesforce epidemiology object

EGFR-mutated NSCLC prevalence by region

Estimated prevalence of EGFR-mutated metastatic NSCLC varies across geographic regions.



Wales

▼

210 Estimated patients per year with EGFR-mutated metastatic NSCLC in the **Wales**

1,750 Metastatic NSCLC cases/year

12% EGFR mutation rate

180 Eligible patients/year

*Data sourced from Salesforce epidemiology object

Many patients face a daily treatment routine

Daily dosing can become part of a patient's everyday burden.



TAP TO START

Many patients face a daily treatment routine

Daily dosing can become part of a patient's everyday burden.



Many patients face a daily treatment routine

Daily dosing can become part of a patient's everyday burden.



Many patients face a daily treatment routine

Daily dosing can become part of a patient's everyday burden.

ONE TABLET. ONCE A WEEK.

A simplified oral treatment routine for appropriate patients with EGFR-mutated metastatic NSCLC.

 **Oncura**
(oncurimab) 80mg



 **Oncura**
(oncurimab) 80mg


Which treatment options would you consider for Maria?

ONCURA



1x 80mg tablet per week

EGRIFEX




14 × 50mg tablets per week
(2 tablets per day)

CAPSORIN



21 × 30mg capsules per week
(3 capsules per day)

INJECTRA



7 × 1mL Injections per week
(daily injection)

*Drug names and doses shown are fictitious and for illustration only.

Which treatment options would you consider for Maria?

1


ONCURA



1x 80mg tablet per week

2

EGRIFEX



14 × 50mg tablets per week
(2 tablets per day)


CAPSORIN



21 × 30mg capsules per week
(3 capsules per day)

3

INJECTRA




7 × 1mL injections per week
(daily injection)

*Drug names and doses shown are fictitious and for illustration only.

Proven efficacy in EGFR-mutated metastatic NSCLC

Oncura demonstrated significant tumour response and durable disease control in patients with EGFR-mutated metastatic non-small cell lung cancer.



Overall response rate*



Median progression-free survival*




Rapid response observed within first 6 weeks*

*Based on phase III clinical trial data.
EGFR: epidermal growth factor receptor, NSCLC: Non-Small Cell Lung Cancer.

Proven efficacy in EGFR-mutated metastatic NSCLC


Oncura demonstrated significant tumour response and durable disease control in patients with EGFR-mutated metastatic non-small cell lung cancer.



Overall response rate*



Median progression-free survival*




Rapid response observed within first 6 weeks*



*Based on phase III clinical trial data.
EGFR: epidermal growth factor receptor, NSCLC: Non-Small Cell Lung Cancer.

Proven efficacy in EGFR-mutated metastatic NSCLC


Oncura demonstrated significant tumour response and durable disease control in patients with EGFR-mutated metastatic non-small cell lung cancer.



Overall response rate*



Median progression-free survival*



Rapid response observed within first 6 weeks*




*Based on phase III clinical trial data.
EGFR: epidermal growth factor receptor, NSCLC: Non-Small Cell Lung Cancer.

Significant improvement in progression-free survival

ASCEND-1 Phase III Study

Randomised, double-blind, phase III study evaluating Oncura versus standard EGFR therapy in patients with EGFR-mutated metastatic NSCLC.



Choose a region

- ☒ Median PFS: 18.4 months Oncura
- ☒ Median PFS: 11.2 months Egrifex
- ☒ Median PFS: 10.1 months Capsorin
- ☒ Median PFS: 8.0 months Injectra

STUDY DESIGN +


*Data from ASCEND-1 phase III study
HR: hazard ratio; PFS: progression-free survival

Significant improvement in progression-free survival

ASCEND-1 Phase III Study

Randomised, double-blind, phase III study evaluating Oncura versus standard EGFR therapy in patients with EGFR-mutated metastatic NSCLC.

COMPETITORS ☐



Choose a region

- ☒ Median PFS: 18.4 months Oncura
- ☐ Median PFS: 11.2 months Egrifex
- ☐ Median PFS: 10.1 months Capsorin
- ☐ Median PFS: 8.0 months Injectra

STUDY DESIGN +

*Data from ASCEND-1 phase III study
HR: hazard ratio; PFS: progression-free survival

Significant improvement in progression-free

STUDY DESIGN



ASCEND-1 STUDY DESIGN

Objective: To evaluate the efficacy and safety of Oncura in patients with EGFR-mutated metastatic NSCLC.

Key inclusion criteria:

- Adults ≥ 18 years
- Confirmed EGFR mutation
- Metastatic NSCLC
- No prior EGFR-targeted therapy

Primary endpoint:

- Progression-free survival (PFS)

Secondary endpoints:

- Overall response rate (ORR)
- Overall survival (OS)
- Duration of response
- Safety and tolerability

Treatment arms:

Oncura 80 mg once weekly

- n = 312

Comparator EGFR therapy

- n = 310

Completed study

- ~92–95% completion rate


*Data from ASCEND-1 phase III study
HR: hazard ratio; PFS: progression-free survival

Consistent and manageable tolerability profile

Oncura demonstrated a tolerability profile consistent with EGFR-targeted therapies, with most adverse events being low grade and manageable.*



Low rates of
treatment
discontinuation
Due to adverse
events



Most adverse
events were
Grade 1–2
Mild to moderate
in severity



Dose
modifications
supported
continued therapy
In most patients



No unexpected
safety signals
observed
In the phase III
study




*Based on phase III clinical trial data.

Consistent and manageable tolerability profile

Oncura demonstrated a tolerability profile consistent with EGFR-targeted therapies, with most adverse events being low grade and manageable.*



Low rates of
treatment
discontinuation
Due to adverse
events



Most adverse
events were
Grade 1–2
Mild to moderate
in severity



Dose
modifications
supported
continued therapy
In most patients



No unexpected
safety signals
observed
In the phase III
study




*Based on phase III clinical trial data.

Simple, once-weekly oral dosing


Oncura is taken as a single oral tablet once weekly, offering a simplified treatment routine for appropriate patients with EGFR-mutated metastatic NSCLC.

Daily therapy vs once-weekly Oncura





80 mg tablet
Taken orally once weekly



With or without food Flexible administration



Simple weekly routine No daily dosing required

*Based on phase III clinical trial data.

Targeting the EGFR driver mutation

Oncura is an oral EGFR inhibitor designed to block signalling that drives tumour cell growth.

1

EGFR mutation

Abnormal EGFR signalling drives uncontrolled cell growth

2

Oncura inhibits EGFR

Blocks downstream signalling pathways

3

Reduced tumour growth

Slows proliferation and supports disease control

Oral, targeted therapy for **EGFR-mutated metastatic NSCLC**


- Designed for **once-weekly dosing**
- Targets the driver pathway underlying tumour growth

MOA VIDEO +

EGFR: epidermal growth factor receptor.

Tamoxifen and the EGFR driver mutation

MOA VIDEO



EGFR: epidermal growth factor receptor.



PATIENT | ORAL BENEFITS | EFFICACY | [TOLERABILITY](#) | DOSING | SUMMARY

 **Oncura**
(oncurimab) 80mg

Adverse events: most common events

The majority of adverse events were low grade and manageable with supportive care and dose modifications.

Adverse	Any grade (%)	Grade $\geq$ 3 (%)
Rash	45	5
Diarrhoea	35	3
Dry skin	28	1
Paronuchia	18	1
Stomatitis	14	1
Fatigue	22	2
Elevated ALT/AST	12	2
Decrease	10	<1

- Low discontinuation due to AEs

- Dose interruption/reduction supported continuation in most patients

- Early recognition + supportive care are important

ALT/AST: liver enzymes.

Supporting adherence with a once weekly routine

Clear guidance and simple routines can help patients stay on track with once-weekly Oncura therapy.

Set a weekly treatment day

Encourage patients to choose the same day each week.

Use reminders

Calendar alerts or phone reminders can support adherence.

Link dosing to a routine

For example, take the tablet with a regular weekly activity.

Provide clear instructions

Ensure patients understand what to do if a dose is missed.

Encourage open communication

Patients should report side effects or concerns promptly.

This reinforces: Weekly rhythm, Simplicity, Predictability*



*For healthcare professionals only
ALT/AST: liver enzymes.



transforming lives with once-weekly targeted therapy


ONCURA® (oncurimab) is taken as a single oral tablet once weekly, offering a simplified treatment routine for appropriate patients with EGFR-mutated metastatic NSCLC.



Proven efficacy
Strong tumour response and durable disease control




Manageable tolerability
Consistent and predictable safety profile



Once-weekly oral dosing A simplified treatment routine

*Indicated for adult patients with EGFR-mutated metastatic NSCLC.

ONCURA® (oncurimab) 80 mg tablet



Prescribing Information content (Fictional but structured to match real oncology PI pages)

Please refer to the full Summary of Product Characteristics (SmPC) before prescribing, particularly in relation to side effects, precautions, and contraindications.

Presentation

Oncura 80 mg film-coated tablets.

Each tablet contains 80 mg oncurimab.

Indication:

Oncura is indicated for the treatment of adult patients with EGFR-mutated metastatic non-small cell lung cancer (NSCLC).

Treatment should be initiated by a healthcare professional experienced in the use of anticancer therapies.

Dosage and administration:

Recommended dose:

80 mg taken orally once weekly.

- Tablets may be taken with or without food.
- Tablets should be swallowed whole with water.
- Do not crush or chew tablets.

Missed dose:

- If a dose is missed, it may be taken within 48 hours.
- If more than 48 hours have passed, skip the missed dose and resume the regular weekly schedule.

Dose modifications:

Dose interruption or reduction may be required in patients experiencing significant adverse reactions. Refer to full prescribing information.

Contraindications

- Hypersensitivity to the active substance or any excipients.

Warnings and precautions

- Interstitial lung disease (ILD) has been reported with EGFR inhibitors.
- Monitor for respiratory symptoms.
- Dermatologic reactions may occur.
- Monitor liver function during treatment.

Interactions

Caution should be exercised when co-administering with:

- Strong CYP3A4 inhibitors
- Strong CYP3A4 inducers

Fertility, pregnancy and lactation

- Not recommended during pregnancy.
- Effective contraception should be used during treatment and for at least 1 month after the last dose.
- Breastfeeding is not recommended during treatment.

Undesirable effects

Very common (≥1/10):	Common (≥1/100 to <1/10):
• Rash • Diarrhoea	• Dry skin • Fatigue • Paronychia • Stomatitis • Elevated liver enzymes • Decreased appetite

Overdose

In the event of overdose, patients should be monitored and supportive treatment initiated as appropriate.

Legal category

Prescription-only medicine (POM)

Adverse event reporting

Adverse events should be reported.
Reporting forms and information can be found at the appropriate national reporting authority website.


Adverse events should also be reported to:
Makana Health Medical Information
Email: safety@makanahealth.com
Tel: +44 (0)0000 000000

Marketing authorisation holder
Makana Health Ltd.
[Address placeholder]


ONCURA RESOURCES




Access materials to support your clinical practice and patient discussions.




Oncura Dosing
guide




Oncura Safety
summary




Oncura Prescribing
Information




Oncura MOA



Oncura Digital
Leavepiece



Oncura Patient
Profile Leavepiece



Makana Health
Website

