Supplementary Material for: A Data-Driven, Component-Level Projection Framework for 21st-Century Sea-Level Rise

Minchew, Meyer, et al.

Contents

1 Materials and Methods **2**
- 1.1 Observational datasets . . . 2
- 1.2 Variables and parameters . . . 4
- 1.3 Component model posteriors . . . 10
- 1.4 Greenland surface mass balance emulator . . . 16
- 1.5 Variance decomposition . . . 20
- 1.6 Temperature scenarios . . . 21
- 1.7 Component-level skill metrics . . . 21

2 Exceedance Probability by Threshold and Scenario **25**

3 Value of Information Analysis **26**
- 3.1 Decision framework . . . 26
  - 3.1.1 Adaptation cost . . . 27
  - 3.1.2 Damage function . . . 27
- 3.2 Two-world structure . . . 27
- 3.3 EVPI calculation . . . 28
  - 3.3.1 Hedging cost (no information) . . . 28
  - 3.3.2 Perfect information cost . . . 28
  - 3.3.3 Annual EVPI . . . 28
- 3.4 Discounting and the VoI surface . . . 29
- 3.5 Key assumptions and limitations . . . 29

1 Materials and Methods

Unless otherwise noted, all reported intervals are 90% Bayesian credible intervals (equal-tailed), and where 1σ is reported, we state explicitly that it is a standard deviation.

1.1 Observational datasets

All component models are calibrated against published observational products (Table S1). We use:

- **Global mean sea level:** NASA satellite altimetry (TOPEX/Jason, 1993–2025; *Beckley et al.*, 2017; *Hamlington et al.*, 2020), with the Frederikse et al. (2020) tide-gauge reconstruction (1900–2018) and the Dangendorf et al. (2024) reconstruction (1900–2021) for pre-satellite validation. None of these records enters any component calibration.

- **Ocean thermal expansion:** NOAA thermosteric sea level (doi:10.7289/V53F4MVP), 0–700 m (1955–2025) and 0–2000 m (2005–2025) (*Levitus et al.*, 2012). Both records are scaled by 1.14 before fitting. This corrects the 14% low bias that *Li et al.* (2022) find in thermosteric trends computed with linear vertical interpolation. A residual structural uncertainty of 5% of the corrected signal is retained. The 700–2000 m layer is carried by the 0–2000 m record from 2005 onward and by the deep layer of the two-layer model at earlier times. A constant 0.12 ± 0.04 mm yr^{-1} (one standard deviation, 1991–2010) (*Desbruyères et al.*, 2016) is added over the full record for the contribution below 2000 m. Two reconstructions of global-mean steric sea level are withheld for validation, from Frederikse et al. (2020) (1900–2018) and Dangendorf et al. (2024) (1900–2021). Over 1957–2018 the Frederikse et al. reconstruction draws on the same archive of in situ profiles as the NOAA record (*Levitus et al.*, 2012; *Ishii et al.*, 2017; *Cheng and Zhu*, 2016), so it is not fully independent of it there. The Dangendorf et al. reconstruction uses no in situ ocean temperature or salinity. Its steric term is the residual of a sea-level budget whose barystatic input is from *Frederikse et al.* (2020). Neither reconstruction carries the 1.14 scaling.

- **Glaciers:** GlaMBIE global consensus (The GlaMBIE Team 2025), combining glaciological field measurements, satellite altimetry, DEM differencing, and GRACE/GRACE-FO gravimetry across all 19 RGI regions. We use 2000–2023 (24 annual points) for calibration. For validation we use the global glacier mass changes of *Zemp et al.* (2019) for 1962–1999 (38 hydrological years).

- **Greenland:** *Mouginot et al.* (2019) SMB and discharge (1972–2018), of which the 41 years from 1978 enter the discharge fit, the years covered by the delayed ocean forcing at every candidate lag. *Mankoff et al.* (2021) discharge (1986–2023) and the IMBIE-3 Greenland record (1972–2023; *Otosaka et al.* (2026)) are withheld for validation. Subsurface ocean temperature from EN4 v4.2.2 with g10 bias corrections (200–500 m, Greenland peripheral waters, 1970–2021; *Good et al.* (2013)). Surface temperature from Berkeley Earth (*Rohde and Hausfather*, 2020). Greenland SMB is projected with a statistical emulator of the regional climate model MARv3.12 driven by seven

CMIP6 global climate models (CESM2, MPI-ESM1-2-HR, NorESM2-MM, UKESM1-0-LL, CNRM-CM6-1, CNRM-ESM2-1, and IPSL-CM6A-LR). These simulations were produced for the PROTECT project (*Boberg et al.*, 2026) and are publicly available at `http://ftp.climato.be/fettweis/MARv3.12/Greenland/PROTECT`. We use the annual SMB of the contiguous ice sheet, excluding peripheral glaciers, over 1980–2014 and over each available SSP to 2100, together with the global mean surface temperature of each driving model from the CMIP6 archive. *Mankoff et al.* (2021) SMB (1986–2023), the mean of three regional climate models driven by reanalysis, is withheld for validation of the emulator.

- **Antarctica:** IMBIE-3 mass balance (1979–2023, 45 annual points per region; *Otosaka et al.* (2026)), partitioned into West Antarctica, East Antarctica, and the Antarctic Peninsula.

- **Terrestrial water storage:** *Frederikse et al.* (2020) reconstruction, used only from 2003 onward, the period over which it derives from GRACE and GRACE-FO observations rather than from models. Prior to 2003, the reconstructed TWS curves are largely unconstrained by independent observations.

- **Temperature scenarios:** IPCC AR6 assessed GMST projections (SSP1-1.9, SSP1-2.6, SSP2-4.5, SSP3-7.0, SSP5-8.5; *Lee et al.* (2021)).

Table S1: Datasets used in the component framework and the role each plays. Records marked *validation* are withheld from every component calibration, including observed GMSL itself. A complete dataset inventory including versions, spatial domains, and processing details is provided in `data/dataset_inventory.xlsx`.

Component	Dataset	Span	Role	Reference
GMSL	NASA altimetry	1993–2025	Validation	Beckley et al. (2017)
GMSL	Frederikse reconstruction	1900–2018	Validation	Frederikse et al. (2020)
GMSL	Dangendorf reconstruction	1900–2021	Validation	Dangendorf et al. (2024)
Thermosteric	NOAA thermosteric, 0–700 m	1955–2025	Calibration	Levitus et al. (2012)
Thermosteric	NOAA thermosteric, 0–2000 m	2005–2025	Calibration	Levitus et al. (2012)
Thermosteric	Frederikse steric	1900–2018	Validation	Frederikse et al. (2020)
Thermosteric	Dangendorf sterodynamic	1900–2021	Validation	Dangendorf et al. (2024)
Glaciers	GlaMBIE global consensus	2000–2023	Calibration	The GlaMBIE Team (2025)
Glaciers	Zemp glaciological and geodetic	1962–1999	Validation	Zemp et al. (2019)
Greenland	Mouginot discharge & SMB	1972–2018	Calibration	Mouginot et al. (2019)
Greenland	EN4 subsurface ocean T	1970–2021	Forcing	Good et al. (2013)
Greenland	Mankoff PROMICE discharge & SMB	1986–2023	Validation	Mankoff et al. (2021)
Greenland	IMBIE-3 Greenland	1972–2023	Validation	Otosaka et al. (2026)
Greenland	MARv3.12 SMB, seven CMIP6 GCMs	1980–2100	Emulator training	Boberg et al. (2026)
Greenland	GMST of the seven driving GCMs	1950–2100	Emulator training	CMIP6 archive
Peninsula	IMBIE-3 Peninsula	1979–2023	Calibration	Otosaka et al. (2026)
EAIS	IMBIE-3 East Antarctica	1979–2023	Diagnostic	Otosaka et al. (2026)
WAIS	IMBIE-3 West Antarctica	1979–2023	Calibration	Otosaka et al. (2026)
TWS	Frederikse TWS (GRACE era)	2003–2018	Forcing	Frederikse et al. (2020)
Temperature	Berkeley Earth surface T	1850–2024	Forcing	Rohde & Hausfather (2020)
Temperature	IPCC AR6 GMST projections	2015–2099	Forcing	Lee et al. (2021)
Projections	IPCC AR6 FACTS	2020–2150	Comparison	Garner et al. (2023)

1.2 Variables and parameters

Table S2 lists every symbol that appears in the main text, with a plain-language description, the value where one is assigned, units, and the source of any assigned value. Symbols used only in this supplement, including those of the value-of-information analysis, are defined where they appear there and are not repeated here.

Table S2: **Symbols used in the main text.** The value column gives the value of parameters that are assigned rather than inferred. Quantities we calibrate are marked *calibrated* and their posterior medians and 90% credible intervals are reported in the main text, since a fitted value is not a prescribed one. Model variables that carry no single value are marked with a dash. Where one symbol carries more than one meaning, each meaning is listed separately and its context named. SLE denotes sea-level equivalent.

Symbol	Description	Value	Units	Source
General				
t	Time	—	yr	—
T	Global mean surface temperature (GMST) anomaly, referenced to 1850–1900	—	°C	*Rohde and Hausfather* (2020)
ΔT	GMST anomaly forcing the EAIS surface-mass-balance model, relative to 1995–2005	—	°C	—
t_0	Study baseline year, to which cumulative contributions are referenced in the component models	2000	yr	—
t_0	Forecast origin, the most recent year of satellite altimetry in the blended forecast	2025	yr	—
i	Index over observations, outlet glaciers, or Monte Carlo samples	—	—	—
z	Depth below the sea surface	—	m	—
Greenland Ice Sheet discharge				
D	Greenland solid-ice discharge, as a contribution to sea level	—	$\mathrm{mm\,SLE\,yr^{-1}}$	—
H_{D}	Cumulative Greenland discharge contribution to GMSL	—	mm SLE	—
γ	Sensitivity of discharge to subsurface ocean temperature	calibrated	$\mathrm{mm\,SLE\,yr^{-1}\,°C^{-1}}$	this study
r_0	Background discharge rate at the baseline ocean temperature	calibrated	$\mathrm{mm\,SLE\,yr^{-1}}$	this study
δ	Lag between ocean warming and the discharge response	6, selected by BIC	yr	this study
T_{ocean}	Subsurface ocean temperature anomaly, 200–500 m, in Greenland's peripheral waters	—	°C	*Good et al.* (2013)
Dimensional scaling for γ				
$\widetilde{\gamma}$	Order-of-magnitude estimate of γ from the scaling argument	0.05–1.5	$\mathrm{mm\,SLE\,yr^{-1}\,°C^{-1}}$	this study
$\Delta\dot{m}_i$	Submarine melt rate of outlet glacier i	—	$\mathrm{m\,yr^{-1}}$	—

Continued on next page.

Table S2 continued from previous page.

Symbol	Description	Value	Units	Source
κ_i	Melt sensitivity to ocean thermal forcing, averaged over the calving face	36–104, median 60	$\mathrm{m\,yr^{-1}\,{}^{\circ}C^{-1}}$	*Xu et al.* (2013); *Rignot et al.* (2016); *Slater and Straneo* (2022); *Wood et al.* (2021)
μ_i	Ratio of the total ice-flux perturbation to the direct submarine melt flux	3–10	—	*Enderlin and Howat* (2013); *Wood et al.* (2021)
$A_{\mathrm{f},i}$	Submerged calving-front area of outlet glacier i	0.9–2.7	$\mathrm{km^2}$	*Rignot et al.* (2016)
$\Delta\dot{Q}_i$	Ice-flux perturbation from outlet glacier i	—	$\mathrm{Gt\,yr^{-1}}$	—
N	Number of marine-terminating outlet glaciers resolved in the record	219	—	*Mouginot et al.* (2019)
ρ_{ice}	Density of ice	900	$\mathrm{kg\,m^{-3}}$	—
ρ_{fw}	Density of freshwater	1000	$\mathrm{kg\,m^{-3}}$	—
A_{ocean}	Global ocean area, setting the convention $1\,\mathrm{mm\,SLE} = 362.5\,\mathrm{Gt}$	3.625×10^{14}	$\mathrm{m^2}$	—
Thermosteric two-layer model				
H_{u}	Steric height of the upper ocean layer	—	mm	—
H_{ud}	Steric height of the upper 2000 m	—	mm	—
H_{full}	Full-depth thermosteric contribution to GMSL	—	mm	—
α_{sw}	Thermal expansion coefficient of seawater	—	$\mathrm{{}^{\circ}C^{-1}}$	—
α'_{sw}	Derivative of α_{sw} with respect to temperature	—	$\mathrm{{}^{\circ}C^{-2}}$	—
$\Delta\theta$	Ocean temperature anomaly profile	—	°C	—
φ	Dimensionless shape factor of the temperature anomaly profile	—	—	—
h_{u}	Thickness of the upper layer	700	m	—
h_{d}	Lower bound of the deep layer	2000	m	—
S_{u}	Thermal state of the upper layer, the amplitude of its temperature anomaly	—	°C	—
S_{d}	Thermal state of the deep layer	—	°C	—
a_{u}	Quadratic expansion coefficient of the upper layer, from the temperature dependence of α_{sw} and other departures from a linear response	calibrated	$\mathrm{mm\,{}^{\circ}C^{-2}}$	this study
b_{u}	Linear expansion coefficient of the upper layer	calibrated	$\mathrm{mm\,{}^{\circ}C^{-1}}$	this study

Continued on next page.

Table S2 continued from previous page.

Symbol	Description	Value	Units	Source
b_{d}	Linear expansion coefficient of the deep layer	calibrated	$\mathrm{mm\,°C^{-1}}$	this study
τ_{u}	Relaxation time of the upper layer toward GMST	calibrated	yr	this study
τ_{d}	Relaxation time of the deep layer toward the upper layer, held fixed	150	yr	*Geoffroy et al.* (2013)
$\dot{H}_{\mathrm{b}}$	Rate of steric expansion below 2000 m, adopted rather than calibrated	0.12 ± 0.04	$\mathrm{mm\,yr^{-1}}$	*Desbruyères et al.* (2016)
t_{ref}	Reference year for the below-2000 m term	2000	yr	—
σ_{extra}	Noise term absorbing misfit beyond the reported observational error	calibrated	mm	this study
$\boldsymbol{\phi}$	Vector of the six fitted thermosteric parameters	—	—	—
$\mathcal{J}$	Objective function, twice the negative log-posterior up to a constant	—	—	—
r_i	Residual between the demeaned model and observation i	—	mm	—
σ_i	Total standard deviation of observation i	—	mm	—
$\mu_{\tau_{\mathrm{u}}}$	Mean of the log-normal prior on $\ln \tau_{\mathrm{u}}$	$\ln 8$	ln(yr)	*Geoffroy et al.* (2013)
$s_{\tau_{\mathrm{u}}}$	Standard deviation of the log-normal prior on $\ln \tau_{\mathrm{u}}$	0.5	—	*Geoffroy et al.* (2013)
Glaciers				
H_{gl}	Cumulative glacier contribution to GMSL	—	mm	—
b_{gl}	Sensitivity of the glacier loss rate to warming	calibrated	$\mathrm{mm\,yr^{-1}\ °C^{-1}}$	this study, prior from *Rounce et al.* (2023)
c_{gl}	Glacier loss rate at the 1995–2005 mean GMST	$-b_{\mathrm{gl}} T_{\mathrm{gl,eq}}$	$\mathrm{mm\,yr^{-1}}$	this study
$T_{\mathrm{gl,eq}}$	GMST at which the glacier loss rate is zero (1850 balance), relative to 1995–2005	calibrated, prior $\mathcal{N}(-0.86,\, 0.15^2)$	°C	*Roe et al.* (2021), this study
$H_{\mathrm{gl},0}$	Glacier level at the baseline year	calibrated	mm	this study
$\varepsilon_{\mathrm{gl}}$	Residual of the glacier fit	—	mm	—
$\sigma_{\mathrm{gl,extra}}$	Model-inadequacy term added in quadrature to the annual GlaMBIE uncertainty	calibrated	$\mathrm{mm\,yr^{-1}}$	this study
Σ_{gl}	Covariance matrix of the cumulative glacier record	—	$\mathrm{mm^2}$	*The GlaMBIE Team et al.* (2025)
Antarctic Peninsula				

Continued on next page.

Table S2 continued from previous page.

Symbol	Description	Value	Units	Source
$\dot{H}_{\mathrm{p}}$	Rate of the Antarctic Peninsula contribution	—	$\mathrm{mm\,yr^{-1}}$	—
a_{p}	Quadratic sensitivity to warming, set to zero because BIC favors the linear model	0	$\mathrm{mm\,yr^{-1}\,{}^{\circ}C^{-2}}$	this study
b_{p}	Sensitivity of the Peninsula rate to warming	calibrated	$\mathrm{mm\,yr^{-1}\,{}^{\circ}C^{-1}}$	this study
c_{p}	Peninsula rate at the 1995–2005 mean GMST	calibrated	$\mathrm{mm\,yr^{-1}}$	this study
$H_{\mathrm{p},0}$	Peninsula level at the first year of the record (1979)	calibrated	mm	this study
$\sigma_{\mathrm{p,extra}}$	Model-inadequacy term for annual scatter beyond the reported uncertainty	calibrated	$\mathrm{mm\,yr^{-1}}$	this study
East Antarctic Ice Sheet				
b_{e}	Sensitivity of the EAIS rate to warming, fitted as a check and not used in the forecast	calibrated	$\mathrm{mm\,yr^{-1}\,{}^{\circ}C^{-1}}$	this study, prior from *Frieler et al.* (2015); *Ligtenberg et al.* (2013)
c_{e}	EAIS rate at the 1995–2005 mean GMST	calibrated	$\mathrm{mm\,yr^{-1}}$	this study
$H_{\mathrm{e},0}$	EAIS level at the first year of the record	calibrated	mm	this study
West Antarctic Ice Sheet				
a_{w}	Acceleration of WAIS mass loss in the quadratic-in-time fit	calibrated	$\mathrm{mm\,yr^{-2}}$	this study
$p(\mathrm{S1})$	Probability assigned to WAIS Scenario 1, the control variable of the scenario mixture	0.90 slow, 0.10 fast	—	this study
λ	Shape parameter of the skew-normal S2 distribution in log space	3	—	this study, motivated by *Robel et al.* (2019)
β	Exponent of the power law in time followed by the S2 trajectory	2	—	this study
Surface mass balance				
$\dot{H}_{\mathrm{smb}}$	Surface mass balance rate. For Greenland it is expressed as a contribution to sea-level rise	—	$\mathrm{Gt\,yr^{-1}}$ (EAIS), $\mathrm{mm\,SLE\,yr^{-1}}$ (Greenland)	—
a_{gris}	Quadratic sensitivity of the Greenland SMB contribution to warming	fitted to MAR for each of seven GCMs (Fig. S6)	$\mathrm{mm\,SLE\,yr^{-1}\,{}^{\circ}C^{-2}}$	this study, from MARv3.12
b_{gris}	Linear sensitivity of the Greenland SMB contribution to warming	fitted to MAR for each of seven GCMs (Fig. S6)	$\mathrm{mm\,SLE\,yr^{-1}\,{}^{\circ}C^{-1}}$	this study, from MARv3.12
c_{gris}	Greenland SMB contribution at zero GMST anomaly, fitted with the drift state	-1.07 [-1.44, -0.78]	$\mathrm{mm\,SLE\,yr^{-1}}$	this study, from *Mouginot et al.* (2019)

Continued on next page.

Table S2 continued from previous page.

Symbol	Description	Value	Units	Source
s	Drift state, the slowly varying departure of observed SMB from the emulator	fitted	$\mathrm{Gt\,yr^{-1}}$	this study, from *Mouginot et al.* (2019)
τ	Relaxation timescale of the drift state, set by a lognormal prior with 90% range 5–25 yr	12 [6, 26]	yr	this study
σ_s	Standard deviation of the drift state	108 [60, 174]	$\mathrm{Gt\,yr^{-1}}$	this study
σ_ε	Standard deviation of year-to-year weather in Greenland SMB	50 [25, 80]	$\mathrm{Gt\,yr^{-1}}$	this study
ϵ_{fb}	Fractional SMB–elevation feedback at the reference simulation, truncated at zero	0.08 ± 0.05	—	*Fettweis et al.* (2013)
C_T	Linear sensitivity of EAIS SMB to GMST, positive for mass gain by accumulation	60 ± 20	$\mathrm{Gt\,yr^{-1}\ {}^\circ C^{-1}}$	*Frieler et al.* (2015); *Ligtenberg et al.* (2013)
C_{T^2}	Quadratic sensitivity of EAIS SMB, set to zero	0	$\mathrm{Gt\,yr^{-1}\ {}^\circ C^{-2}}$	this study
$\dot{M}_0$	Baseline EAIS SMB, set to zero because IMBIE-3 shows EAIS near balance	0	$\mathrm{Gt\,yr^{-1}}$	*Otosaka et al.* (2026)
Greenland temperature chain				
$T^{\mathrm{Gr}}_{\mathrm{surface}}$	Greenland regional surface temperature anomaly, averaged over the Berkeley Earth land cells in Greenland	—	°C	*Rohde and Hausfather* (2020)
ξ	Slope of the surface-to-ocean transfer function, fitted by ordinary least squares	fitted, 0.147 ± 0.023	—	this study
χ	Intercept of the surface-to-ocean transfer function	fitted, -0.056 ± 0.048	°C	this study
$\varepsilon_{\mathrm{go}}$	Zero-mean residual of the transfer function	—	°C	—
AA	Arctic amplification factor, the ratio of Greenland to global warming, used to convert GMST to Greenland regional temperature for the discharge model.	2.5 ± 0.5	—	*Rantanen et al.* (2022)
Forecast construction				
$\dot{H}_{\mathrm{forecast}}$	Blended forecast rate of GMSL rise	—	$\mathrm{mm\,yr^{-1}}$	—
H_{forecast}	Forecast GMSL level	—	mm	—
$\dot{H}_{\mathrm{obs}}$	Rate from the satellite-era quadratic fit to altimetry	—	$\mathrm{mm\,yr^{-1}}$	*Hamlington et al.* (2024); *Nerem et al.* (2022)
$\dot{H}_{\mathrm{comp}}$	Rate of the summed calibrated components	—	$\mathrm{mm\,yr^{-1}}$	—

Continued on next page.

Table S2 continued from previous page.

Symbol	Description	Value	Units	Source
H_k	Level-space projection ensemble for component k	—	mm	—
w	Weight blending the observed and component rates	—	—	—
Θ	Standard logistic function	—	—	—
t_r	Midpoint year of the blend transition	2035 forecast, 2015 hindcast	yr	this study
τ_r	Width of the blend transition	5	yr	this study
Diagnostics				
$\dot{H}$	Rate of GMSL rise	—	$\mathrm{mm\,yr^{-1}}$	—
α	Transient sensitivity of the rate of GMSL rise to warming, $d\dot{H}/dT$	—	$\mathrm{mm\,yr^{-1}\,{}^\circ C^{-1}}$	—
R^2	Coefficient of determination	—	—	—
χ^2/dof	Reduced chi-square, the error-weighted sum of squared residuals per degree of freedom	—	—	—

1.3 Component model posteriors

This section collects the parameter posteriors of every calibrated component model (Figs. S1–S5). Greenland discharge is fitted by weighted least squares to the annual increments of its cumulative record over a discrete set of delays, so its posterior is a mixture of normals weighted by the Bayesian Information Criterion and is shown as marginal densities. The remaining components are sampled by Markov chain Monte Carlo and are shown as corner plots, with marginal histograms on the diagonal and pairwise joint densities on the off-diagonal. Posterior medians and their 90% credible intervals are reported in the main text, and they are the values carried into the projections.




Figure S1: **Greenland discharge delay model parameter posteriors.** Marginal posteriors for the two fitted parameters and the selected delay of the discharge delay model calibrated against *Mouginot et al.* (2019). **(a)** The sensitivity of the discharge rate to the delayed ocean temperature anomaly, γ, and **(b)** the background discharge rate, r_0. **(c)** The delay δ is resolved only to integer years, so its posterior is the set of weights over the candidate delays, which place 98.0% of the weight on $\delta = 6\,\mathrm{yr}$. Panel titles give the median with the 90% credible interval.




Figure S2: **Two-layer thermosteric model posterior correlation structure.** The joint posterior for the five parameters of the two-layer energy balance model fitted to NOAA 0–700 m and 0–2000 m thermosteric sea level data. Diagonal panels show marginal posterior histograms (red), with dashed vertical lines marking the median and 90% credible interval. Off-diagonal panels show the joint posterior density, with contours denoting 1, 2, and 3 standard deviations. Only τ_u has an informative prior (black curve) while the other four parameters have flat, uninformative priors (not shown).




Figure S3: **Glacier model posterior correlation structure.** The joint posterior for the four free parameters of the glacier model (b_{gl}, $T_{gl,eq}$, $\sigma_{gl,extra}$, and $H_{gl,0}$) fitted to the GlaMBIE record (*The GlaMBIE Team et al.*, 2025), with the full covariance of the cumulative record in the likelihood. Diagonal panels show marginal posterior histograms (gray), with dashed vertical lines marking the median and 90% credible interval. Off-diagonal panels show the joint posterior density, with contours denoting 1, 2, and 3 standard deviations. Black curves show the two priors: b_{gl}, which is centered on the sensitivity of *Rounce et al.* (2023) and weakly informative, and $T_{gl,eq}$, which is centered on the 1850–1879 mean GMST and fixes the rate at the 1995–2005 mean GMST, $c_{gl} = -b_{gl}T_{gl,eq}$, while $\sigma_{gl,extra}$ and $H_{gl,0}$ have wide priors that are not shown.




Figure S4: **Antarctic Peninsula model posterior correlation structure.** The joint posterior for the four parameters of the Peninsula model fitted to the IMBIE-3 Peninsula record (*Otosaka et al.*, 2026), with the full covariance of the cumulative record in the likelihood. The fourth parameter is the model-inadequacy term σ_{extra}, fitted here, for glaciers, and for WAIS, but not for EAIS. Diagonal panels show marginal posterior histograms (blue), with dashed vertical lines marking the median and 90% credible interval. Off-diagonal panels show the joint posterior density, with contours denoting 1, 2, and 3 standard deviations. Only b_{AP} has a prior shown (black curve), a symmetric normal that is flat over the range the data support, while the remaining three parameters have wide priors that are not shown.




Figure S5: **EAIS model posterior correlation structure.** The joint posterior for the three parameters of the East Antarctic model fitted to the IMBIE-3 East Antarctic record (*Otosaka et al.*, 2026), with the full covariance of the cumulative record in the likelihood. This fit serves as a diagnostic, the projection instead adopting a literature surface-mass-balance sensitivity. Diagonal panels show marginal posterior histograms (pale blue), with dashed vertical lines marking the median and 90% credible interval. Off-diagonal panels show the joint posterior density, with contours denoting 1, 2, and 3 standard deviations. Only b_{EAIS} has an informative prior (black curve), centered on the literature sensitivity $C_T = 60 \pm 20$ Gt yr^{-1} °C^{-1} (*Frieler et al.*, 2015; *Ligtenberg et al.*, 2013) and wide enough that the data set the posterior, while c_{EAIS} and $H_{\mathrm{EAIS,0}}$ have wide priors that are not shown.

1.4 Greenland surface mass balance emulator

We project Greenland SMB with a statistical emulator of the regional climate model MARv3.12 driven by seven CMIP6 global climate models (*Boberg et al.*, 2026): CESM2, MPI-ESM1-2-HR, NorESM2-MM, UKESM1-0-LL, CNRM-CM6-1, CNRM-ESM2-1, and IPSL-CM6A-LR. For each GCM, we fit the annual SMB contribution of the ice sheet as a quadratic in that GCM's own GMST anomaly T, relative to 1995–2005 and smoothed with an 11-year centered mean,

$$\dot{H}_{\mathrm{smb}} = a_{\mathrm{gris}}\, T^2 + b_{\mathrm{gris}}\, T + c_{\mathrm{gris}}, \tag{1}$$

with $\dot{H}_{\mathrm{smb}}$ in $\mathrm{mm\,SLE\,yr^{-1}}$ and positive for sea-level rise. Each fit pools the history run (1980–2014) with every SSP run available for that GCM (2015–2100). Each run is a separate segment with first-order autoregressive residuals. The posterior of a_{gris}, b_{gris}, and the intercept is analytic under a weak normal-inverse-gamma prior, and the autocorrelation is integrated over a grid weighted by marginal likelihood. We tested adding a lag between warming and the SMB response for CESM2 and MPI-ESM1-2-HR, the two GCMs with all three scenarios. It did not improve the fits, so the emulator has none.

In projections, we replace the intercept fitted to MAR with a level c_{gris} and a drift state $s(t)$, both fitted to the *Mouginot et al.* (2019) SMB. The emulator supplies the change in SMB with warming, and the observations supply its present level and its slow departures from the emulator. For each member, the residual of the observed SMB about that member's emulator response is modeled as $c_{\mathrm{gris}} + s(t) + \varepsilon(t)$, where s is a first-order autoregressive process with relaxation timescale τ and standard deviation σ_s, and ε combines year-to-year weather, with standard deviation σ_ε, and the reported observational error. The hyperparameters $(\tau, \sigma_s, \sigma_\varepsilon)$ are estimated on a grid from the Kalman-filter likelihood of the residual about the median emulator, with a lognormal prior on τ whose 90% range is 5–25 yr and a weak half-normal prior on σ_s. Each member draws $(\tau, \sigma_s, \sigma_\varepsilon)$ from this posterior and one joint path of c_{gris} and s through 1972–2018 by forward filtering and backward sampling. After 2018 the state follows its autoregressive process and relaxes toward zero, and before 1972 it is continued backward in the same way. The projections add white weather noise with the member's σ_ε and the SMB–elevation feedback of *Fettweis et al.* (2013). That feedback is an additional loss equal to a fraction $\epsilon_{\mathrm{fb}} \sim \mathcal{N}(0.08, 0.05^2)$, truncated at zero, of the loss at their reference simulation, scaled by the member's cumulative SMB anomaly since 2000. Each member also draws one GCM, with equal weight, and one posterior draw of a_{gris} and b_{gris} for that GCM. Because MAR is the only regional climate model in the ensemble, the spread between regional climate models is not sampled. Under the same forcing, RACMO gives less SMB loss than MAR (*Glaude et al.*, 2024).

We test the emulator out of sample by leaving out each GCM in turn and predicting its runs from the other six (Fig. S7). As in the projections, the prediction is anchored to the withheld run's own 1995–2005 mean. Over 2015–2100, all 13 withheld runs fall within the 90% predictive range. The differences between MAR and the emulator range from −19.7 to +28.7 mm SLE, with a median absolute difference of 14.0 mm. The largest are for UKESM1-0-LL, at +17.6 mm (SSP2-4.5) and +28.7 mm (SSP5-8.5), where the other six GCMs underpredict its mass loss. Fig. S8 shows the same comparison year by year. The emulator is trained on MAR alone, so it does not sample the differences between regional climate models. Forced by the same CESM2 SSP5-8.5 run, RACMO gives an SMB contri-




Figure S6: **Greenland SMB emulator fits.** **(a–g)** Annual SMB contribution of the contiguous Greenland ice sheet simulated by MARv3.12 driven by each of seven CMIP6 GCMs, relative to the 1995–2005 mean of that GCM's history run, against the GCM's own GMST anomaly T (11-year centered mean, relative to 1995–2005). Points are colored and shaped by experiment. Black curves show the posterior median of the fitted quadratic, including the fitted intercept, and gray bands its 90% credible interval. **(h)** 90% ellipse of the joint posterior of a_{gris} and b_{gris} for each GCM, labeled with the letter of its panel. All quantities are contributions to GMSL.




Figure S7: **Leave-one-GCM-out test of the Greenland SMB emulator.** Each GCM is withheld in turn, and its SSP runs are predicted by the emulator fitted to the other six GCMs, driven by the withheld GCM's GMST and anchored to its own 1995–2005 mean. **(a)** Cumulative SMB contribution over 2015–2100 relative to the 1995–2005 rate, MAR against the emulator. Horizontal bars show the 90% predictive range, which includes the spread of the six GCMs, the coefficient uncertainty, autoregressive weather noise, and the weather noise of the 1995–2005 anchor. The dashed line is the 1:1 line. **(b)** MAR minus the emulator median for each withheld run, with the same 90% range. Positive values mean that the emulator underpredicts the contribution to sea-level rise.

bution of 87 mm over 2015–2099 and MAR gives 145 mm, against an emulator prediction of 117 [88, 146] mm. The two regional models differ by 58 mm, as much as the width of the emulator's 90% range (Fig. S8n).

Driven by observed GMST, the emulator's forced response declines more slowly than the observed SMB (Fig. S9). Its median SMB falls from 387 $\mathrm{Gt\,yr^{-1}}$ over 1972–1985 to 358 $\mathrm{Gt\,yr^{-1}}$ over 2010–2018, while the *Mouginot et al.* (2019) SMB falls from 466 to 252 $\mathrm{Gt\,yr^{-1}}$. Every Mouginot year lies within the emulator's 90% range once weather noise is included, and 2 of the 38 years of the withheld *Mankoff et al.* (2021) SMB lie outside it (2012 and 2019). The emulator represents the response of SMB to global warming. It does not reproduce multi-year departures from that response, such as the low SMB of 2000–2018.

The drift state described above accounts for these departures. Fitted to the *Mouginot et al.* (2019) SMB over 1972–2018, it raises the maximum log likelihood from −293.5 for a constant level to −284.2. The likelihood is nearly flat in τ from about 10 to more than 100 yr, so τ follows its prior. Out of sample, the drift state helps little. Fitted to 1972–2004, it




Figure S8: **Greenland SMB projections of regional climate models against the emulator, 2015–2100. (a–m)** Cumulative SMB contribution since 2015 of each MARv3.12 scenario run (*Boberg et al.*, 2026) (green), relative to the 1995–2005 mean of that GCM's history run, with the prediction of the emulator fitted to the other six GCMs and driven by the withheld GCM's GMST (dashed median and shaded 90% range, colored by scenario). The range includes the spread of the six GCMs, the coefficient uncertainty, autoregressive weather noise, and the weather noise of the 1995–2005 anchor, and the values at 2100 are those of Fig. S7. **(n)** RACMO2.3p2 (*Noël et al.*, 2021) (purple) and MAR (green dotted), both forced by a CESM2 SSP5-8.5 branch run, and HIRHAM (orange dash-dot), forced by CESM2 r1i1p1f1 under SSP5-8.5, all from *Glaude et al.* (2024), integrated over the contiguous ice sheet and taken relative to their own 1995–2005 means. RACMO ends in 2099. The emulator in (n) is fitted to the six GCMs other than CESM2 and driven by the CMIP6 CESM2 SSP5-8.5 GMST, because the GMST of the branch run is not archived. All quantities are contributions to GMSL.




Figure S9: **Greenland SMB emulator driven by observed GMST.** The emulator's forced response (dark band, 90% range, and median), driven by the Berkeley Earth GMST anomaly (*Rohde and Hausfather*, 2020) with the 11-year centered mean used in training, is compared with the *Mouginot et al.* (2019) SMB, whose 1995–2005 mean anchors the emulator in this comparison, and the withheld *Mankoff et al.* (2021) SMB. The forced-response range includes the GCM, the coefficients, and the uncertainty of that anchor. The drift state is not included. The light band adds autoregressive weather noise with the parameters fitted to each GCM. The centered mean for 2020–2024 uses the IPCC AR6 2.7 °C GMST path (*Lee et al.*, 2021) for 2025–2029. The right axis gives the same SMB in Gt yr^{-1}.

forecasts the 2005–2018 SMB no better than a fixed 1995–2005 anchor, and both overpredict it. Fitted to 1972–2018, its forecast of the 2019–2023 mean SMB is slightly closer than the fixed anchor to the *Mankoff et al.* (2021) and IMBIE-3 (*Otosaka et al.*, 2026) records, and both again overpredict it. At 2100 the drift state leaves the median SMB contribution essentially unchanged and widens its 90% interval, mostly through the uncertainty of c_{gris}. A prior on τ that is uniform in $\log \tau$ over 3–150 yr moves the 2100 medians by 2–3 mm. Continued backward before 1972, the state adds 6 mm of SMB mass gain over 1900–1971 relative to a fixed anchor, because the record begins in a period of high SMB. This moves our Greenland hindcast further from the *Frederikse et al.* (2020) estimate, whose Greenland term before 1972 is the reconstruction of *Kjeldsen et al.* (2015).

1.5 Variance decomposition

We decompose the variance of the 2100 forecast separately for each warming trajectory. Pooling the trajectories would require assigning each one a probability. The total here is the sum of the component models before blending with the observed trend, excluding ter-

restrial water storage. Under the fast-WAIS weighting, WAIS accounts for 98.3%, 97.0%, and 93.3% of the total variance at 1.8, 2.7, and 3.6 °C. The thermosteric, glacier, and Greenland contributions together account for 1.1%, 2.6%, and 5.8%, a share that includes the covariance among them from the warming path they share. EAIS and the Antarctic Peninsula account for the small remainder. WAIS does not depend on the warming trajectory or on the parameters of any other component, so these shares nearly sum to one. Under the slow-WAIS weighting, $p(\mathrm{S1}) = 0.90$, the WAIS variance is that of the two-branch mixture, $p\,\mathrm{var}(S_1) + (1-p)\,\mathrm{var}(S_2) + p(1-p)\,(\bar{S}_1 - \bar{S}_2)^2$, and WAIS accounts for 94.2%, 86.8%, and 73.7% of the total variance at 1.8, 2.7, and 3.6 °C.

1.6 Temperature scenarios

The component projections are driven by the IPCC AR6 projections of global mean surface temperature for each SSP (*Lee et al.*, 2021). As in the main text, we label each scenario by the AR6 central estimate of its warming over 2081–2100 relative to 1850–1900 (*Lee et al.*, 2021, Table 4.5): 1.4 °C (SSP1-1.9), 1.8 °C (SSP1-2.6), 2.7 °C (SSP2-4.5), 3.6 °C (SSP3-7.0), and 4.4 °C (SSP5-8.5). Each SSP has a median trajectory and a 5–95% range, which comes mostly from uncertainty in the response of the climate to a given emissions pathway. A world that warms faster than the median tends to stay warmer for the whole century. We therefore treat this range as a set of warming paths rather than as independent noise in each year.

Each ensemble member k is assigned one value z_k drawn from a standard normal distribution. Its temperature is the median trajectory plus an offset $\delta_k(t) = z_k\,[h(t) - h(t_0)]$ for $t > t_0$, and zero before t_0. When $z_k > 0$, $h(t)$ is the distance from the median to the 95th percentile divided by 1.645. When $z_k < 0$, it is the distance from the 5th percentile to the median divided by 1.645. This keeps the asymmetry of the AR6 range. We set $t_0 = 2024$ because warming up to the present is observed, so the offset adds spread only to future warming. As a result, the spread at 2100 is somewhat narrower than the full AR6 range. For the 2.7 °C scenario the upper half-width at 2099 is 0.64 °C rather than 0.82 °C.

Member k uses the same z_k in every component and every SSP. Each member therefore represents one world, in which the ocean, the glaciers, and the ice sheets all respond to the same warming. The z_k are drawn with their own random seed, so the parameter draws of each component do not change. Sampling the warming range widens the 2100 intervals of the temperature-driven components, most strongly for thermal expansion. At 2.7 °C its 90% interval grows from 82 to 157 mm.

1.7 Component-level skill metrics

Table S3 collects the goodness-of-fit statistics for every component fitted to data. We report reduced chi-square rather than a coefficient of determination. Every component is calibrated against a cumulative record whose variance is dominated by its trend, so R^2 in that space approaches unity for any model that reproduces the trend at all and says little about the quality of the fit. A straight line in time, with no temperature dependence whatsoever, scores 0.983 against the 0–2000 m thermosteric record.

Reduced chi-square compares the residuals with the uncertainty the calibration data are reported to carry, so a value near unity indicates both that the model reproduces the observations to within their stated errors and that the error model used in the likelihood is well specified. The normalization is not identical in every case, and we note where it differs. Glaciers, the Antarctic Peninsula, EAIS, and WAIS are fitted to cumulative records whose errors are serially correlated, and their χ^2 is therefore evaluated against the full covariance of the cumulative record built from the reported annual uncertainties, not against a diagonal approximation, which would understate the residual scatter that the reported errors can absorb. All four sit between 0.94 and 1.05. Glaciers and the Peninsula reach those values only with fitted model-inadequacy terms, $\sigma_{\mathrm{gl,extra}}$ and $\sigma_{\mathrm{p,extra}}$, described in the main text. Without them, the reduced chi-square is 4.1 for glaciers, with the GlaMBIE 95% uncertainties converted to standard deviations, and 2.64 for the Peninsula. Both records therefore carry real annual scatter beyond their reported per-year uncertainties. EAIS does not need a model-inadequacy term. For the thermosteric model the quoted value includes the fitted excess-noise term σ_{extra} in the point variance and is normalized by the number of data points rather than by the degrees of freedom, so it sits slightly below the corresponding χ^2/dof. The 0–2000 m value of 0.37 indicates that the 21-year record is fitted to well inside its reported error. The Greenland discharge value of 1.45 is evaluated on the annual increments of the cumulative record, which is equivalent to using its full covariance, before the error scale is applied. It is the factor by which the parameter covariance is inflated, so it is a statement about the reported errors being too small rather than about the model missing the data. It has 38 degrees of freedom, the 40 annual increments between the 41 fitted points minus two for the fitted parameters. The *Mouginot et al.* (2019) discharge record runs from 1972, but EN4 begins in 1970, so the delayed ocean forcing is undefined in the first years of the record. Every candidate lag is therefore fit to the same 41 years, 1978 to 2018, those covered by the longest candidate lag of 8 yr, and the first six years of the discharge record do not enter the fit.

Two components need further comment. WAIS is fitted in time rather than in temperature, but with the same annual-increment likelihood as the Peninsula, including a fitted model-inadequacy term, so its reduced chi-square of 0.99 is comparable with the other rows. A cubic in time is not supported by the record, with $\Delta\mathrm{BIC} = +3.6$ relative to the quadratic. Snowfall anomalies also modulate the record. In the Amundsen Sea Embayment, persistently low snowfall added 51 ± 4 Gt yr^{-1} to the mass loss during 2009–2013, and extreme snowfall reduced it by 60 ± 16 Gt yr^{-1} in 2019 and 2020 (*Davison et al.*, 2023). To test their effect on the fit, we removed both anomalies from the IMBIE-3 record, converting at 362.5 Gt per mm SLE, and refitted the S1 model with the same likelihood. With the anomalies removed, the contribution at 2100 relative to 2000 falls from 86 to 79 mm. This test is approximate, because *Davison et al.* (2023) report the anomalies for the Amundsen Sea Embayment alone and as multi-year means. The 7 mm change is small relative to the S1 90% interval of [−10, 182] mm, so we fit the record as reported.

EAIS is listed as a diagnostic because the East Antarctic Ice Sheet is close to balance over the IMBIE-3 record, leaving little signal for a temperature-dependent model to recover. The fit is consistent with the reported errors, but the projection adopts a literature surface-mass-balance sensitivity, $C_T = 60 \pm 20$ Gt yr^{-1} °C^{-1} (*Frieler et al.*, 2015; *Ligtenberg et al.*, 2013), and uses the regression only as a check on that value.

One further fit enters the framework without being a component model in its own right. The linear surface-to-ocean transfer function given in the main text, which carries SSP temperature trajectories into the Greenland discharge model, explains a modest fraction of the interannual variance, $R^2 = 0.35$ over 51 common annual values, with a residual standard deviation of 0.20 °C that is propagated into the projections. It is the weakest statistical link in the Greenland projection chain, and it is retained rather than replaced because the quantity it must supply is the multi-decadal trend in subsurface temperature rather than its year-to-year variability. Greenland SMB is passed through as the RACMO-derived *Mouginot et al.* (2019) SMB over the historical period and projected with the emulator of MARv3.12 described in Section 1.4, whose response to warming is not fitted to observations. Its level and its slow departures from the emulator are fitted to the *Mouginot et al.* (2019) SMB through the drift state. The emulator's out-of-sample test is the leave-one-GCM-out comparison in Fig. S7, and the drift state's are the forecast tests in Section 1.4.

These are all within-sample statistics, and on their own they establish only that each model is consistent with the record it was fitted to. The tests that bear on forecasting skill are out of sample. The thermosteric model is compared against the *Frederikse et al.* (2020) steric reconstruction, withheld from calibration, with an RMS difference of 3.4 mm over 1957–2018 and 2.4 mm before 1957, where that reconstruction is itself constrained by models rather than by subsurface observations. When the thermosteric model is refitted with the 0–700 m record before 1990 withheld, retaining the full 0–2000 m record, which begins in 2005, it gives $b_\mathrm{u} = 52.9$ [46.9, 69.0] mm °C^{-1}, compared with 51.8 [43.8, 70.0] mm °C^{-1} from the full record. It predicts a rate of 0.24 [0.17, 0.29] mm yr^{-1} over the withheld years 1955–1989, compared with 0.20 [0.16, 0.24] mm yr^{-1} observed. In the prediction, the withheld years lie 15.0 [12.0, 17.6] mm below the 1995–2005 mean, compared with 12.0 [11.4, 12.6] mm observed (Fig. S10). GMST rose by only 0.3 to 0.4 °C over the withheld years, so this test constrains the level of the fit more than its sensitivity to warming. The early record also constrains the curvature of the response. Without it, $a_\mathrm{u} = 9.1$ [1.1, 23.6] mm °C^{-2}, compared with 18.8 [12.0, 27.8] mm °C^{-2} from the full record, so the projections use the full-record fit. The glacier model is compared against the global glacier mass changes of *Zemp et al.* (2019) for 1962–1999, withheld from calibration, and gives a mean rate of 0.30 [0.17, 0.40] mm yr^{-1}, compared with an observed 0.32 [−0.02, 0.66] mm yr^{-1}. For this comparison the posterior rate $b_\mathrm{gl}T + c_\mathrm{gl}$ is evaluated for each hydrological year (October to September) with T weighted 0.25 from the preceding calendar year and 0.75 from the current one. The uncertainties of *Zemp et al.* (2019), reported as 95% intervals, are converted to standard deviations and combined following their prescription, with the glaciological, interpolation, and area terms added in quadrature across years and the geodetic term added linearly. Like the GlaMBIE global total, their record includes the Greenland and Antarctic peripheral glaciers. Fitted to 1962–1989 alone, the *Zemp et al.* (2019) record shows no detectable sensitivity to GMST, unlike 1990–2016, and neither a delayed glacier response nor a quadratic dependence on GMST accounts for the difference. Greenland discharge is compared against the *Mankoff et al.* (2021) record (1986–2023) and the IMBIE-3 time series (*Otosaka et al.*, 2026), neither of which enters the fit. The model falls within error of both for discharge and for total mass balance. As noted in the main text, the *Mankoff et al.* (2021) comparison is the weaker of the two, because that record shares its underlying raw data with the *Mouginot et al.* (2019) calibration and differs from it in processing rather than in provenance. IMBIE-3 is

the more informative test. It combines mass-balance estimates from satellite observations of ice flow, surface elevation, and gravimetry. Its input-output estimates, which are built from ice flow, include *Mouginot et al.* (2019) and *Mankoff et al.* (2021). Its altimetry estimates (from 1992) and gravimetry estimates (from 2002) are independent of the calibration, so the independent part of the test is the total mass balance after 1992. When the discharge model is refitted to the 1990–2018 part of the *Mouginot et al.* (2019) record alone, it gives $\gamma = 0.365$ [0.302, 0.426] mm SLE $\mathrm{yr}^{-1}\,^{\circ}\mathrm{C}^{-1}$ and predicts 15.14 [14.86, 15.41] mm SLE of cumulative discharge over the withheld years 1978–1990, compared with 15.23 [14.96, 15.49] mm SLE observed. The tests for the framework as a whole rather than one component at a time are budget closure, in which the summed components are compared with satellite altimetry measurements that no component calibration saw, and the hindcast of the level, rate, and transient sensitivity of GMSL since 1950. Both are presented in the Results of the main text.




Figure S10: **Early-record holdout of the two-layer thermosteric model. (a)** NOAA 0–700 m thermosteric sea level (circles, with 90% intervals of the total point error) with the model fitted to the full 1955–2025 record (red) and to 1990–2025 alone (blue), each shown as the posterior median and 90% credible interval. The model and the data are referenced to their 1995–2005 means, and the years withheld from the 1990–2025 fit are shaded. The 1990–2025 fit is dashed over the withheld years. **(b)** The same quantities relative to the median of the full time-series fit. The bands show parameter uncertainty only and do not include the fitted observation noise σ_{extra}.

Table S3: Goodness-of-fit for the component models fitted to data. For glaciers, the Peninsula, EAIS, and WAIS, χ^2/dof is evaluated against the full covariance of the cumulative record. For the thermosteric model it is normalized by the number of data points rather than by the degrees of freedom and includes the fitted σ_{extra} in the point variance. For Greenland discharge it is evaluated on the annual increments of the cumulative record, which is equivalent to using its full covariance, before the error scale is applied. Dataset references are given in Table S1. Greenland SMB, whose emulator is fitted to regional climate model output rather than to observations (Section 1.4), and terrestrial water storage are omitted.

Component	Calibration data	Reduced χ^2	Model
Thermosteric (0–700 m)	NOAA, 1955–2025 (71)	1.13	Two-layer, 5 fitted
Thermosteric (0–2000 m)	NOAA, 2005–2025 (21)	0.37	Joint with 0–700 m
Glaciers	GlaMBIE, 2000–2023 (24)	1.05	Linear rate–T, 1850 balance, σ_{extra}
Greenland discharge	Mouginot, 1978–2018 (41)	1.45	Delay, 2 fitted
Antarctic Peninsula	IMBIE-3, 1979–2023 (45)	0.97	Linear rate–T
EAIS (diagnostic)	IMBIE-3, 1979–2023 (45)	0.94	Linear rate–T
WAIS	IMBIE-3, 1979–2023 (45)	0.99	Linear rate in time, σ_{extra}

2 Exceedance Probability by Threshold and Scenario

Table S4 reports the probability of exceeding selected global mean sea-level thresholds (relative to pre-industrial) by 2100 under three warming trajectories, for the slow-WAIS case, $p(\text{S1}) = 0.90$, and the fast-WAIS case, $p(\text{S1}) = 0.10$. Together these columns isolate the contribution of WAIS tail risk: the difference between them is the effect of shifting the scenario weight from very unlikely to very likely continued mass loss along the observed trajectory. The slow-WAIS column retains 10% weight on the accelerating branch, so probabilities there fall but do not vanish.

The tabulated values are the fractions of posterior samples above each threshold. Fig. 10b of the main text plots the same quantity as a continuous curve over sea level. Those curves are smoothed for display only, by replacing each sample with a Gaussian kernel whose width is one quarter of Scott's rule for that ensemble, about 24 mm for the fast-WAIS case and 6 to 7 mm for the slow-WAIS case. At every threshold and warming trajectory in Table S4, the smoothed curves differ from the unrounded sample fractions by at most 0.6 percentage points.

Table S4: **Probability (%) of exceeding global mean sea-level thresholds by 2100.** Thresholds are relative to the pre-industrial baseline (≈1850–1900). The $p(\mathrm{S1}) = 90\%$ column is the slow-WAIS case, which keeps 10% weight on the accelerating branch; the $p(\mathrm{S1}) = 10\%$ column is the fast-WAIS case. Values are computed from posterior samples under each warming trajectory.

SLR threshold (m, pre-ind.)	Warming trajectory	SPECTRUM p(s1)=90%	SPECTRUM p(s1)=10%
0.5	1.8 °C	94.1	99.2
	2.7 °C	99.9	100.0
	3.6 °C	100.0	100.0
0.75	1.8 °C	7.5	49.7
	2.7 °C	50.7	84.3
	3.6 °C	95.6	99.0
1.0	1.8 °C	2.5	23.0
	2.7 °C	4.2	33.6
	3.6 °C	29.7	67.3
1.5	1.8 °C	0.9	7.8
	2.7 °C	1.1	10.1
	3.6 °C	1.7	15.3
2.0	1.8 °C	0.4	3.5
	2.7 °C	0.4	3.9
	3.6 °C	0.6	5.5

3 Value of Information Analysis

The value of information (VoI) analysis quantifies the economic benefit of resolving the binary uncertainty about whether WAIS retreat remains slow (Scenario 1) or allows for the possibility of rapid retreat (Scenario 2). We compute the expected value of perfect information (EVPI), the perfect-information limit of the sample-information framework of *Raiffa and Schlaifer* (1961). No signal model is specified, so any test that discriminates the two scenarios imperfectly would return less than the values reported here.

3.1 Decision framework

We consider an illustrative coastal adaptation decision where a planner must choose a defense height d (meters of sea-level protection) to minimize total expected costs, evaluated against the sea-level rise distribution S at a representative target year of 2100. For a given sea-level rise distribution S at 2100, the total annual cost of defense height d is

$$C(d) = C_{\mathrm{adapt}}(d) + \mathbb{E}\big[D\big(\max(S - d,\, 0)\big)\big], \tag{2}$$

where $C_{\mathrm{adapt}}(d) = c_{\mathrm{a}} \cdot d$ is the annualized adaptation cost and $\mathbb{E}\big[D(\cdot)\big]$ is the expected value of the residual damage D from SLR exceeding the defense height (overtopping).

3.1.1 Adaptation cost

The total capital cost of coastal adaptation is estimated at $C_{\text{cap}} = \$4\,\text{T/m}$ of GMSL rise. This is derived from Tiggeloven et al. (2020), who estimate that 3.4% of the global coastline (∼54,400 km of ∼1.6 million km total) warrants dike heightening under cost-benefit optimality, at a unit cost of \$7 M per km per meter of dike height (the median cost raising dikes in New Orleans post-Hurricane Katrina). This yields ∼\$400 B/m for dike raising alone along the optimal 3.4% of coastline. We apply a conservative 10× multiplier to account for the full societal adaptation burden—including seawalls, nature-based solutions, managed retreat, and infrastructure relocation—across a broader fraction of the global coastline.

Because the cost function $C(d)$ (Eq. 2) combines adaptation costs with annual flood damages from Jevrejeva et al. (2018), both terms need to be expressed in consistent units. Adaptation is dominated by capital investment deployed over the planning horizon, so we conservatively annualize C_{cap} using the capital recovery factor (CRF):

$$c_{\text{a}} = C_{\text{cap}} \times \text{CRF}, \qquad \text{CRF} = \frac{r\,(1+r)^n}{(1+r)^n - 1}, \tag{3}$$

where $r = 0.03$ is the real discount rate and $n = 75\,\text{yr}$ is the planning horizon. This gives CRF ≈ 0.034 and an annualized adaptation cost of $c_{\text{a}} \approx \$135\,\text{B/(m·yr)}$ of GMSL rise. The resulting $C(d)$ is in \$/yr, the EVPI is in \$/yr, and the present-value conversion (Eq. 7) produces a present-value total in \$.

3.1.2 Damage function

Residual flood damages are computed from the estimates of Jevrejeva et al. (2018), which relate GMSL rise to annual global flood costs without additional adaptation, based on estimated asset exposure. We interpolate linearly between six points. Four are values stated in their text, \$10.2, 11.7, 14.3, and 27.0 T/yr at 0.52, 0.63, 0.86, and 1.80 m of rise. One is zero cost at zero rise. The sixth, \$2.5 T/yr at 0.20 m, is read from their Fig. 4a, where the curves for their 1.5 °C, 2 °C, and RCP8.5 median scenarios pass through \$2.3–2.7 T/yr at that level of rise. Beyond 1.80 m we extrapolate linearly using the slope of the final segment, \$13.5 T/yr per meter.

3.2 Two-world structure

The forecast distribution is decomposed into two conditional distributions, conditioned directly on WAIS scenario membership rather than on a percentile-threshold proxy:

- **The S1 world** (S_{S1}) samples from the blended forecast conditional on WAIS Scenario 1, representing continued mass loss along the observed trajectory.
- **The S2 world** (S_{S2}) samples from the full forecast conditional on WAIS Scenario 2, representing outcomes that allow for rapid WAIS retreat.

Because the WAIS mixture comprises exactly these two scenarios, the two worlds partition it completely, but they are built differently. Every Monte Carlo sample's WAIS scenario assignment is recorded at the point of sampling and carried through the component-summation

and rate-space-blending pipeline as a per-sample label, and S_{S2} is the 1,805 samples of the blended forecast carrying the S2 label, an exact conditional subset rather than a percentile-threshold proxy. S_{S1} is not the 195 samples carrying the S1 label, which would be too few to resolve the tail; it is a separate 2,000-member ensemble in which WAIS is replaced by its S1 trajectory for every sample, with the non-WAIS components unchanged. The two branches therefore have different sample sizes, 2,000 and 1,805.

The VoI takes these two conditional worlds as its inputs and recombines them below with the prior p_{s} as the weight, so the slow- and fast-WAIS cases of the main text, at $p(\mathrm{S1}) = 0.90$ and 0.10, are mixtures of these worlds rather than inputs to the calculation. Because the value depends on the prior, we report it as a function of p_{s} rather than at a single value.

3.3 EVPI calculation

For a given prior $p_{\mathrm{s}} \equiv p(\mathrm{S1})$ on the S1 world, the EVPI is the annual benefit of learning which scenario holds before choosing the defense height for coastal development.

3.3.1 Hedging cost (no information)

Without additional information, the planner hedges by choosing the single defense height d^*_{hedge} that minimizes the prior-weighted expected cost:

$$C^*_{\mathrm{hedge}} = \min_d \left[\, p_{\mathrm{s}} \cdot C_{\mathrm{S1}}(d) + (1 - p_{\mathrm{s}}) \cdot C_{\mathrm{S2}}(d) \,\right], \tag{4}$$

where $C_{\mathrm{S1}}(d) = C_{\mathrm{adapt}}(d) + \mathbb{E}[D(\max(S_{\mathrm{S1}} - d, 0))]$ and $C_{\mathrm{S2}}(d)$ is defined analogously using S_{S2}.

3.3.2 Perfect information cost

With perfect information, the planner would tailor the defense height to whichever world is realized:

$$C^*_{\mathrm{perfect}} = p_{\mathrm{s}} \cdot \min_d C_{\mathrm{S1}}(d) \;+\; (1 - p_{\mathrm{s}}) \cdot \min_d C_{\mathrm{S2}}(d). \tag{5}$$

3.3.3 Annual EVPI

The annual value of perfect information is the difference between the hedging cost and the perfect-information cost:

$$\mathrm{EVPI}_{\mathrm{ann}} = C^*_{\mathrm{hedge}} - C^*_{\mathrm{perfect}}. \tag{6}$$

This is the annual cost penalty of having to compromise on a single design rather than tailoring to the true state. It is non-negative by construction, because C^*_{hedge} minimizes a weighted sum of the two cost curves while C^*_{perfect} minimizes each of them separately. As a minimum over d of functions affine in p_{s}, C^*_{hedge} is concave in p_{s}, and subtracting the affine C^*_{perfect} leaves a concave function that vanishes at $p_{\mathrm{s}} = 0$ and $p_{\mathrm{s}} = 1$, so the maximum is interior. Its location is set by the geometry of the two cost curves and need not be at $p_{\mathrm{s}} = 0.5$. Here the two worlds are strongly asymmetric, with optimal designs of 0.85 m under S1 against 3.10 m under S2, and the maximum falls at $p_{\mathrm{s}} = 0.74$.

3.4 Discounting and the VoI surface

The annual EVPI is converted to a present value at the year of arrival by discounting the stream of benefits from the year information arrives (t_a) to the target year ($t_\mathrm{T} = 2100$) at a real discount rate $r = 3\%$:

$$\mathrm{VoI}(t_\mathrm{a}, p_\mathrm{s}) = \mathrm{EVPI}_\mathrm{ann}(p_\mathrm{s})\ \max\left(0,\ \frac{1-(1+r)^{-(t_\mathrm{T}-t_\mathrm{a})}}{r}\right). \tag{7}$$

The annuity factor $[1-(1+r)^{-n}]/r$ captures the number of discounted years over which the improved decision yields benefits.

The VoI surface (Fig. 11c of the main text) is computed over a grid of arrival years ($t_\mathrm{a} \in [2026, 2150]$) and prior probabilities ($p_\mathrm{s} \in [0.01, 0.99]$). All calculations use the 2.7 °C warming trajectory, the scenario closest to current policies (*Hausfather*, 2025). The choice of warming trajectory has qualitatively little impact on the results given all the uncertainties involved. Note that our cost estimates are likely to be underestimates (see §3.1.1), but the takeaway message remains even if the true costs or damages are an order of magnitude lower. The main text quotes the maximum of this surface over both axes, US$4.7 trillion, which is attained at $p_\mathrm{s} = 0.74$ for information arriving in 2026. For the same arrival year it also quotes the values at the priors of the two storylines, US$3.9 trillion at $p_\mathrm{s} = 0.90$ (slow-WAIS) and US$0.9 trillion at $p_\mathrm{s} = 0.10$ (fast-WAIS).

3.5 Key assumptions and limitations

The VoI calculation is intended to be simple analysis to make a simple point, not to stand for detailed financial decisions. As such, we make several simplifying assumptions:

1. **Binary resolution.** Information perfectly resolves the slow/fast dichotomy. Partial information (e.g., updating p_s from 0.10 to 0.30 without full resolution) would yield lower VoI.

2. **Single decision.** The planner makes one irreversible design choice. In practice, adaptive management with staged decisions would reduce the hedging penalty, lowering VoI.

3. **Global aggregation.** Costs are aggregated globally. Local VoI would differ based on coastal exposure and elevation.

4. **Linear adaptation cost.** We assume C_adapt scales linearly with defense height. Economies or diseconomies of scale would modify the optimal designs.

5. **Discount rate.** The 3% real discount rate is standard for infrastructure but lower rates (reflecting intergenerational equity) would increase VoI. Higher discount rates, that would lower VoI, are unlikely given the long timeline of risk exposure from sea level rise.

Despite these simplifications, the analysis provides a rough order-of-magnitude estimate of the economic value of resolving WAIS uncertainty, supporting the argument that targeted investment in ice-sheet observations and process understanding has large expected returns.

References

Beckley, B. D., P. S. Callahan, I. Hancock, David W., G. T. Mitchum, and R. D. Ray (2017), On the "Cal-Mode" correction to TOPEX satellite altimetry and its effect on the global mean sea level time series, *Journal of Geophysical Research: Oceans*, *122*(11), 8371–8384, doi:10.1002/2017JC013090.

Boberg, F., X. Fettweis, N. Hansen, R. Mottram, and M. R. van den Broeke (2026), 21st century change in precipitation on the Greenland Ice Sheet using high resolution regional climate models, *The Cryosphere*, *20*, 2947–2960, doi:10.5194/tc-20-2947-2026.

Cheng, L., and J. Zhu (2016), Benefits of CMIP5 multimodel ensemble in reconstructing historical ocean subsurface temperature variations, *Journal of Climate*, *29*, 5393–5416, doi:10.1175/JCLI-D-15-0730.1.

Dangendorf, S., Q. Sun, T. Wahl, P. Thompson, J. X. Mitrovica, and B. Hamlington (2024), Probabilistic reconstruction of sea-level changes and their causes since 1900, *Earth System Science Data*, *16*(7), 3471–3494, doi:10.5194/essd-16-3471-2024.

Davison, B. J., A. E. Hogg, R. Rigby, S. Veldhuijsen, J. M. van Wessem, M. R. van den Broeke, P. R. Holland, H. L. Selley, and P. Dutrieux (2023), Sea level rise from West Antarctic mass loss significantly modified by large snowfall anomalies, *Nature Communications*, *14*, 1479, doi:10.1038/s41467-023-36990-3.

Desbruyères, D. G., S. G. Purkey, E. L. McDonagh, G. C. Johnson, and B. A. King (2016), Deep and abyssal ocean warming from 35 years of repeat hydrography, *Geophysical Research Letters*, *43*, 10,356–10,365, doi:10.1002/2016GL070413.

Enderlin, E. M., and I. M. Howat (2013), Submarine melt rate estimates for floating termini of Greenland outlet glaciers (2000–2010), *Journal of Glaciology*, *59*(213), 67–75, doi:10.3189/2013JoG12J049.

Fettweis, X., B. Franco, M. Tedesco, J. H. van Angelen, J. T. M. Lenaerts, M. R. van den Broeke, and H. Gallée (2013), Estimating the Greenland ice sheet surface mass balance contribution to future sea level rise using the regional atmospheric climate model MAR, *The Cryosphere*, *7*, 469–489, doi:10.5194/tc-7-469-2013.

Frederikse, T., F. Landerer, L. Caron, S. Adhikari, D. Parkes, V. W. Humphrey, S. Dangendorf, P. Hogarth, L. Zanna, L. Cheng, and Y.-H. Wu (2020), The causes of sea-level rise since 1900, *Nature*, *584*, 393–397, doi:10.1038/s41586-020-2591-3.

Frieler, K., P. U. Clark, F. He, C. Buizert, R. Reese, S. R. M. Ligtenberg, M. R. van den Broeke, R. Winkelmann, and A. Levermann (2015), Consistent evidence of increasing Antarctic accumulation with warming, *Nature Climate Change*, *5*(4), 348–352, doi:10.1038/nclimate2574.

Garner, G. G., T. H. J. Hermans, R. E. Kopp, A. B. A. Slangen, T. L. Edwards, A. Levermann, S. Nowicki, M. D. Palmer, C. Smith, B. Fox-Kemper, H. T. Hewitt, C. Xiao,

G. Aðalgeirsdóttir, S. S. Drijfhout, N. R. Golledge, M. Hemer, G. Krinner, A. Mix, D. Notz, I. S. Nurhati, L. Ruiz, J.-B. Sallée, Y. Yu, L. Hua, T. Palmer, and B. Pearson (2023), IPCC AR6 sea level projections, *Zenodo*, doi:10.5281/zenodo.6382554, version 20230209.

Geoffroy, O., D. Saint-Martin, D. J. L. Olivié, A. Voldoire, G. Bellon, and S. Tytéca (2013), Transient climate response in a two-layer energy-balance model. Part I: Analytical solution and parameter calibration using CMIP5 AOGCM experiments, *Journal of Climate*, *26*, 1841–1857, doi:10.1175/JCLI-D-12-00195.1.

Glaude, Q., B. Noël, M. Olesen, M. R. van den Broeke, W. J. van de Berg, R. Mottram, N. Hansen, A. Delhasse, C. Amory, C. Kittel, H. Goelzer, and X. Fettweis (2024), A factor two difference in 21st-century Greenland Ice Sheet surface mass balance projections from three regional climate models under a strong warming scenario (SSP5-8.5), *Geophysical Research Letters*, *51*(22), e2024GL111,902, doi:10.1029/2024GL111902.

Good, S. A., M. J. Martin, and N. A. Rayner (2013), EN4: Quality controlled ocean temperature and salinity profiles and monthly objective analyses with uncertainty estimates, *Journal of Geophysical Research: Oceans*, *118*, 6704–6716, doi:10.1002/2013JC009067.

Hamlington, B. D., A. S. Gardner, E. R. Ivins, J. T. M. Lenaerts, J. T. Reager, D. S. Trossman, E. D. Zaron, S. Adhikari, A. Arendt, A. Aschwanden, B. D. Beckley, D. P. S. Bekaert, G. Blewitt, L. Caron, D. P. Chambers, H. A. Chandanpurkar, K. Christianson, B. Csatho, R. I. Cullather, R. M. DeConto, J. T. Fasullo, T. Frederikse, J. T. Freymueller, D. M. Gilford, M. Girotto, W. C. Hammond, R. Hock, N. Holschuh, R. E. Kopp, F. Landerer, E. Larour, D. Menemenlis, M. Merrifield, J. X. Mitrovica, R. S. Nerem, I. J. Nias, V. Nieves, S. Nowicki, K. Pangaluru, C. G. Piecuch, R. D. Ray, D. R. Rounce, N.-J. Schlegel, H. Seroussi, M. Shirzaei, W. V. Sweet, I. Velicogna, N. Vinogradova, T. Wahl, D. N. Wiese, and M. J. Willis (2020), Understanding of contemporary regional sea-level change and the implications for the future, *Reviews of Geophysics*, *58*, e2019RG000,672, doi:10.1029/2019RG000672.

Hamlington, B. D., A. Bellas-Manley, J. K. Willis, S. Fournier, N. Vinogradova, R. S. Nerem, C. G. Piecuch, P. R. Thompson, and R. E. Kopp (2024), The rate of global sea level rise doubled during the past three decades, *Communications Earth & Environment*, *5*, 601, doi:10.1038/s43247-024-01761-5.

Hausfather, Z. (2025), An assessment of current policy scenarios over the 21st century and the reduced plausibility of high-emissions pathways, *Dialogues on Climate Change*, *2*(1), 26–32, doi:10.1177/29768659241304854.

Ishii, M., Y. Fukuda, S. Hirahara, S. Yasui, T. Suzuki, and K. Sato (2017), Accuracy of global upper ocean heat content estimation expected from present observational data sets, *Scientific Online Letters on the Atmosphere*, *13*, 163–167, doi:10.2151/sola.2017-030.

Jevrejeva, S., L. P. Jackson, A. Grinsted, D. Lincke, and B. Marzeion (2018), Flood damage costs under the sea level rise with warming of 1.5 °C and 2 °C, *Environmental Research Letters*, *13*(7), 074,014, doi:10.1088/1748-9326/aacc76.

Kjeldsen, K. K., N. J. Korsgaard, A. A. Bjørk, S. A. Khan, J. E. Box, S. Funder, N. K. Larsen, J. L. Bamber, W. Colgan, M. van den Broeke, M.-L. Siggaard-Andersen, C. Nuth, A. Schomacker, C. S. Andresen, E. Willerslev, and K. H. Kjær (2015), Spatial and temporal distribution of mass loss from the Greenland Ice Sheet since AD 1900, *Nature*, *528*, 396–400, doi:10.1038/nature16183.

Lee, J.-Y., J. Marotzke, G. Bala, L. Cao, S. Corti, J. P. Dunne, F. Engelbrecht, E. Fischer, J. C. Fyfe, C. Jones, A. Maycock, J. Mutemi, O. Ndiaye, S. Panickal, and T. Zhou (2021), Future global climate: scenario-based projections and near-term information, in *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the IPCC*, Cambridge University Press, doi:10.1017/9781009157896.006.

Levitus, S., J. I. Antonov, T. P. Boyer, O. K. Baranova, H. E. Garcia, R. A. Locarnini, A. V. Mishonov, J. R. Reagan, D. Seidov, E. S. Yarosh, and M. M. Zweng (2012), World ocean heat content and thermosteric sea level change (0–2000 m), 1955–2010, *Geophysical Research Letters*, *39*, L10,603, doi:10.1029/2012GL051106.

Li, Y., J. A. Church, T. J. McDougall, and P. M. Barker (2022), Sensitivity of observationally based estimates of ocean heat content and thermal expansion to vertical interpolation schemes, *Geophysical Research Letters*, *49*, e2022GL101,079, doi:10.1029/2022GL101079.

Ligtenberg, S. R. M., W. J. van de Berg, M. R. van den Broeke, J. G. L. Rae, and E. van Meijgaard (2013), Future surface mass balance of the Antarctic ice sheet and its influence on sea level change, simulated by a regional atmospheric climate model, *Climate Dynamics*, *41*(3–4), 867–884, doi:10.1007/s00382-013-1749-1.

Mankoff, K. D., X. Fettweis, P. L. Langen, M. Stendel, K. K. Kjeldsen, N. B. Karlsson, B. Noël, M. R. van den Broeke, A. Solgaard, W. Colgan, J. E. Box, S. B. Simonsen, M. D. King, A. P. Ahlstrøm, S. B. Andersen, and R. S. Fausto (2021), Greenland ice sheet mass balance from 1840 through next week, *Earth System Science Data*, *13*, 5001–5025, doi:10.5194/essd-13-5001-2021.

Mouginot, J., E. Rignot, A. A. Bjørk, M. van den Broeke, R. Millan, M. Morlighem, B. Noël, B. Scheuchl, and M. Wood (2019), Forty-six years of Greenland Ice Sheet mass balance from 1972 to 2018, *Proceedings of the National Academy of Sciences*, *116*(19), 9239–9244, doi:10.1073/pnas.1904242116.

Nerem, R. S., T. Frederikse, and B. D. Hamlington (2022), Extrapolating empirical models of satellite-observed global mean sea level to estimate future sea level change, *Earth's Future*, *10*, e2021EF002,290, doi:10.1029/2021EF002290.

Noël, B., L. van Kampenhout, J. T. M. Lenaerts, W. J. van de Berg, and M. R. van den Broeke (2021), A 21st century warming threshold for sustained Greenland ice sheet mass loss, *Geophysical Research Letters*, *48*, e2020GL090,471, doi:10.1029/2020GL090471.

Otosaka, I. N., A. Shepherd, C. Amory, M. Horwath, E. R. Ivins, M. D. King, S. Nowicki, A. J. Payne, E. Rignot, L. S. Sørensen, N.-J. Schlegel, K. M. Simon, B. E. Smith, T. C. Sutterley, M. R. Van Den Broeke, I. Velicogna, G. A, C. Agosta, P. Ditmar, T. Döhne, M. E. Engdahl, X. Fettweis, R. Forsberg, A. S. Gardner, L. Gilbert, H. Goelzer, N. Gourmelen, A. Groh, N. Hansen, C. Harig, V. Helm, S. A. Khan, C. Kittel, P. L. Langen, M. Larsen, B. D. Loomis, M. McMillan, B. Medley, D. Melini, R. H. Mottram, A. Muir, J. Nilsson, B. Noël, M. E. Pattle, M. Roca I Aparici, I. Sasgen, H. V. Save, B. Scheuchl, E. J. O. Schrama, L. Schröder, K.-W. Seo, S. B. Simonsen, T. Slater, G. Spada, B. D. Vishwakarma, N. Wever, D. N. Wiese, and B. Wouters (2026), Mass balance of the Greenland and Antarctic ice sheets from the 1970s to 2023, *Scientific Data*, *13*, 1301, doi: 10.1038/s41597-026-08088-0.

Raiffa, H., and R. Schlaifer (1961), *Applied Statistical Decision Theory*, Division of Research, Graduate School of Business Administration, Harvard University, Boston.

Rantanen, M., A. Y. Karpechko, A. Lipponen, K. Nordling, O. Hyvärinen, K. Ruosteenoja, T. Vihma, and A. Laaksonen (2022), The Arctic has warmed nearly four times faster than the globe since 1979, *Communications Earth & Environment*, *3*, 168, doi:10.1038/s43247-022-00498-3.

Rignot, E., Y. Xu, D. Menemenlis, J. Mouginot, B. Scheuchl, X. Li, M. Morlighem, H. Seroussi, M. van den Broeke, I. Fenty, C. Cai, L. An, and B. de Fleurian (2016), Modeling of ocean-induced ice melt rates of five west Greenland glaciers over the past two decades, *Geophysical Research Letters*, *43*, 6374–6382, doi:10.1002/2016GL068784.

Robel, A. A., H. Seroussi, and G. H. Roe (2019), Marine ice sheet instability amplifies and skews uncertainty in projections of future sea-level rise, *Proceedings of the National Academy of Sciences*, *116*(30), 14,887–14,892, doi:10.1073/pnas.1904822116.

Roe, G. H., J. E. Christian, and B. Marzeion (2021), On the attribution of industrial-era glacier mass loss to anthropogenic climate change, *The Cryosphere*, *15*, 1889–1905, doi: 10.5194/tc-15-1889-2021.

Rohde, R. A., and Z. Hausfather (2020), The Berkeley Earth land/ocean temperature record, *Earth System Science Data*, *12*, 3469–3479, doi:10.5194/essd-12-3469-2020.

Rounce, D. R., R. Hock, F. Maussion, R. Hugonnet, W. Kochtitzky, M. Huss, E. Berthier, D. Brinkerhoff, L. Compagno, L. Copland, D. Farinotti, B. Menounos, and R. W. McNabb (2023), Global glacier change in the 21st century: Every increase in temperature matters, *Science*, *379*, 78–83, doi:10.1126/science.abo1324.

Slater, D. A., and F. Straneo (2022), Submarine melting of glaciers in Greenland amplified by atmospheric warming, *Nature Geoscience*, *15*(10), 794–799, doi:10.1038/s41561-022-01035-9.

The GlaMBIE Team, M. Zemp, L. Jakob, I. Dussaillant, S. U. Nussbaumer, N. Gourmelen, S. Dubber, G. A, S. Abdullahi, L. M. Andreassen, E. Berthier, A. Bhattacharya,

A. Blazquez, L. F. Boehm Vock, T. Bolch, J. Box, M. H. Braun, F. Brun, E. Cicero, W. Colgan, N. Eckert, D. Farinotti, C. Florentine, D. Floricioiu, A. Gardner, C. Harig, J. Hassan, R. Hugonnet, M. Huss, T. Jóhannesson, C.-C. A. Liang, C.-Q. Ke, S. A. Khan, O. King, M. Kneib, L. Krieger, F. Maussion, E. Mattea, R. McNabb, B. Menounos, E. Miles, G. Moholdt, J. Nilsson, F. Pálsson, J. Pfeffer, L. Piermattei, S. Plummer, A. Richter, I. Sasgen, L. Schuster, T. Seehaus, X. Shen, C. Sommer, T. Sutterley, D. Treichler, I. Velicogna, B. Wouters, H. Zekollari, and W. Zheng (2025), Community estimate of global glacier mass changes from 2000 to 2023, *Nature*, *639*, 382–388, doi:10.1038/s41586-024-08545-z, glaMBIE consortium.

Tiggeloven, T., H. de Moel, H. C. Winsemius, D. Eilander, G. Erkens, E. Gebremedhin, A. Diaz Loaiza, A. Bouwman, J. van Huijstee, W. Ligtvoet, S. Kuzma, T. Luo, C. Iceland, and P. J. Ward (2020), Global-scale benefit–cost analysis of coastal flood adaptation to different flood risk drivers using structural measures, *Natural Hazards and Earth System Sciences*, *20*, 1025–1044, doi:10.5194/nhess-20-1025-2020.

Wood, M., E. Rignot, I. Fenty, L. An, A. Bjork, M. Van Den Broeke, C. Cai, E. Kane, D. Menemenlis, R. Millan, M. Morlighem, J. Mouginot, B. Noël, B. Scheuchl, I. Velicogna, J. K. Willis, and H. Zhang (2021), Ocean forcing drives glacier retreat in Greenland, *Science Advances*, *7*(1), eaba7282, doi:10.1126/sciadv.aba7282.

Xu, Y., E. Rignot, I. Fenty, D. Menemenlis, and M. Flexas (2013), Subaqueous melting of Store Glacier, west Greenland from three-dimensional, high-resolution numerical modeling and ocean observations, *Geophysical Research Letters*, *40*, 4648–4653, doi:10.1002/grl.50825.

Zemp, M., M. Huss, E. Thibert, N. Eckert, R. McNabb, J. Huber, M. Barandun, H. Machguth, S. U. Nussbaumer, I. Gärtner-Roer, L. Thomson, F. Paul, F. Maussion, S. Kutuzov, and J. G. Cogley (2019), Global glacier mass changes and their contributions to sea-level rise from 1961 to 2016, *Nature*, *568*(7752), 382–386, doi:10.1038/s41586-019-1071-0.