

Orthogonal risks in 21st-century sea-level rise: independent roles for emissions reduction and ice-sheet stabilization

Minchew et al.

Abstract

Sea-level projections are conventionally presented as functions of emission scenario, implying that emissions reduction is the primary lever for managing coastal risk. We show that this framing obscures the dominant source of uncertainty. Using a data-driven Bayesian forecast calibrated against independent observations of each sea-level contributor, we project 1.0–1.3 m of global mean sea-level rise by 2100 (medians across policy-relevant pathways relative to pre-industrial), exceeding IPCC AR6 estimates by 70–79%. A variance decomposition reveals that 92% of total forecast uncertainty originates from a single source—the dynamical response of the West Antarctic Ice Sheet (WAIS)—while emission-pathway differences account for only 8%. These two dimensions of risk are approximately orthogonal: emissions reduction compresses the predictable, temperature-sensitive components (ocean thermal expansion, glaciers, Greenland) but leaves the WAIS-dominated tail largely unchanged, while interventions targeting ice-sheet stability would collapse the dominant uncertainty without requiring further emissions cuts. Even under aggressive mitigation (2 °C trajectory), we find a 7% probability of exceeding 2 m; under 5 °C warming, this rises to only 19%, confirming that the tail risk is largely emission-insensitive. These results reframe sea-level risk management as a two-lever problem: emissions reduction to shift the central estimate, and ice-sheet research and intervention to reduce the catastrophic tail. Neither alone is sufficient; both are necessary.

Rates of global mean sea-level rise (GMSLR) have doubled twice in the past 50 years [1, 2] (Fig. 1). The first doubling took about 16 years and 0.34 °C of warming [3]. The second took about 21 years and 0.41 °C of warming (Fig. 1). The rate could double again around 2080 if the current observed sea-level acceleration continues, and by 2040 if it takes no more warming than the previous two doublings. The 40-year gap illustrates the uncertainty that coastal planning and adaptation must absorb.

Every centimeter of sea-level rise exposes roughly one million additional people to annual coastal flooding [7], approximately 90% of whom live in low- and middle-income countries [8]. Without additional adaptation, annual flood damages are projected to reach US\$14–18 trillion per year for 0.9–1.1 m of rise [9]. Estimates of total adaptation costs per unit sea level rise are not available but a lower-bound, rough-order-of-magnitude estimate of US\$4 trillion per meter of sea-level rise is reasonable based on a projected cost of US\$400 billion simply to raise dikes by 1 meter along only 3.4% of coastlines [10]. Developing countries currently

receive US\$21 billion per year in total adaptation finance across all sectors, not just coastal [11], raising serious questions about the feasibility of coastal adaptation at relevant scales and highlighting the extreme financial costs, to say nothing of human and ecological impacts, arising from large uncertainties.

The rate of reduction in the uncertainties is a crucial consideration because effective planning requires reliable sea-level forecasts be delivered before key decisions need to be made. Uncertainty in the rates of sea-level rise compounds uncertainty in inundation thresholds and paralyzes anticipatory spatial planning, routinely defaulting policy toward *in situ* maladaptations that compound sunk costs and strand assets [12], or delaying action until forced by disaster [13]. Unconstrained timelines encourage fragmented, market-based incrementalism that deploys public capital inefficiently [14] and systemically transfers the financial burden of reactive retreat onto vulnerable municipalities and demographics [15].

Much of the current coastal adaptation planning relies on projections from the IPCC Sixth Assessment Report (AR6), which estimates 0.4–0.7 m of rise by 2100 relative to the 1995–2014 baseline [16, 17]. These projections depend largely on process-based ice sheet models that diverge from satellite observations within years of initialization [18, 19]. As a result, observed GMSL trends (0.66 m median at 2100, 0.41–0.91 m 90% CI) already exceed AR6 medium-confidence projections under all policy-relevant emission scenarios [2, 20]. AR6 itself documents this in a single figure but without comment: extrapolating the satellite-era rate and acceleration—which reflect the current warming trajectory that could lead to 3 °C (SSP2-4.5)—produces sea-level rise that matches or slightly exceeds the process-based projections under the very high emissions scenario (SSP5-8.5), which is now regarded as generating an implausible level of warming (AR6 Figure 9.27; [16, 21]). So while the data show that rates of sea level rise double with approximately 0.4 °C of warming (Fig. 1b), IPCC models need roughly an order of magnitude more warming just to reproduce current observed rates of sea-level rise.

Disagreement between observations and ice sheet models arise from known structural biases, including the absence of leading-order physical processes like iceberg calving [19] and an assumed ice viscosity that is less sensitive to stress than observations indicate [22]. The latter produces a roughly 30% underestimate of projected 21st century ice loss, exceeding the uncertainty due to warming trajectories [23]. More broadly, decades of general forecasting research shows that models more complex than their calibration data can constrain produce worse predictions than simpler models [24, 25]. Current ice sheet models have millions of degrees of freedom representing spatially distributed parameters and uncertain parameterizations. The observations used to initialize these models inform only a small fraction of these degrees of freedom, with the remainder determined by prior assumptions and model design [26, 27]. A forecasting system based on models representing diverse approaches and a hierarchy of complexity is needed to fill the needs of coastal planning for sea-level rise.

Data-driven forecasting

We construct a data-driven forecast that focuses on different sea-level contributors, each with minimal physical models constrained by independent observational records spanning decades. Observed GMSLR and its sensitivity to observed global mean surface temperature (GMST) are diagnostics, and we show that the sum of our model components agrees within

6% of 2000–2025 observed GMSL in hindcasting tests (Supplementary Material). Unless otherwise noted, all sea-level values in the main text are global-mean 21st-century contributions referenced to the year 2000, while values in the abstract are relative to pre-industrial for consistency with existing projections and literature. Throughout, we label emission scenarios by their approximate end-of-century global mean surface temperature: 2 °C (SSP1-2.6), 3 °C (SSP2-4.5), and 4 °C (SSP3-7.0).

Our framework builds on two established lines of data-driven forecasting. The simplest is the “naive” forecast, which requires only a fit to the observed GMSL rate and acceleration. The satellite altimetry record provides GMSLR measurements from 1993–2025 that show a constant acceleration of 0.08 mm yr^{-2} ($0.02\text{--}0.14 \text{ mm yr}^{-2}$) and rate 4.5 mm yr^{-1} ($3.5\text{--}5.5 \text{ mm yr}^{-1}$) in 2025 [2, 20]. Extrapolating these values to 2100 yields 66 cm (41–91 cm) of 21st-century rise (Fig. 3a). Assuming the naive forecast has predictive skill over about half the calibration window, the extrapolation of the 32-year satellite record provides predictive skill over the next 15–20 years. Beyond that horizon, the naive forecast is less reliable but nevertheless establishes a scenario-independent, data-driven reference (Fig. 3a).

At the next level of complexity are the semi-empirical models that are based on the observed sensitivity of the rate of GMSLR to changes in GMST (Fig. 1b). Under current warming trends, Rahmstorf (2007) [5] predicts 0.67 m (0.63–0.71 m) of 21st-century rise, matching the extrapolation of modern observed rates within error [2]. An updated semi-empirical model [6], which adds a rate-of-warming term, predicts 0.96 m (0.80–1.12 m), in close agreement with the results of this study. Although physically motivated by the fit to observed sensitivities, semi-empirical models were assessed as *low confidence* by IPCC AR5 because they do not resolve individual contributing processes and because extrapolation beyond the calibration period assumes a stationary temperature–sea-level relationship [28]. This reasoning is carried forward by AR6 but violated by the decision to favor process models that omit key physical processes and were not tested against observed GMSL, the very quantity they claim to predict, prior to publication (cf. AR6 Figure 9.27 in [16]). Our own results show that the simplest semi-empirical models have much greater forecasting skill for the components of sea-level rise that are directly sensitive to warming.

Component model calibration

We decompose global mean sea level into separate components: ocean thermal expansion (thermosteric), Greenland Ice Sheet (GrIS) surface mass balance (SMB) and glacier discharge, West Antarctic Ice Sheet (WAIS) discharge, East Antarctic Ice Sheet (EAIS) SMB, Antarctic Peninsula total ice loss, glaciers (land ice not in the ice sheets), and terrestrial water storage. Our component-summation approach overlaps with the methodology of previous studies [29–31], while extending to more recent data, rate-space blending against satellite-era observations to ensure alignment between the forecast and observed data trends (Supplementary Materials), and a structured deep-uncertainty treatment of WAIS discharge. The sensitivities of glaciers, GrIS discharge, and Antarctic Peninsula to warming are calibrated with independent observational datasets spanning 24–71 years (Table S1). Surface mass balance for GrIS and EAIS are adopted from published model sensitivities rather than data calibration because they lack sufficient data coverage for our approach [32, 33]. For completeness, we adopt the IPCC projection of terrestrial water storage, a decision driven by a

lack of suitable data and its negligible contribution to 21st century GMSLR.

The thermosteric component—the largest single contributor to observed sea-level rise but a small contributor to forecast uncertainty—is modeled with a two-layer energy balance [36] fitted jointly to NOAA in situ thermosteric sea level measurements at 0–700 m (1957–2025) and 0–2000 m (2005–2025) [37] (Fig. 2a). In our model, the upper ocean responds to GMST warming through a thermal state variable S_u with relaxation time τ_u . The deep ocean relaxes toward S_u on a longer timescale $\tau_d = 150$ yr taken from the literature [36].

The Greenland Ice Sheet is currently the largest land-ice contributor [1] but a small contributor to forecast uncertainty. We treat SMB and ice discharge separately. SMB observations from a regional climate model (RCM), RACMO [34, 40], are used directly for the hindcast. The forecasts scale with temperature using sensitivities from regional climate model intercomparisons [32, 33]. Our choice of RCM does not materially affect our results, and we capture the structural uncertainty in the inter-RCM spread [43] in our forecast priors (Supplementary Material).

GrIS discharge is modeled as the cumulative response to time-delayed subsurface ocean warming, with a sensitivity γ , a delay δ between ocean warming and ice sheet response, and a secular drift r_0 (Supplementary Material). Cross-correlation of EN4 subsurface ocean temperature [44] against the Mouginot et al. [34] discharge record and Bayesian Information Criterion (BIC) model selection yields $\delta = 5$ yr (Fig. 2b). The fitted sensitivity $\gamma = 0.37 \text{ mm yr}^{-1} \text{ }^\circ\text{C}^{-1}$ agrees with physical scaling estimates from ocean and glacier dynamics ($0.36 \text{ mm yr}^{-1} \text{ }^\circ\text{C}^{-1}$). An independently derived discharge record (1986–2023) [45] and NOAA Arctic Report Card estimates [46], withheld from calibration, fall within the 90% credible interval.

Mountain glacier mass loss is modeled as a rate–temperature relationship calibrated against the GlaMBIE global consensus [2000–2023; 47]. The data only support a linear sensitivity (Supplementary Material), which we find to be $b_{\text{gl}} = 0.66 \text{ mm yr}^{-1} \text{ }^\circ\text{C}^{-1}$ (Fig. 2c). This is approximately five times the value embedded in IPCC AR6 median projections ($\sim 0.13 \text{ mm yr}^{-1} \text{ }^\circ\text{C}^{-1}$) and $1.3\text{--}1.7\times$ the effective sensitivity of the PyGEM process model [48], but is overall a small contributor to sea-level rise (Supplementary Materials).

We compare the sum of these independently calibrated components to observations of GMSL from satellite altimetry as a diagnostic (Fig. 2d,e). The diagnostic is possible because each component is calibrated against its own observational record. Total GMSL is not used as a calibration target. The component sum closes 90–94% of the observed GMSLR rate over 1993–2025, with a variance-weighted attribution assigning the residual mainly to the thermosteric (61%) and terrestrial water storage (21%) components, both within their stated uncertainties, and all components are within 90% CI of their independent estimates (Supplementary Material).

West Antarctic Ice Sheet. WAIS is by far the largest source of forecast uncertainty, accounting for 92% of within-emissions-scenario variance at 2100 (Figs. 2f and 3b). The ice sheet contains ~ 3.3 m of sea level equivalent on retrograde beds susceptible to internal feedbacks commonly called the marine ice sheet instability (MISI; [49]). Ocean thermal forcing sufficient to drive retreat is likely committed regardless of emissions pathway [50, 51] and may be driven as much by oceanographic processes local to the ice sheet as those further

afield [?]. Based on these lines of evidence, we make our WAIS projections independent of the warming trajectory, meaning the range of 21st-century outcomes is controlled by ice dynamics and, to first order, is no longer sensitive to emissions reductions.

No physical process models have been proven to produce reliable projections of WAIS (Fig. 2f). To provide a representative range of uncertainties, we construct a scenario-mixture framework with three outcomes that are grounded in the scientific literature (Supplementary Material). Scenario 1: ‘slow WAIS’ where rates of discharge continue to increase within the range of present-day rates, contributing 25–85 mm GMSLR by 2100. Based on the dramatic changes in WAIS discharge over the past decades, we take this scenario to be unlikely, and assign it a 10% probability. Scenario 2 represents the role of MISI-driven acceleration with grounding-line retreat on retrograde bedrock, with a GMSL contribution of 150–1000 mm by 2100. The lower bound is anchored to transient-calibrated ice sheet model projections [52] and the observed 33 km grounding-line retreat at Pine Island Glacier since 1992 [53] while the upper bound is set by structured expert judgment [56]. We assign Scenario 2 an 80% probability. Scenario 3 represents MISI combined with marine ice cliff instability and its values defined by the IPCC AR6 low-confidence 83rd (600 mm) and 95th (1300 mm) percentile ranges. We assign a 10% probability to this scenario. Within each scenario, the 2100 endpoint is drawn from a skew-normal distribution in log-space with physically motivated asymmetry [54]. A multiplicative rheology correction (R with median $R = 1.3$) is applied to all scenarios, reflecting the underestimation of ice loss when models assume the ice-viscosity stress exponent $n = 3$, which is contrary to the observationally constrained value $n = 4.1 \pm 0.4$ [22, 23, 55]. The resulting mixture has a median 21st century sea-level rise of 0.41 m with a 0.07 to 1.56 m 90% CI at 2100. Our choices of scenario weights have small quantitative impacts on our forecast results in the high-end tails of the distribution, but do not qualitatively impact the findings of this study (Supplementary Materials).

Forecasts

We focus on warming trajectories of 2 °C (SSP1-2.6) through 4 °C (SSP3-7.0), the pathways most consistent with current policies and stated pledges (Fig. 3). Under the 2 °C target, the blended forecast projects 0.78 m of 21st-century sea-level rise [0.44–1.92 m], which is 79% above the IPCC AR6 medium-confidence estimate of 0.44 m. Under 3 °C warming, approximately the current trajectory, the median is 0.95 m [0.61–2.08 m], 70% above IPCC’s 0.56 m. For the 4 °C trajectory, the median reaches 1.15 m [0.80–2.29 m]. For reference, the 5 °C warming trajectory (SSP5-8.5) yields a median of 1.34 m [0.98–2.48 m]. All sea-level values given relative to year 2000.

Relative to pre-industrial GMSL (adding ~ 0.19 m of observed rise through 2000), these forecasts correspond to medians of 0.97–1.34 m of global mean sea level rise across the policy-relevant pathways. The probability of exceeding 1 m relative to pre-industrial by 2100 is 46% under 2 °C warming, 74% under 3 °C, and 95% under 4 °C warming trajectory. Averaged across these pathways, the probability of exceeding 1 m of sea level rise relative to pre-industrial levels is 73%, and is largely insensitive to the relative weighting of scenarios (Fig. 3) because forecasting spread is driven by the dynamics of WAIS glaciers. If, on the other hand, the ‘slow WAIS’ scenario were true, the probability of exceeding 1 m of sea level rise relative to pre-industrial levels drops to zero if warming stays below 3 °C (Fig. 3b).

Orthogonal dimensions of sea-level risk

The within-scenario 90% uncertainty range (1.47 m for 3 °C) is four times larger than the spread across scenario medians (0.37 m from 2–4 °C). A variance decomposition (law of total variance across policy scenarios) attributes 92% of total forecast variance to within-scenario uncertainty—predominantly WAIS—and 8% to between-scenario differences. This is not merely a statement about the relative magnitude of uncertainties. It reveals that sea-level risk decomposes into two approximately orthogonal dimensions, each addressed by a distinct class of intervention.

The first dimension is *emission-sensitive*. The temperature-dependent components—ocean thermal expansion, glacier mass loss, Greenland surface mass balance and glacier discharge, and the Antarctic Peninsula—respond to global temperature through well-characterized physical relationships calibrated against decades of observations. Their collective contribution shifts systematically across warming trajectories and is directly reducible by emissions cuts. Conditional on a stable WAIS, the forecast narrows to a 90% range of approximately 140 mm, a level of precision comparable to the thermosteric projection alone (Fig. 3a, purple band). In this regime, emissions policy is the primary lever: the difference between 2 °C and 4 °C warming corresponds to roughly 200 mm of additional rise, nearly all of which is attributable to temperature-sensitive components.

The second dimension is *emission-insensitive*. WAIS dynamics accounts for 91% of total forecast variance and is largely decoupled from the emission pathway because ocean thermal forcing sufficient to sustain grounding-line retreat is already committed [50, 51]. The tail of the forecast distribution—the outcomes that drive the most severe human and economic consequences—is controlled almost entirely by whether WAIS retreat remains gradual or transitions to rapid, irreversible collapse. Emissions reduction compresses the median but barely touches this tail: the probability of exceeding 2 m relative to pre-industrial is 7% under the 2 °C trajectory and 19% under 5 °C. This factor-of-three difference is meaningful, but it does not eliminate the risk. Under any plausible emission pathway, there remains a non-negligible probability of catastrophic sea-level outcomes that emissions reduction alone cannot address.

The orthogonality of these two risk dimensions has a direct implication for resource allocation. Emissions reduction and ice-sheet stabilization are not competing priorities—they are complementary interventions operating on independent components of the problem. Emissions reduction without ice-sheet stability yields a well-constrained forecast centered on a potentially catastrophic outcome. Ice-sheet stability without emissions reduction yields a safe ice-sheet contribution but a wider envelope from thermally driven components. Adequate risk management requires both.

Implications for ice-sheet intervention

The concentration of forecast uncertainty in a single physical system—and specifically in a single binary question about the reversibility of WAIS retreat—points to the value of research and potential interventions aimed at ice-sheet stabilization. Proposed interventions, including submarine barriers to restrict warm water access to ice-sheet grounding zones [?], targeted cooling of critical ice shelves, and grounding-line stabilization through sediment

deposition, remain at an early conceptual stage. Our results provide quantitative motivation for advancing this research: if any intervention could shift probability mass from the unstable to the stable WAIS regime, the expected reduction in sea-level rise—and its associated human and economic costs—would be large. The stable-WAIS forecast (90% range of 540–680 mm) versus the full forecast (90% range of 600–2100 mm) represents a reduction in upper-bound risk of approximately 1.4 m, corresponding to hundreds of millions of fewer people exposed to coastal flooding [7] and trillions of dollars in avoided damages [9].

Furthermore, the forecast uncertainty is not smoothly distributed across outcomes but is concentrated in two distinct regimes. Conditional on a stable WAIS, the 90% range is approximately 140 mm. Conditional on ice-sheet instability, it expands to approximately 1500 mm with a median nearly 400 mm higher. The unconditional forecast—a probability-weighted mixture of these regimes—produces a wide confidence interval that reflects the superposition of two narrower distributions rather than a single diffuse uncertainty. A coastal community designing for the unconditional median prepares for an outcome that falls between the two most probable futures, potentially underbuilding for instability while overbuilding for stability. Adaptation strategies should therefore be designed in stages: sized for the stable regime now, with identified triggers and pre-planned expansions if observations confirm the onset of instability.

Reducing this structural uncertainty requires resolving a specific physical question: whether the observed acceleration in West Antarctic mass loss reflects a transient response to ocean forcing or the early stages of irreversible retreat. This is, by a wide margin, the single most consequential open question in sea-level science, and the one where targeted investment in observations, process understanding, and intervention research would yield the greatest return in forecast precision and risk reduction.

Broader implications

The data-driven framework presented here is not a replacement for process models, which remain essential for understanding the mechanisms of ice-sheet change. Our framework is complementary: an independent, data-driven estimate that makes its assumptions and limitations explicit, provides an observational benchmark against which process models can be tested, and produces conditional distributions that reflect what the data currently support. We consider robust, well-calibrated process models to be high-value targets for research and development, but the evidence suggests these models are not ready to be used for planning that costs billions of dollars and impacts millions of lives.

We emphasize that the disconnect between the high costs of sea-level rise and the large uncertainties in current sea-level projections is a consequence of chronic under-investment compounded by the lack of institutional homes able to focus on translating knowledge gained through basic scientific research into forecasts for coastal communities in the way that, for example, the National Weather Service has traditionally done [57]. A specific example is that modeling efforts for ISMIP6, which informed IPCC AR6 projections, were largely unfunded, leading to a dearth of model runs and near-absence of model parameter-sensitivity studies [18]. The world is going to endure high costs from sea-level rise (Fig. 3b). The questions are: how high will those costs be, and will we invest now to understand where and how to target funding to reduce the harm to people, ecosystems, and infrastructure, or

continue on our current path? Our analysis shows that this investment must be directed along two independent axes—reducing emissions to compress the predictable components, and understanding and potentially stabilizing the West Antarctic Ice Sheet to address the dominant, emission-insensitive uncertainty that defines the upper bound of 21st-century sea-level rise.

References

- [1] T. Frederikse, *et al.*, The causes of sea-level rise since 1900. *Nature* **584**, 393–397 (2020), doi:10.1038/s41586-020-2591-3.
- [2] B. D. Hamlington, *et al.*, The rate of global sea level rise doubled during the past three decades. *Communications Earth & Environment* **5**, 601 (2024), doi:10.1038/s43247-024-01761-5.
- [3] R. A. Rohde, Z. Hausfather, The Berkeley Earth land/ocean temperature record. *Earth System Science Data* **12**, 3469–3479 (2020), doi:10.5194/essd-12-3469-2020.
- [4] B. D. Beckley, P. S. Callahan, D. W. Hancock, G. T. Mitchum, R. D. Ray, On the “Cal-Mode” Correction to TOPEX Satellite Altimetry and Its Effect on the Global Mean Sea Level Time Series. *Journal of Geophysical Research: Oceans* **122** (11), 8371–8384 (2017), doi:10.1002/2017JC013090.
- [5] S. Rahmstorf, A semi-empirical approach to projecting future sea-level rise. *Science* **315**, 368–370 (2007), doi:10.1126/science.1135456.
- [6] M. Vermeer, S. Rahmstorf, Global sea level linked to global temperature. *Proceedings of the National Academy of Sciences* **106** (51), 21527–21532 (2009), doi:10.1073/pnas.0907765106.
- [7] S. A. Kulp, B. H. Strauss, New elevation data triple estimates of global vulnerability to sea-level rise and coastal flooding. *Nature Communications* **10**, 4844 (2019), doi:10.1038/s41467-019-12808-z.
- [8] B. Jongman, P. J. Ward, J. C. J. H. Aerts, Global exposure to river and coastal flooding: Long term trends and changes. *Global Environmental Change* **22** (4), 823–835 (2012), doi:10.1016/j.gloenvcha.2012.07.004.
- [9] S. Jevrejeva, L. P. Jackson, A. Grinsted, D. Lincke, B. Marzeion, Flood damage costs under the sea level rise with warming of 1.5 °C and 2 °C. *Environmental Research Letters* **13** (7), 074014 (2018), doi:10.1088/1748-9326/aacc76.
- [10] T. Tiggeloven, *et al.*, Global-scale benefit–cost analysis of coastal flood adaptation. *Natural Hazards and Earth System Sciences* **20**, 1025–1044 (2020), doi:10.5194/nhess-20-1025-2020.
- [11] United Nations Environment Programme, *Adaptation Gap Report 2023: Underfinanced. Underprepared.*, Tech. rep., UNEP (2023), doi:10.59117/20.500.11822/43796.

- [12] T. O'Donnell, Managed retreat and planned retreat: a systematic literature review. *Philosophical Transactions of the Royal Society B* **377**, 20210129 (2022), doi:10.1098/rstb.2021.0129.
- [13] F. Tubridy, M. Scott, M. Lennon, Managed retreat in response to flooding: lessons from the past for contemporary climate change adaptation. *Planning Perspectives* **36** (6), 1249–1268 (2021), doi:10.1080/02665433.2021.1939115.
- [14] A. R. Siders, I. Ajibade, D. Casagrande, Transformative potential of managed retreat as climate adaptation. *Current Opinion in Environmental Sustainability* **50**, 272–280 (2021), doi:10.1016/j.cosust.2021.06.007.
- [15] A. R. Siders, I. Ajibade, Introduction: Managed retreat and environmental justice in a changing climate. *Journal of Environmental Studies and Sciences* **11**, 287–293 (2021), doi:10.1007/s13412-021-00700-6.
- [16] B. Fox-Kemper, *et al.*, Ocean, cryosphere and sea level change, in *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge University Press) (2021), doi:10.1017/9781009157896.011.
- [17] D. Hirschfeld, *et al.*, Global survey shows planners use widely varying sea-level rise projections for coastal adaptation. *Communications Earth & Environment* **4**, 102 (2023), doi:10.1038/s43247-023-00703-x.
- [18] A. Aschwanden, T. C. Bartholomaus, D. J. Brinkerhoff, M. Truffer, Brief communication: A roadmap towards credible projections of ice sheet contribution to sea level. *The Cryosphere* **15**, 5705–5715 (2021), doi:10.5194/tc-15-5705-2021.
- [19] H. A. Fricker, *et al.*, Antarctica in 2025: Drivers of deep uncertainty in projected ice loss. *Science* **387** (6734), 601–609 (2025), doi:10.1126/science.adt9619.
- [20] B. D. Hamlington, *et al.*, Observation-based trajectory of future sea level for the coastal United States tracks near high-end model projections. *Communications Earth & Environment* **3**, 230 (2022), doi:10.1038/s43247-022-00537-z.
- [21] Z. Hausfather, G. P. Peters, Emissions – the ‘business as usual’ story is misleading. *Nature* **577**, 618–620 (2020), doi:10.1038/d41586-020-00177-3.
- [22] J. D. Millstein, B. M. Minchew, S. S. Pegler, Ice viscosity is more sensitive to stress than commonly assumed. *Communications Earth & Environment* **3**, 57 (2022), doi:10.1038/s43247-022-00385-x.
- [23] B. Getraer, M. Morlighem, Increasing the Glen-Nye power-law exponent accelerates ice-loss projections for the Amundsen Sea Embayment, West Antarctica. *Geophysical Research Letters* **52** (7), e2024GL112516 (2025), doi:10.1029/2024GL112516.
- [24] K. C. Green, J. S. Armstrong, Simple versus complex forecasting: The evidence. *Journal of Business Research* **68**, 1678–1685 (2015), doi:10.1016/j.jbusres.2015.03.026.

- [25] S. Makridakis, E. Spiliotis, V. Assimakopoulos, The M4 Competition: 100,000 time series and 61 forecasting methods. *International Journal of Forecasting* **36**, 54–74 (2020), doi:10.1016/j.ijforecast.2019.04.014.
- [26] T. Isaac, N. Petra, G. Stadler, O. Ghattas, Scalable and efficient algorithms for the propagation of uncertainty from data through inference to prediction for large-scale problems, with application to flow of the Antarctic ice sheet. *Journal of Computational Physics* **296**, 348–368 (2015), doi:10.1016/j.jcp.2015.04.047.
- [27] N. Petra, J. Martin, G. Stadler, O. Ghattas, A computational framework for infinite-dimensional Bayesian inverse problems, Part II: Stochastic Newton MCMC with application to ice sheet flow inverse problems. *SIAM Journal on Scientific Computing* **36** (4), A1525–A1555 (2014), doi:10.1137/130934805.
- [28] J. A. Church, *et al.*, Sea level change, in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the IPCC* (Cambridge University Press) (2013), doi:10.1017/CBO9781107415324.026.
- [29] R. E. Kopp, *et al.*, Probabilistic 21st and 22nd century sea-level projections at a global network of tide-gauge sites. *Earth’s Future* **2** (8), 383–406 (2014), doi:10.1002/2014EF000239.
- [30] M. Mengel, *et al.*, Future sea level rise constrained by observations and long-term commitment. *Proceedings of the National Academy of Sciences* **113** (10), 2597–2602 (2016), doi:10.1073/pnas.1500515113.
- [31] T. E. Wong, *et al.*, BRICK v0.2, a simple, accessible, and transparent model framework for climate and regional sea-level projections. *Geoscientific Model Development* **10**, 2741–2760 (2017), doi:10.5194/gmd-10-2741-2017.
- [32] X. Fettweis, *et al.*, GrSMBMIP: intercomparison of the modelled 1980–2012 surface mass balance over the Greenland Ice Sheet. *The Cryosphere* **14**, 3935–3958 (2020), doi:10.5194/tc-14-3935-2020.
- [33] B. Noël, L. van Kampenhout, J. T. M. Lenaerts, W. J. van de Berg, M. R. van den Broeke, A 21st century warming threshold for sustained Greenland ice sheet mass loss. *Geophysical Research Letters* **48**, e2020GL090471 (2021), doi:10.1029/2020GL090471.
- [34] J. Mouginot, *et al.*, Forty-six years of Greenland Ice Sheet mass balance from 1972 to 2018. *Proceedings of the National Academy of Sciences* **116** (19), 9239–9244 (2019), doi:10.1073/pnas.1904242116.
- [35] H. Seroussi, *et al.*, ISMIP6 Antarctica: a multi-model ensemble of the Antarctic ice sheet evolution over the 21st century. *The Cryosphere* **14**, 3033–3070 (2020), doi:10.5194/tc-14-3033-2020.
- [36] O. Geoffroy, *et al.*, Transient climate response in a two-layer energy-balance model. Part I: Analytical solution and parameter calibration using CMIP5 AOGCM experiments. *Journal of Climate* **26**, 1841–1857 (2013), doi:10.1175/JCLI-D-12-00195.1.

- [37] S. Levitus, *et al.*, World ocean heat content and thermosteric sea level change (0–2000 m), 1955–2010. *Geophysical Research Letters* **39**, L10603 (2012), doi:10.1029/2012GL051106.
- [38] S. G. Purkey, G. C. Johnson, Warming of global abyssal and deep Southern Ocean waters between the 1990s and 2000s: Contributions to global heat and sea level rise budgets. *Journal of Climate* **23**, 6336–6351 (2010), doi:10.1175/2010JCLI3682.1.
- [39] Y. Li, J. A. Church, T. J. McDougall, P. M. Barker, Sensitivity of observationally based estimates of ocean heat content and thermal expansion to vertical interpolation schemes. *Geophysical Research Letters* **49**, e2022GL101079 (2022), doi:10.1029/2022GL101079.
- [40] B. Noël, *et al.*, Modelling the climate and surface mass balance of polar ice sheets using RACMO2—Part 1: Greenland (1958–2016). *The Cryosphere* **12**, 811–831 (2018), doi:10.5194/tc-12-811-2018.
- [41] P. M. Alexander, *et al.*, Greenland Ice Sheet seasonal and spatial mass variability from model simulations and GRACE (2003–2012). *The Cryosphere* **10**, 1259–1277 (2016), doi:10.5194/tc-10-1259-2016.
- [42] B. Noël, W. J. van de Berg, S. Lhermitte, M. R. van den Broeke, Rapid ablation zone expansion amplifies North Greenland mass loss. *Science Advances* **5**, eaaw0123 (2019), doi:10.1126/sciadv.aaw0123.
- [43] Q. Glaude, *et al.*, A factor two difference in 21st-century Greenland Ice Sheet surface mass balance projections from three regional climate models under a strong warming scenario (SSP5-8.5). *Geophysical Research Letters* **51** (22), e2024GL111902 (2024), doi:10.1029/2024GL111902.
- [44] S. A. Good, M. J. Martin, N. A. Rayner, EN4: Quality controlled ocean temperature and salinity profiles and monthly objective analyses with uncertainty estimates. *Journal of Geophysical Research: Oceans* **118**, 6704–6716 (2013), doi:10.1002/2013JC009067.
- [45] K. D. Mankoff, *et al.*, Greenland ice sheet mass balance from 1840 through next week. *Earth System Science Data* **13**, 5001–5025 (2021), doi:10.5194/essd-13-5001-2021.
- [46] K. Poinar, *et al.*, Greenland Ice Sheet, in *Arctic Report Card 2025* (NOAA) (2026), doi:10.25923/rbsh-t897.
- [47] M. Zemp, *et al.*, Community estimate of global glacier mass changes from 2000 to 2023. *Nature* **639**, 382–388 (2025), glaMBIE consortium, doi:10.1038/s41586-024-08545-z.
- [48] D. R. Rounce, *et al.*, Global glacier change in the 21st century: Every increase in temperature matters. *Science* **379**, 78–83 (2023), doi:10.1126/science.abo1324.
- [49] I. Joughin, B. E. Smith, B. Medley, Marine ice sheet collapse potentially under way for the Thwaites Glacier Basin, West Antarctica. *Science* **344**, 735–738 (2014), doi:10.1126/science.1249055.

- [50] K. A. Naughten, P. R. Holland, J. De Rydt, Unavoidable future increase in West Antarctic ice-shelf melting over the twenty-first century. *Nature Climate Change* **13**, 1222–1228 (2023), doi:10.1038/s41558-023-01818-x.
- [51] S. H. R. Rosier, G. H. Gudmundsson, A. Jenkins, K. A. Naughten, Calibrated sea level contribution from the Amundsen Sea sector, West Antarctica, under RCP8.5 and Paris 2C scenarios. *The Cryosphere* **19**, 2527–2557 (2025), doi:10.5194/tc-19-2527-2025.
- [52] D. N. Goldberg, M. Morlighem, N. Gourmelen, Recent observations of Thwaites Glacier, West Antarctica are consistent with high rates of loss in next 50 years. *Geophysical Research Letters* **53** (5), e2025GL118823 (2026), doi:10.1029/2025GL118823.
- [53] E. Rignot, *et al.*, Thirty years of glacier grounding line retreat in Antarctica. *Proceedings of the National Academy of Sciences* **123** (10), e2524380123 (2026), doi:10.1073/pnas.2524380123.
- [54] A. A. Robel, H. Seroussi, G. H. Roe, Marine ice sheet instability amplifies and skews uncertainty in projections of future sea-level rise. *Proceedings of the National Academy of Sciences* **116** (30), 14887–14892 (2019), doi:10.1073/pnas.1904822116.
- [55] D. F. Martin, *et al.*, Rates of sea-level rise are highly sensitive to ice viscosity parameters in model benchmarks. *AGU Advances* **7** (2), e2025AV001946 (2026), doi:10.1029/2025AV001946.
- [56] J. L. Bamber, M. Oppenheimer, R. E. Kopp, W. P. Aspinall, R. M. Cooke, Ice sheet contributions to future sea-level rise from structured expert judgment. *Proceedings of the National Academy of Sciences* **116** (23), 11195–11200 (2019), doi:10.1073/pnas.1817205116.
- [57] G. Catania, T. Moon, A. Aschwanden, Glacial knowledge gaps impede resilience to sea level rise. *Eos* **103** (2022), doi:10.1029/2022EO220238.




Figure 1: **Observed global mean sea-level rise and its response to global mean surface temperature.** (a) GMSLR (blue, 5-year smooth; [1]) and NASA satellite altimetry (light blue; [4]), with the satellite-era extrapolation (dashed; [2]). GMST (red, 15-year smooth [3]) is referenced to the righthand axis. (b) Rate of GMSLR ($\dot{S} = dGMSLR/dt$) versus GMST anomaly, T . Observed values are shown in circles, colored by year, and the component model of this is colored by instantaneous transient sensitivity $\alpha = d\dot{S}/dT$. Other semi-empirical model results are shown as dashed [5] and dash-dot lines [6]. Horizontal and vertical blue lines mark the 1–2 and 2–4 mm yr⁻¹ rate doublings that occurred during 1976–1992 (1959–1993 with $\pm 1\sigma$) and 1992–2013 (1992–2014, $\pm 1\sigma$), respectively. The corresponding warming increments (ΔT) show that the first doubling required less warming (0.34 ± 0.04 °C) than the second (0.41 ± 0.03 °C). The vertical red band indicates the transition from thermosteric- (ocean thermal expansion) to barystatic-dominated sea-level rise (changes in glacier ice and water storage). The sinuous tail in the early observations is largely due to dam impoundment of water rather than a thermodynamic signal [1].




Figure 2: **Model calibration, validation, and WAIS uncertainty.** (a) Two-layer energy balance model fitted jointly to NOAA thermosteric sea level at 0–700 m (1957–2025) and 0–2000 m (2005–2025). (b) Greenland ice discharge modeled as a cumulative time-delay response to subsurface ocean warming ($\delta = 5$ yr, $R^2 = 0.995$). Calibration data (Mouginit et al. (2019) [34], filled circles) and withheld validation data (NOAA Arctic Report Card, diamonds) both fall within the 90% credible interval. (c) Global glacier contribution fitted to GlaMBIE consensus observations (2000–2023) using a linear temperature-driven model ($R^2 = 0.995$). Grey band shows the 90% credible interval. (d,e) Budget closure: temperature-forced component hindcast (stacked) compared with NASA altimetry and reconstructions. The component sum closes the 2000–2025 budget to within 6%. (f) WAIS: IMBIE satellite observations (black) compared with the ISMIP6 ensemble median and interquartile range (green; [35]), our full scenario mixture (blue), the stable-WAIS scenario (purple), and IPCC AR6 total Antarctic projection (red).




Figure 3: Distribution and exceedance probability of global mean sea-level projections. (a) Probability densities for the composite forecast of this study (blue), extrapolation of observations (grey; [2]), and IPCC AR6 projections (red; [16]). Vertical dashed lines indicate the respective medians. The purple band shows the 90% confidence interval for the case that WAIS does not experience a runaway acceleration, and sea level rise is due primarily to thermosteric, Greenland, and glaciers. We plot a purple band rather than distribution because the distribution is much tighter, and therefore taller, than the other distributions. (b) Probability of exceeding a given sea-level threshold for the 2°C (dash-dot), 3°C (solid), and 4°C (dashed) warming trajectories including instabilities in the West Antarctic Ice Sheet. Purple line shows the 3°C trajectory with a stable West Antarctic Ice Sheet. Lower axes show the corresponding number of additional people on land below flood level [7] and global annual flood costs [9], both assuming no additional coastal adaptation.