

Recitation 29: Lines and curves in space

Warm up:

Find a vector-valued function for the line segment connecting the points $P = (-3, 7, 6)$ and $Q = (5, -4, 7)$ in such a way that the value at $t = 0$ is P and the value at $t = 1$ is Q . Also, find the point two-thirds of the way from P to Q .

Group work:

Problem 1 Find a vector-valued function for the line through the point $(1, -2, 3)$ that is perpendicular to the lines

$$\vec{r}_1(t) = \langle 7, 8, -2 \rangle + t\langle 3, 5, 7 \rangle \quad \text{and} \quad \vec{r}_2(s) = \langle 4, -3, -7 \rangle + s\langle 4, 9, -1 \rangle$$

Problem 2 Find the distance from the point $P(-1, 4, 3)$ to the line $\langle 8 + t, 3 - 3t, -26t \rangle$. *Hint: The distance from the point to the line is the distance from the point P and the closest point on the line.*

Problem 3 Show that the curve $\vec{r} = \langle t \cos t, t \sin t, t \rangle$ lies completely on the cone $z^2 = x^2 + y^2$.

Problem 4 Match each of the following curves to the corresponding vector-valued function.

(a) $\langle 3, t^2, 5 \rangle$

(c) $\langle 3, \sin t, \cos t \rangle$

(e) $\langle \sin t, \cos t, 2 \cos t \rangle$

(b) $\langle 3, t^2, t \rangle$

(d) $\langle 3t, 5 \sin t, 5 \cos t \rangle$

(f) $\langle 2 \cos t, \sin t, \cos(3t) \rangle$

Recitation 29: Lines and curves in space




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6