

Particle Physics Phenomenology exercise 7

1. Study the space-time picture for a yo-yo which originally has

$$\begin{aligned} p_q &= k \frac{E_{\text{cm}}}{2} (1; 0, 0, 1) \\ p_{\bar{q}} &= \frac{1}{k} \frac{E_{\text{cm}}}{2} (1; 0, 0, -1) \end{aligned}$$

for $k \neq 1$, by applying $dp_z/dt = \pm\kappa$.

Show that it agrees with what is obtained by boosting an original system with $k = 1$.

2. Consider the string piece spanned between two points (t_1, z_1) and (t_2, z_2) along the string, where $z_1 > z_2$. Show that the relations

$$\begin{aligned} E &= \kappa(z_1 - z_2) \\ p_z &= \kappa(t_1 - t_2) \end{aligned}$$

hold for a string piece “at rest”, i.e. with $t_1 = t_2$. Thereafter boost along the z axis to show that it holds generally.

3. The charm mass varies between different approaches (e.g. “current algebra” vs. “constituent”), in the range 1.2 – 1.6 GeV. Estimate the range of tunneling suppression factors that would correspond to.
4. Assume that all string branchings produce particles with $m_\perp = 1$ GeV and that the fragmentation function is $f(z) = \delta(z - 1/2)$. Estimate the typical $\Gamma = (\kappa\tau)^2$ value, and from there the typical proper time at which strings break.

Hint: use the recursive formula

$$\Gamma_i = (1 - z_i) \left(\Gamma_{i-1} + \frac{m_{\perp i}^2}{z_i} \right)$$

with equilibrium corresponding to $\Gamma_i = \Gamma_{i-1}$.

5. Consider a back-to-back two-parton system, e.g. from e^+e^- annihilation. Show that the absolute momentum of a jet is approximately $|\mathbf{p}_{\text{jet}}| \approx E_{\text{jet}} - \text{constant}$, where $E_{\text{jet}} \approx E_{\text{cm}}/2$. How is the constant related to the typical Γ value introduced above? What does it imply about the jet mass?

Hint: consider how one string break in the middle of the event splits the system into two parts.

6. Compare the average multiplicity of a “Mercedes” three-jet event ($E_q = E_{\bar{q}} = E_g = E_{\text{cm}}/3$) with a two-jet event of the same energy ($E_q = E_{\bar{q}} = E_{\text{cm}}/2$). Assume that the multiplicity per string piece is proportional to its effective rapidity range in its rest frame, defined for an average transverse hadron mass $\langle m_\perp \rangle$.