

# North Atlantic surface water mass transformation contributions to AMOC in eddy-parameterized and eddy-resolving simulations

OS11D-0707

Monday, December 9

8:30-12:20

Ben Moore-Maley<sup>1</sup>, Julie McClean<sup>2</sup>, Sarah Gille<sup>2</sup>, Alice Barthel<sup>1</sup>, Luke Van Roeke<sup>1</sup>

<sup>1</sup>Los Alamos National Laboratory, Los Alamos, New Mexico

<sup>2</sup>Scripps Institution of Oceanography/University of California San Diego, La Jolla, California

Contact: bmoorema@lanl.gov

## Motivation

The Atlantic Meridional Overturning Circulation (AMOC) is an important component of the global climate system. Surface water mass transformation in the subpolar North Atlantic preconditions the deep water formation that contributes to the deep equatorward branch of AMOC. Surface transformation is thus a valuable diagnostic for interpreting observations of overturning circulation and evaluating model AMOC performance.




Figure 1 Map of the subpolar North Atlantic showing bathymetry, study regions and subpolar gyre circulation. Regions are bounded equatorward by the Overturning in the Subpolar North Atlantic Program (OSNAP) observing lines (Lozier et al., 2019).

## Research question

How does surface water mass transformation contribute to subpolar overturning circulation in an eddy-parameterized vs. eddy-resolving ocean simulation?

## Key points

1. Water mass analysis in temperature-salinity (TS) coordinates can be a powerful diagnostic for evaluating ocean model AMOC performance.
2. In the eddy-parameterized (LR) simulation presented here, an early surface freshwater bias suppresses surface transformation in the Irminger and Labrador Seas
3. This suppressed transformation is accompanied by poorly-formed TS properties of the subpolar gyre boundary currents.

## Method

Surface water mass transformation is based on the Walin (1982) framework.

### (1) Density transformation\*

$$Tr(\rho) = -\frac{1}{\Delta\rho} \iint_A \left[ \frac{\alpha H_{net}}{C_p} + \beta SF_{net} \right] \times \nabla [\rho, \rho'(x, y)] dA$$

Bin size      Heat flux component      Freshwater flux component      Binning function

### (2) Temperature transformation\*\*

$$Tr(T) = \frac{1}{\Delta T} \iint_A \frac{H_{net}}{\rho' C_p} \times \nabla [T, T'(x, y)] dA$$

### (3) Salinity transformation\*\*

$$Tr(S) = -\frac{1}{\Delta S} \iint_A S' F_{net} \times \nabla [S, S'(x, y)] dA$$

\*Speer and Tziperman (1992), \*\*Evans et al. (2014)

## Simulations

DOE Energy Exascale Earth System Model (E3SM) (Golaz et al., 2022)

- Forced configurations with coupled ocean and sea ice components
  - Model for Prediction Across Scales (MPAS-Ocean, MPAS-Seice)
- Unstructured hexagonal mesh, allows regional resolution refinement

## Configurations

- Low Resolution (LR), 30-60 km
  - Gent and McWilliams (1990) + Redi (1982) mesoscale eddy parameterizations
- High Resolution (HR), 6-18 km
  - mesoscale eddy resolving

## Forcing and initialization

- Coordinated Ocean-ice Reference Experiments II (CORE-II) protocol (1948-2009) (Large and Yeager, 2009)
- Polar Science Center Hydrographic Climatology (PHC2) initialization (Steele et al., 2001)

## AMOC

- AMOC weakens from the initial state in the LR simulation over the first 10 years




Figure 2 AMOC stream function time series at 26.5°N and time mean years 51-60.

## Early simulation adjustment

- LR: surface freshening, mixed layer shoaling, and westward subpolar gyre shift




Figure 3 Low Resolution (LR): Jan-Mar (JFM) surface salinity, JFM mixed layer depth, and annual 0-500 m velocity fields for years 1, 2, and 3-7. JFM sea ice extent (white) and 27.3 kg m⁻³ surface density (red) also shown.

- HR: no significant adjustment, except Greenland Sea mixed layer deepening




Figure 4 As in Figure 3, but for the High Resolution (HR) simulation.

## Surface density transformation

- At higher densities, surface transformation is increasingly present in the Irminger and Labrador Seas at HR, but not at LR




Figure 6 Surface density transformation maps (integrand of Eq. 1) at three potential density bins during years 3-7. Min/max JFM surface outcrops are also shown for each density bin.

## TS transformation and overturning

- In both simulations, transformation proceeds from warm/salty (Iceland Basin) to cold/fresh (Labrador Sea)
- Transformation is present between 27.6 and 27.8 kg m⁻³ at HR, but not at LR
- This transformation accompanies a robust Labrador Sea Water export at HR




Figure 7 Surface temperature-salinity (TS) transformation (arrows) and net import (blue) and export (red) transports by region (years 3-7). Potential density in gray.