

Blood Pressure Log

Name or label: _____ Dates: _____

Use your clinician's monitoring schedule

Rest seated for five minutes. Keep your back and arm supported, feet flat, and cuff on bare skin at heart level. Avoid exercise, nicotine and caffeine for 30 minutes beforehand. Take two readings one minute apart and write down **both numbers from both readings**. Do not keep only the lowest.

My schedule: _____ My BP goal: _____

Date / time	Reading 1 Top / bottom	Reading 2 Top / bottom	Pulse (optional)	Notes / symptoms
	___ / ___	___ / ___		
	___ / ___	___ / ___		
	___ / ___	___ / ___		
	___ / ___	___ / ___		
	___ / ___	___ / ___		
	___ / ___	___ / ___		
	___ / ___	___ / ___		
	___ / ___	___ / ___		
	___ / ___	___ / ___		
	___ / ___	___ / ___		

When to act

If systolic BP is above 180 **or** diastolic BP is above 120 and you have chest pain, breathlessness, new weakness, vision changes or trouble speaking, call 911. If you have no concerning symptoms, wait at least one minute and repeat; contact your healthcare professional immediately if it stays this high. Severe symptoms need urgent care even below these BP numbers.

Bring or send readings according to your care plan. Do not wait 12 weeks to report a concern.

Source: American Heart Association: Home Blood Pressure Monitoring