

## Discussion 4 – Thursday, September 2nd

1. Find the distance between the point  $P(2, 1, 3)$  and the line  $\mathbf{l}(t) = (2, 3, -2) + t(-1, 1, -2)$ .

**Solution:** See #4 from Discussion 3 (Tuesday August 31) worksheet.

2. Calculate the area of the parallelogram having vertices  $(1, -2)$ ,  $(4, -3)$ ,  $(3, -6)$  and  $(0, -5)$ .

**Solution:** The two (distinct) vector forms of edges are  $(4, -3) - (1, -2) = (3, -1)$  and  $(4, -3) - (3, -6) = (1, 3)$ , so the volume is given by

$$\begin{vmatrix} 3 & -1 \\ 1 & 3 \end{vmatrix} = (3)(3) - (-1)(1) = 10.$$

3. Consider the triangle where two of the sides are given by the vectors  $\mathbf{a} = \mathbf{i} - 3\mathbf{j}$  and  $\mathbf{b} = 4\mathbf{i} + 3\mathbf{j}$ . Compute the area of the triangle.

**Solution:** The area is half of the area of the parallelogram spanned by these vectors, so is

$$\frac{1}{2} \begin{vmatrix} 1 & -3 \\ 4 & 3 \end{vmatrix} = \frac{1}{2}((1)(3) - (-3)(4)) = \frac{15}{2}.$$

4.

(a) Find the volume of the parallelepiped formed by the vectors  $\mathbf{a} = (-2, 0, 5)$ ,  $\mathbf{b} = (4, -1, 0)$  and  $\mathbf{c} = (3, 1, 1)$ .

**Solution:** Again we take a determinant,

$$\begin{aligned} \begin{vmatrix} -2 & 0 & 5 \\ 4 & -1 & 0 \\ 3 & 1 & 1 \end{vmatrix} &= (-2) \begin{vmatrix} -1 & 0 \\ 1 & 1 \end{vmatrix} - (0) \begin{vmatrix} 4 & 0 \\ 3 & 1 \end{vmatrix} + (5) \begin{vmatrix} 4 & -1 \\ 3 & 1 \end{vmatrix} \\ &= (-2)(-1) + 0 + (5)(7) \\ &= 37. \end{aligned}$$

(b) Find the volume of the parallelepiped formed by the vectors  $\mathbf{a} = (-2, 0, 5)$ ,  $\mathbf{b} = (4, -1, 0)$  and  $\mathbf{d} = (8, -1, -10)$ . Interpret your answer geometrically. Specifically, what can you say about these three vectors?

**Solution:**

$$\begin{aligned} \begin{vmatrix} -2 & 0 & 5 \\ 4 & -1 & 0 \\ 8 & -1 & -10 \end{vmatrix} &= (-2) \begin{vmatrix} -1 & 0 \\ -1 & -10 \end{vmatrix} - (0) \begin{vmatrix} 4 & 0 \\ 8 & -10 \end{vmatrix} + (5) \begin{vmatrix} 4 & -1 \\ 8 & -1 \end{vmatrix} \\ &= (-2)(10) + 0 + (5)(4) \\ &= 0. \end{aligned}$$

If the volume of the parallelepiped is zero, it must be “flat”; indeed, the three vectors given here all lie in the same plane  $\mathbf{p}(t) = t\mathbf{a} + s\mathbf{b}$ , for all  $s, t \in \mathbb{R}$ . Notice that  $\mathbf{d} = -2\mathbf{a} + \mathbf{b}$ .

5.

(a) Calculate  $\mathbf{a} \times \mathbf{b}$  if  $\mathbf{a} = (3, -2, 3)$  and  $\mathbf{b} = (2, -4, 1)$ . Calculate  $\mathbf{b} \times \mathbf{a}$ . How do these two

cross products compare?

**Solution:**

$$\begin{aligned} \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & -2 & 3 \\ 2 & -4 & 1 \end{vmatrix} &= \begin{vmatrix} -2 & 3 \\ -4 & 1 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 3 & 3 \\ 2 & 1 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 3 & -2 \\ 2 & -4 \end{vmatrix} \mathbf{k} \\ &= 10\mathbf{i} + 3\mathbf{j} - 8\mathbf{k}. \\ \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -4 & 1 \\ 3 & -2 & 3 \end{vmatrix} &= \begin{vmatrix} -4 & 1 \\ -2 & 3 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 2 & 1 \\ 3 & 3 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 2 & -4 \\ 3 & -2 \end{vmatrix} \mathbf{k} \\ &= -10\mathbf{i} - 3\mathbf{j} + 8\mathbf{k}. \end{aligned}$$

Notice that these are *not* equal!  $\mathbf{a} \times \mathbf{b} = -\mathbf{b} \times \mathbf{a}$ .

**The Right Hand Rule:** If the fingers of your right hand curl in the direction of a rotation from  $\mathbf{a}$  to  $\mathbf{b}$ , then your thumb points in the direction of  $\mathbf{a} \times \mathbf{b}$ .

(b) Without doing any calculation, use the right hand rule to find a vector parallel to  $\mathbf{i} \times \mathbf{j}$ .

*Answer:*  $\mathbf{k}$

(c) Find a vector parallel to  $\mathbf{j} \times \mathbf{k}$ .

*Answer:*  $\mathbf{i}$

(d) Find a vector parallel to  $-\mathbf{k} \times \mathbf{i}$

*Answer:*  $-\mathbf{j}$

(e) Find a vector parallel to  $(\mathbf{i} + \mathbf{k}) \times (\mathbf{k} - \mathbf{i})$ .

*Answer:*  $-\mathbf{j}$

(f) Calculate  $(\mathbf{i} + \mathbf{k}) \times (\mathbf{k} - \mathbf{i})$ . Are your answers to (e) and (f) the same? What is different?

**Solution:**


$$\begin{aligned} \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 0 & 1 \\ -1 & 0 & 1 \end{vmatrix} &= \begin{vmatrix} 0 & 1 \\ 0 & 1 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 1 & 1 \\ -1 & 1 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 1 & 0 \\ -1 & 0 \end{vmatrix} \mathbf{k} \\ &= -2\mathbf{j}. \end{aligned}$$

The vector from (e) has the same direction as this vector, but has different magnitude.

## 6. The Triangle Inequality.

Let  $\mathbf{a}$  and  $\mathbf{b}$  be any vectors in  $\mathbb{R}^n$ . The triangle inequality states that  $\|\mathbf{a} + \mathbf{b}\| \leq \|\mathbf{a}\| + \|\mathbf{b}\|$ .

(a) Give a geometric interpretation of the triangle inequality. (i.e. Draw a picture in  $\mathbb{R}^2$  or  $\mathbb{R}^3$  that represents this inequality.)



**Solution:**

The triangle inequality says that the sum of the lengths of two sides of a triangle is at least as large as the length of the third side.

(b) Use the Cauchy-Schwarz inequality to prove the triangle inequality. (Hint:  $\|\mathbf{a} + \mathbf{b}\|^2 = (\mathbf{a} + \mathbf{b}) \cdot (\mathbf{a} + \mathbf{b})$  )

$$\begin{aligned}\|\mathbf{a} + \mathbf{b}\|^2 &= (\mathbf{a} + \mathbf{b}) \cdot (\mathbf{a} + \mathbf{b}) \\ &= \mathbf{a} \cdot \mathbf{a} + 2\mathbf{a} \cdot \mathbf{b} + \mathbf{b} \cdot \mathbf{b} \\ &= \|\mathbf{a}\|^2 + 2\mathbf{a} \cdot \mathbf{b} + \|\mathbf{b}\|^2 \\ &\leq \|\mathbf{a}\|^2 + 2|\mathbf{a} \cdot \mathbf{b}| + \|\mathbf{b}\|^2 \\ &\leq \|\mathbf{a}\|^2 + 2\|\mathbf{a}\| \|\mathbf{b}\| + \|\mathbf{b}\|^2 \\ &= (\|\mathbf{a}\| + \|\mathbf{b}\|)^2,\end{aligned}$$

and now taking square roots on each side gives us the triangle inequality.