**Tuesday, October 2** \*\* Taylor series, the 2<sup>nd</sup> derivative test, and changing coordinates.

- 1. Consider  $f(x, y) = 2\cos x y^2 + e^{xy}$ .
  - (a) Show that (0,0) is a critical point for f.
  - (b) Calculate each of  $f_{xx}$ ,  $f_{xy}$ ,  $f_{yy}$  at (0,0).
  - (c) For x, y fixed consider the function h(t) = f(tx, ty). Recall from Calc II that the  $2^{\text{nd}}$ -order Taylor approximation for h(t) at 0 is  $h(t) \approx h(0) + \frac{h'(0)}{1}t + \frac{h''(0)}{2}t^2$ . Use this (with t = 1) and the Chain Rule to write out the  $2^{\text{nd}}$ -order Taylor approximation g(x, y) for f at (0, 0), which should be a polynomial of degree two in x and y.
  - (d) To make sure the next two problems go smoothly, check your answer to (c) with the instructor.
- 2. Let g(x, y) be the approximation you obtained for f(x, y) near (0,0) in 1(c). It's not clear from the formula whether g, and hence f, has a min, max, or a saddle at (0,0). Test along several lines until you are convinced you've determined which type it is. In the next problem, you'll confirm your answer in two ways.
- 3. Consider alternate coordinates (u, v) on  $\mathbb{R}^2$  given by (x, y) = (u v, u + v).
  - (a) Sketch the u- and v-axes relative to the usual x- and y-axes, and draw the points whose (u, v)-coordinates are: (-1, 2), (1, 1), (1, -1).
  - (b) Express g as a function of u and v, and expand and simplify the resulting expression.
  - (c) Explain why your answer in 3(b) confirms your answer in 2.
  - (d) Sketch a few level sets for g. What do the level sets of f look like near (0,0)?
  - (e) It turns out that there is always a similar change of coordinates so that the Taylor series of a function f which has a critical point at (0,0) looks like  $f(u,v) \approx f(0,0) + au^2 + bv^2$ . In fact this is why the  $2^{\text{nd}}$  derivative test works.

Double check your answer in 2 by applying the  $2^{\mathrm{nd}}$ -derivative test directly to f.

- 4. Consider the function  $f(x, y) = 3xe^y x^3 e^{3y}$ .
  - (a) Check that f has only one critical point, which is a local maximum.
  - (b) Does f have an absolute maxima? Why or why not? Check your answer with the instructor.