557_Project_2BS


Ben Straub, Hillary Koch, Jiawei Huang, Arif Masrur 3/15/2017

No Command Lines Ever. Whoa


function (..., recursive = FALSE) .Primitive("c")

What the Factor Variables look like


Class/Response Distribution




Seismic Distribution



Seismoacoustic Distribution




Ghazard Distribution




What the Continuous Variables look like

density.default(x = seismic\$genergy)




density.default(x = seismic\$gpuls)




N = 2584 Bandwidth = 66.84

density.default(x = seismic\$gdenergy

density.default(x = seismic\$gdpuls)







N = 2584 Bandwidth = 10.47

N = 2584 Bandwidth = 9.244

density.default(x = seismic\$maxenerg\$nsity.default(x = seismic\$nbumps, adjus

Density







N = 2584 Bandwidth = 279.1

N = 2584 Bandwidth = 1.395

nsity.default(x = seismic\$nbumps2, adjusnsity.default(x = seismic\$nbumps3, adjus





N = 2584 Bandwidth = 1.395

N = 2584 Bandwidth = 1.395

Call:

```
lm(formula = class ~ ., data = seismic)
```

Residuals:


Coefficients:


Estimate Std. Error t value Pr(>|t|) (Intercept) -2.393e-02 2.565e-02 -0.933 0.35090 seismic 1.869e-02 1.076e-02 1.737 0.08254 . seismoacoustic 2.610e-03 1.002e-02 0.260 0.79457 6.190e-04 1.157e-02 0.054 0.95732 shift genergy -8.698e-08 3.459e-08 -2.514 0.01199 * 1.019e-04 1.670e-05 6.102 1.2e-09 *** gpuls -6.943e-05 1.006e-04 -0.690 0.49009 gdenergy gdpuls -1.942e-04 1.368e-04 -1.420 0.15583 -1.394e-02 1.608e-02 -0.867 0.38618 ghazard nbumps 4.674e-01 1.680e-01 2.783 0.00543 ** nbumps2 -4.282e-01 1.682e-01 -2.546 0.01096 * -4.260e-01 1.681e-01 -2.535 0.01131 * nbumps3 nbumps4 -4.622e-01 1.708e-01 -2.706 0.00685 ** nbumps5 -2.963e-01 2.332e-01 -1.270 0.20408 2.536e-07 2.395e-06 0.106 0.91568 energy -1.054e-06 2.333e-06 -0.452 0.65164 maxenergy

Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1


Residual standard error: 0.2371 on 2568 degrees of freedom Multiple R-squared: 0.09128, Adjusted R-squared: 0.08597 F-statistic: 17.2 on 15 and 2568 DF, p-value: < 2.2e-16

nsity.default(x = seismic\$nbumps4, adjusnsity.default(x = seismic\$nbumps5, adjus





Histogram of res




Lots of multicollinearity to worry about during variable selection

vif(fit)

##	seismic	seismoacoustic	shift	genergy	gpuls
##	1.209814	1.286183	1.411216	2.889651	4.057018
##	gdenergy	gdpuls	ghazard	nbumps	nbumps2
##	3.000282	3.430524	1.395598	2414.689538	798.964152
##	nbumps3	nbumps4	nbumps5	energy	maxenergy
##	769.131960	104.402690	11.562237	110.283444	93.762895

Correlation of the Variables




\$r

	genergy	gpuls	nbumps4	${\tt nbumps3}$	${\tt nbumps}$	${\tt nbumps2}$	${\tt nbumps5}$	gdenergy
genergy	1							
gpuls	0.75	1						
nbumps4	0.15	0.26	1					
nbumps3	0.19	0.23	0.18	1				
nbumps	0.22	0.3	0.4	0.8	1			
nbumps2	0.14	0.21	0.16	0.35	0.8	1		
nbumps5	-0.0099	0.049	-0.017	0.046	0.07	-0.0053	1	
gdenergy	0.049	0.29	0.037	-0.012	0.03	0.041	0.12	1

```
0.072 0.38
                          0.066
                                  0.015 0.058
                                                  0.051
                                                           0.14
                                                                    0.81
gdpuls
         gdpuls
genergy
gpuls
nbumps4
nbumps3
nbumps
nbumps2
nbumps5
gdenergy
gdpuls
              1
$p
         genergy gpuls nbumps4 nbumps3 nbumps nbumps2 nbumps5 gdenergy
genergy
               0
gpuls
                     0
nbumps4 1.4e-14
                     0
                              0
               0
                     0
                              0
                                      0
nbumps3
                     0
                              0
                                      0
                                             0
nbumps
               0
                                                      0
nbumps2 2.2e-13
                     0
                              0
                                      0
                                             0
            0.62 0.012
                                  0.018 4e-04
                                                   0.79
nbumps5
                            0.4
                                                              0
                                                  0.036 3.3e-10
gdenergy
           0.014
                          0.061
                                   0.54
                                          0.13
                                                                       0
         0.00027
                     0 0.00076
                                   0.45 0.0032 0.0094 5.9e-13
                                                                       0
gdpuls
         gdpuls
genergy
gpuls
nbumps4
nbumps3
nbumps
nbumps2
nbumps5
gdenergy
              0
gdpuls
$sym
         genergy gpuls nbumps4 nbumps3 nbumps nbumps2 nbumps5 gdenergy
genergy
gpuls
nbumps4
                        1
nbumps3
                                1
nbumps
                                        1
nbumps2
                                               1
nbumps5
                                                        1
gdenergy
                                                                1
gdpuls
         gdpuls
genergy
gpuls
nbumps4
nbumps3
nbumps
nbumps2
nbumps5
gdenergy
```

```
gdpuls 1
attr(,"legend")
[1] 0 ' ' 0.3 '.' 0.6 ',' 0.8 '+' 0.9 '*' 0.95 'B' 1
```



```
$r
        row
              column
                         cor
1
               gpuls
                      0.7500 0.0e+00
    genergy
2
    genergy
             nbumps4
                      0.1500 1.4e-14
                      0.2600 0.0e+00
3
             nbumps4
      gpuls
4
                      0.1900 0.0e+00
    genergy
             nbumps3
5
             nbumps3
                      0.2300 0.0e+00
     gpuls
6
    nbumps4
             nbumps3
                      0.1800 0.0e+00
7
              nbumps
                      0.2200 0.0e+00
    genergy
8
      gpuls
              nbumps
                      0.3000 0.0e+00
9
    nbumps4
              nbumps
                     0.4000 0.0e+00
10
    nbumps3
              nbumps
                      0.8000 0.0e+00
             nbumps2
                      0.1400 2.2e-13
11
    genergy
12
      gpuls
             nbumps2
                      0.2100 0.0e+00
13
             nbumps2 0.1600 0.0e+00
    nbumps4
14
    nbumps3
             nbumps2
                      0.3500 0.0e+00
             nbumps2 0.8000 0.0e+00
15
     nbumps
16
             nbumps5 -0.0099 6.2e-01
    genergy
17
      gpuls
             nbumps5 0.0490 1.2e-02
18
             nbumps5 -0.0170 4.0e-01
    nbumps4
19
    nbumps3
             nbumps5 0.0460 1.8e-02
20
     nbumps
             nbumps5 0.0700 4.0e-04
    nbumps2
             nbumps5 -0.0053 7.9e-01
```

```
22 genergy gdenergy 0.0490 1.4e-02
23
     gpuls gdenergy 0.2900 0.0e+00
24 nbumps4 gdenergy 0.0370 6.1e-02
25 nbumps3 gdenergy -0.0120 5.4e-01
   nbumps gdenergy 0.0300 1.3e-01
27 nbumps2 gdenergy 0.0410 3.6e-02
28 nbumps5 gdenergy 0.1200 3.3e-10
29 genergy gdpuls 0.0720 2.7e-04
30 gpuls gdpuls 0.3800 0.0e+00
31 nbumps4 gdpuls 0.0660 7.6e-04
32 nbumps3 gdpuls 0.0150 4.5e-01
   nbumps gdpuls 0.0580 3.2e-03
33
34 nbumps2 gdpuls 0.0510 9.4e-03
35 nbumps5 gdpuls 0.1400 5.9e-13
36 gdenergy gdpuls 0.8100 0.0e+00
$р
NULL
$sym
NULL
```


[1] 646 16

Separating into Test and Training Sets


```
## Setting up Test and Training Sets
##-----
n <- dim(seismic)[1]</pre>
p <- dim(seismic)[2]</pre>
set.seed(2016)
test <- sample(n, round(n/4))</pre>
train <- (1:n)[-test]
seismic.train <- seismic[train,]</pre>
seismic.test <- seismic[test,]</pre>
dim(seismic)
[1] 2584
           16
dim(seismic.train)
[1] 1938
           16
dim(seismic.test)
```


Linear regression of an indicator matrix

pc.comp



PC1 vs PC3






lda.class 0 1 0 1771 108 1 36 23

[1] 0.1755725


[1] 0.9800775



lda.class 0 1 0 591 34 1 16 5

[1] 0.1282051

[1] 0.9736409



Logistic Regression on the Training and Test Sets

	Estimate	Std. Error	z value	Pr(> z)	
(Intercept)	-6.343e+00	7.721e-01	-8.215	< 2e-16	***
seismic	4.808e-01	2.111e-01	2.278	0.022727	*
${\tt seismoacoustic}$	2.159e-01	1.993e-01	1.084	0.278524	
shift	1.179e+00	3.573e-01	3.301	0.000965	***
genergy	-2.471e-07	5.044e-07	-0.490	0.624239	
gpuls	7.095e-04	2.474e-04	2.868	0.004136	**
gdenergy	-1.904e-04	2.177e-03	-0.087	0.930292	
gdpuls	-2.997e-03	3.093e-03	-0.969	0.332500	
ghazard	-2.335e-01	3.509e-01	-0.666	0.505671	
nbumps	1.807e+01	5.354e+02	0.034	0.973080	
nbumps2	-1.773e+01	5.354e+02	-0.033	0.973590	
nbumps3	-1.771e+01	5.354e+02	-0.033	0.973611	
nbumps4	-1.806e+01	5.354e+02	-0.034	0.973097	

```
nbumps5 -1.604e+01 5.354e+02 -0.030 0.976095
energy 1.622e-06 4.033e-05 0.040 0.967929
maxenergy -7.101e-06 3.969e-05 -0.179 0.858012
---
Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
```

(Dispersion parameter for binomial family taken to be 1)

Null deviance: 958.82 on 1937 degrees of freedom Residual deviance: 813.40 on 1922 degrees of freedom AIC: 845.4


Number of Fisher Scoring iterations: 12

The predictors that are significant in our logistic model are seismic, shift and gpuls. The predictors nbumps6, nbumps7 and nbumps89 were removed as they did not provide any data.

[1] 0.9329205

[1] 0.04580153

[1] 0.997233



The diagonal elements of the confusion matrix indicate correct predictions, while the off-diagonals represent incorrect predictions. Hence our model on the training data set correctly predicted that the seismic activity would be of no harzard on 1786 observations and that it would be of hazard on 0 observations, for a total of 1786 + 0 = 1786 correct predictions. The mean() function can be used to compute the fraction of hazards for which the prediction was correct. In this case, logistic regression correctly predicted the class of hazard 92 percent of the time. The bad part about this 92 percent of the time is that it did not get any of our actual real hazards observations correct!!!

```
## [1] 0.9349845

## ## glm.pred 0 1

## 0.602 37

## 1 5 2

## [1] 0.05128205

## [1] 0.9917628
```

The diagonal elements of the confusion matrix indicate correct predictions, while the off-diagonals represent incorrect predictions. Hence our model on the testing data set correctly predicted that the seismic activity would be of no harzard on 605 observations and that it would be hazardous on 2 observations, for a total of 602 + 2 = 604 correct predictions. The mean() function can be used to compute the fraction of seismic activity for which the prediction was correct. In this case, logistic regression correctly predicted class of hazard 93.5 % of the time. However, again worrisome, is that the model miss 5 observations that were hazardous instances and 37 that were not hazardous.

Recall that the logistic regression model had only 3 predictors that were significant from an avaiable 19. Perhaps by removing the variables that appear not to be helpful in predicting seismic hazard, we can obtain a more effective model. After all, using predictors that have no relationship with the response tends to cause a deterioration in the test error rate (since such predictors cause an increase in variance without a corresponding decrease in bias), and so removing such predictors may in turn yield an improvement [straight from the book]