

# 557\_Project\_2BS


*Ben Straub, Hillary Koch, Jiawei Huang, Arif Masrur*

*3/15/2017*


## No Command Lines Ever. Whoa

What the Factor Variables look like


**Class/Response Distribution**




**Seismic Distribution**



**Seismoacoustic Distribution**




**Ghazard Distribution**




What the Continuous Variables look like


**density.default(x = seismic\$genergy)**




**density.default(x = seismic\$gpuls)**





**density.default(x = seismic\$gdenergy)**





**density.default(x = seismic\$gdpuls)**



**density.default(x = seismic\$maxenergy, adjus**



**nsity.default(x = seismic\$nbumps2, adjus**



Call:

```
lm(formula = class ~ ., data = seismic)
```

Residuals:

	Min	1Q	Median	3Q	Max
	-0.57549	-0.07778	-0.03812	-0.00950	1.03232

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	-2.393e-02	2.565e-02	-0.933	0.35090
seismic	1.869e-02	1.076e-02	1.737	0.08254 .
seismoacoustic	2.610e-03	1.002e-02	0.260	0.79457
shift	6.190e-04	1.157e-02	0.054	0.95732
genergy	-8.698e-08	3.459e-08	-2.514	0.01199 *
gpuls	1.019e-04	1.670e-05	6.102	1.2e-09 ***
gdenergy	-6.943e-05	1.006e-04	-0.690	0.49009
gdpuls	-1.942e-04	1.368e-04	-1.420	0.15583
ghazard	-1.394e-02	1.608e-02	-0.867	0.38618
nbumps	4.674e-01	1.680e-01	2.783	0.00543 **
nbumps2	-4.282e-01	1.682e-01	-2.546	0.01096 *
nbumps3	-4.260e-01	1.681e-01	-2.535	0.01131 *
nbumps4	-4.622e-01	1.708e-01	-2.706	0.00685 **
nbumps5	-2.963e-01	2.332e-01	-1.270	0.20408
energy	2.536e-07	2.395e-06	0.106	0.91568
maxenergy	-1.054e-06	2.333e-06	-0.452	0.65164

---









Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1








Residual standard error: 0.2371 on 2568 degrees of freedom

Multiple R-squared: 0.09128, Adjusted R-squared: 0.08597

F-statistic: 17.2 on 15 and 2568 DF, p-value: < 2.2e-16

nsity.default(x = seismic\$nbumps4, adjusnsity.default(x = seismic\$nbumps5, adjus






Lots of multicollinearity to worry about during variable selection

```
vif(fit)
```

```
##      seismic seismoacoustic      shift      genergy      gpuls
##      1.209814      1.286183      1.411216      2.889651      4.057018
##      gdenergy      gdpuls      ghazard      nbumps      nbumps2
##      3.000282      3.430524      1.395598      2414.689538      798.964152
##      nbumps3      nbumps4      nbumps5      energy      maxenergy
##      769.131960      104.402690      11.562237      110.283444      93.762895
```

## Correlation of the Variables



```
$r
      genergy gpuls nbumps4 nbumps3 nbumps nbumps2 nbumps5 gdenergy
genergy      1
gpuls      0.75      1
nbumps4      0.15 0.26      1
nbumps3      0.19 0.23 0.18      1
nbumps      0.22 0.3 0.4 0.8      1
nbumps2      0.14 0.21 0.16 0.35 0.8      1
nbumps5     -0.0099 0.049 -0.017 0.046 0.07 -0.0053      1
gdenergy      0.049 0.29 0.037 -0.012 0.03 0.041 0.12      1
```

gdpuls	0.072	0.38	0.066	0.015	0.058	0.051	0.14	0.81
gdpuls								

genergy  
gpuls  
nbumps4  
nbumps3  
nbumps  
nbumps2  
nbumps5  
gdenenergy  
gdpuls

1

\$p

	genergy	gpuls	nbumps4	nbumps3	nbumps	nbumps2	nbumps5	gdenenergy
genergy	0							
gpuls	0	0						
nbumps4	1.4e-14	0	0					
nbumps3	0	0	0	0				
nbumps	0	0	0	0	0			
nbumps2	2.2e-13	0	0	0	0	0		
nbumps5	0.62	0.012	0.4	0.018	4e-04	0.79	0	
gdenenergy	0.014	0	0.061	0.54	0.13	0.036	3.3e-10	0
gdpuls	0.00027	0	0.00076	0.45	0.0032	0.0094	5.9e-13	0
gdpuls								

genergy  
gpuls  
nbumps4  
nbumps3  
nbumps  
nbumps2  
nbumps5  
gdenenergy  
gdpuls

0

\$sym

	genergy	gpuls	nbumps4	nbumps3	nbumps	nbumps2	nbumps5	gdenenergy
genergy	1							
gpuls	,	1						
nbumps4			1					
nbumps3				1				
nbumps		.	,	1				
nbumps2			.	,	1			
nbumps5						1		
gdenenergy							1	
gdpuls		.					+	
gdpuls								

genergy  
gpuls  
nbumps4  
nbumps3  
nbumps  
nbumps2  
nbumps5  
gdenenergy

```
gdpuls 1
attr("legend")
[1] 0 ' ' 0.3 '.' 0.6 ',' 0.8 '+' 0.9 '*' 0.95 'B' 1
```

## Separating into Test and Training Sets

```
##-----
## Setting up Test and Training Sets
##-----

n <- dim(seismic)[1]
p <- dim(seismic)[2]

set.seed(2016)
test <- sample(n, round(n/4))
train <- (1:n)[-test]
seismic.train <- seismic[train,]
seismic.test <- seismic[test,]

dim(seismic)
```

```
[1] 2584 16
```

```
dim(seismic.train)
```

```
[1] 1938 16
```

```
dim(seismic.test)
```

```
[1] 646 16
```

```
#View(seismic.train)
#View(seismic.test)
```

## Linear regression of an indicator matrix

```
##-----
## Linear regression of indicator matrix
##-----

responseY <- seismic$class
predictorX <- seismic[,16]

# Following Le Bao's code
class1 <- which(responseY==1)
class0 <- which(responseY==0)
```



```

Y <- matrix(data = rep(0,length(responseY)*2),nrow = length(responseY))
Y[class0,1] <- 1
Y[class1,2] <- 1

betaHat <- solve(t(as.matrix(predictorX))%*%as.matrix(predictorX))%*%t(as.matrix(predictorX))%*%Y
Y1 <- as.matrix(predictorX)%*%betaHat[,1]
Y2 <- as.matrix(predictorX)%*%betaHat[,2]

pred.mx <- cbind(Y1,Y2)
pred <- rep(NA,length(Y1))
for(i in 1:length(Y1)){
  pred[i] <- which.max(pred.mx[i,]) - 1
}

# Confusion matrix
mx <- cbind(pred,responseY,pred-responseY)

confusion <- matrix(rep(NA,4), nrow = 2)
correct <- which(mx[,3] == 0)
confusion[1,1] <- length(which(mx[correct,1] == 0))
confusion[2,2] <- length(which(mx[correct,1] == 1))
confusion[1,2] <- length(which(mx[,3] == -1))
confusion[2,1] <- length(which(mx[,3] == 1))
confusion

##      [,1] [,2]
## [1,] 2411 169
## [2,]    3    1

sensitivity <- confusion[2,2]/sum(confusion[,2])
specificity <- confusion[1,1]/sum(confusion[,1])
error.rate <- (confusion[1,2] + confusion[2,1])/sum(confusion)
c(sensitivity, specificity, error.rate)

```

```
## [1] 0.005882353 0.998757249 0.066563467
```

## Linear Discriminant Analysis on full model

## Quadratic Discriminant Analysis -INCOMPLETE

```

##-----
## Fit QDA model
##-----

## Currently, can't perform QDA. This is probably due to multicollinearity in the model
## (can't invert covariance matrix) but should be possible after variable selection

#qda.fit <- qda(class~., data = seismic, subset = train)

```

## Regularized Discriminant Analysis -INCOMPLETE

```
##-----  
## Fit RDA model  
##-----  
  
## Currently, can't perform RDA. This is probably due to multicollinearity in the model  
## (can't invert covariance matrix) but should be possible after variable selection
```

## Logistic Regression

```
Call:  
glm(formula = class ~ ., family = binomial, data = seismic.train)
```

Deviance Residuals:

	Min	1Q	Median	3Q	Max
	-1.8471	-0.3860	-0.2851	-0.1566	3.0825

Coefficients:

	Estimate	Std. Error	z value	Pr(> z )
(Intercept)	-6.343e+00	7.721e-01	-8.215	< 2e-16 ***
seismic	4.808e-01	2.111e-01	2.278	0.022727 *
seismoacoustic	2.159e-01	1.993e-01	1.084	0.278524
shift	1.179e+00	3.573e-01	3.301	0.000965 ***
genergy	-2.471e-07	5.044e-07	-0.490	0.624239
gpuls	7.095e-04	2.474e-04	2.868	0.004136 **
gdenergy	-1.904e-04	2.177e-03	-0.087	0.930292
gdpuls	-2.997e-03	3.093e-03	-0.969	0.332500
ghazard	-2.335e-01	3.509e-01	-0.666	0.505671
nbumps	1.807e+01	5.354e+02	0.034	0.973080
nbumps2	-1.773e+01	5.354e+02	-0.033	0.973590
nbumps3	-1.771e+01	5.354e+02	-0.033	0.973611
nbumps4	-1.806e+01	5.354e+02	-0.034	0.973097
nbumps5	-1.604e+01	5.354e+02	-0.030	0.976095
energy	1.622e-06	4.033e-05	0.040	0.967929
maxenergy	-7.101e-06	3.969e-05	-0.179	0.858012

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for binomial family taken to be 1)

Null deviance: 958.82 on 1937 degrees of freedom  
Residual deviance: 813.40 on 1922 degrees of freedom  
AIC: 845.4

Number of Fisher Scoring iterations: 12

[1] 0.9329205

```
glm.pred  0  1
          0 1802 125
          1   5   6
```

```
[1] 0.04580153
```


```
[1] 0.997233
```

```
[1] 0.9349845
```


```
glm.pred  0  1
          0 604 39
          1   3  0
```

```
[1] 0
```

```
[1] 0.9950577
```



## Variable Selection-LASSO



[1] 8.670049

(Intercept)	seismic	seismoacoustic	shift	genergy
-8.144581e-03	8.800484e-03	0.000000e+00	7.977504e-03	0.000000e+00

gpuls	gdenergy	gdpuls	ghazard	nbumps
4.677101e-05	0.000000e+00	0.000000e+00	0.000000e+00	3.117955e-02
nbumps2	nbumps3	nbumps4	nbumps5	energy
0.000000e+00	0.000000e+00	0.000000e+00	0.000000e+00	0.000000e+00
maxenergy				
0.000000e+00				



## Variables selected through LASSO

(Intercept)	seismic	shift	gpuls	nbumps
-8.144581e-03	8.800484e-03	7.977504e-03	4.677101e-05	3.117955e-02


## Principal Component Analysis from the Book - INCOMPLETE



Importance of components:

	PC1	PC2	PC3	PC4	PC5	PC6
Standard deviation	1.9629	1.5284	1.5089	1.17618	1.05902	1.02457
Proportion of Variance	0.2408	0.1460	0.1423	0.08646	0.07009	0.06561
Cumulative Proportion	0.2408	0.3868	0.5291	0.61559	0.68568	0.75129
	PC7	PC8	PC9	PC10	PC11	PC12
Standard deviation	0.96145	0.9165	0.81413	0.76650	0.71908	0.45631
Proportion of Variance	0.05777	0.0525	0.04143	0.03672	0.03232	0.01301
Cumulative Proportion	0.80907	0.8616	0.90299	0.93971	0.97203	0.98504
	PC13	PC14	PC15	PC16		
Standard deviation	0.36522	0.3174	0.07039	0.01562		
Proportion of Variance	0.00834	0.0063	0.00031	0.00002		
Cumulative Proportion	0.99338	0.9997	0.99998	1.00000		



pr.out





Data: X dimension: 2584 15  
Y dimension: 2584 1  
Fit method: svdpc  
Number of components considered: 15


#### VALIDATION: RMSEP

Cross-validated using 10 random segments.

	(Intercept)	1 comps	2 comps	3 comps	4 comps	5 comps	6 comps
CV	0.248	0.2428	0.2411	0.2387	0.2387	0.2388	0.2389
adjCV	0.248	0.2428	0.2416	0.2387	0.2387	0.2388	0.2389
	7 comps	8 comps	9 comps	10 comps	11 comps	12 comps	13 comps
CV	0.2388	0.2388	0.2388	0.2388	0.2385	0.2381	0.2385
adjCV	0.2387	0.2387	0.2387	0.2388	0.2385	0.2380	0.2384
	14 comps	15 comps					
CV	0.2387	0.2398					
adjCV	0.2386	0.2396					

#### TRAINING: % variance explained

	1 comps	2 comps	3 comps	4 comps	5 comps	6 comps	7 comps
X	25.306	40.680	55.704	64.926	72.401	79.396	85.185
class	4.225	5.285	7.573	7.577	7.584	7.592	7.792
	8 comps	9 comps	10 comps	11 comps	12 comps	13 comps	14 comps
X	89.627	93.557	97.005	98.398	99.294	99.97	99.998
class	7.917	8.022	8.026	8.289	8.847	8.87	8.872
	15 comps						
X	100.000						
class	9.128						



[1] 0.05357258

Data: X dimension: 1938 15  
 Y dimension: 1938 1  
 Fit method: svdpc  
 Number of components considered: 15

VALIDATION: RMSEP

Cross-validated using 10 random segments.

	(Intercept)	1 comps	2 comps	3 comps	4 comps	5 comps	6 comps
CV	0.2512	0.2454	0.2417	0.2413	0.2417	0.2417	0.242
adjCV	0.2512	0.2453	0.2415	0.2413	0.2416	0.2417	0.242

	7 comps	8 comps	9 comps	10 comps	11 comps	12 comps	13 comps
CV	0.2421	0.2421	0.2421	0.2422	0.2422	0.2420	0.2422
adjCV	0.2420	0.2420	0.2420	0.2421	0.2421	0.2419	0.2421

	14 comps	15 comps
CV	0.2431	477953634
adjCV	0.2429	453530602

TRAINING: % variance explained

	1 comps	2 comps	3 comps	4 comps	5 comps	6 comps	7 comps
X	25.528	41.325	56.263	65.253	72.800	79.615	85.224
class	4.815	7.847	7.952	7.953	7.967	8.025	8.255


  

	8 comps	9 comps	10 comps	11 comps	12 comps	13 comps	14 comps
X	89.653	93.574	97.048	98.456	99.324	99.978	99.999
class	8.362	8.466	8.502	8.574	8.905	8.943	8.981


  

	15 comps
X	100.000
class	9.781

## Variable Selection - PCA - INCOMPLETE




## Variable Selection - PCA - INCOMPLETE






**PC1 vs PC3**



**PC2 vs PC3**



Data: X dimension: 2584 15  
Y dimension: 2584 1  
Fit method: svdpc  
Number of components considered: 15

# VALIDATION: RMSEP


Cross-validated using 10 random segments.


	(Intercept)	1 comps	2 comps	3 comps	4 comps	5 comps	6 comps
CV	0.248	0.2428	0.2411	0.2387	0.2387	0.2388	0.2389
adjCV	0.248	0.2428	0.2416	0.2387	0.2387	0.2388	0.2389
	7 comps	8 comps	9 comps	10 comps	11 comps	12 comps	13 comps
CV	0.2388	0.2388	0.2388	0.2388	0.2385	0.2381	0.2385
adjCV	0.2387	0.2387	0.2387	0.2388	0.2385	0.2380	0.2384
	14 comps	15 comps					
CV	0.2387	0.2398					
adjCV	0.2386	0.2396					

## TRAINING: % variance explained

	1 comps	2 comps	3 comps	4 comps	5 comps	6 comps	7 comps
X	25.306	40.680	55.704	64.926	72.401	79.396	85.185
class	4.225	5.285	7.573	7.577	7.584	7.592	7.792
	8 comps	9 comps	10 comps	11 comps	12 comps	13 comps	14 comps
X	89.627	93.557	97.005	98.398	99.294	99.97	99.998
class	7.917	8.022	8.026	8.289	8.847	8.87	8.872
	15 comps						
X	100.000						
class	9.128						

**class**





```
[1] 0.05357258
```

## Logistic Regression after Variable Selection

Call:

```
glm(formula = class ~ seismic + shift + gpuls + nbumps, family = binomial,
     data = seismic.train)
```

Deviance Residuals:

Min	1Q	Median	3Q	Max
-1.6270	-0.3846	-0.2947	-0.1627	2.9781

Coefficients:

	Estimate	Std. Error	z value	Pr(> z )
(Intercept)	-5.9508244	0.6490468	-9.169	< 2e-16 ***
seismic	0.3641160	0.1944250	1.873	0.061098 .
shift	1.1371057	0.3402674	3.342	0.000832 ***
gpuls	0.0004913	0.0001283	3.829	0.000129 ***
nbumps	0.3231048	0.0507286	6.369	1.9e-10 ***

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for binomial family taken to be 1)

Null deviance:	958.82	on 1937	degrees of freedom
Residual deviance:	828.98	on 1933	degrees of freedom

AIC: 838.98

Number of Fisher Scoring iterations: 6

[1] 0.9318885

```
glm.pred    0    1
           0 1803 128
           1    4    3
```

[1] 0.02290076


[1] 0.9977864

[1] 0.9380805

```
glm.pred    0    1
           0 606  39
           1    1    0
```

[1] 0

[1] 0.9983526



## Quadratic Discriminant Analysis after variable selection

```
##-----  
## Fit QDA model after variable selection  
##-----  
  
# Model 1  
qda.fit <- qda(class~seismic+shift+gpuls+nbumps, data=seismic.train)  
qda.class=predict(qda.fit,seismic.test)$class  
confusion <- table(qda.class ,seismic.test$class)  
  
sensitivity <- confusion[2,2]/sum(confusion[,2])  
specificity <- confusion[1,1]/sum(confusion[,1])  
  
confusion
```

```
##  
## qda.class    0    1  
##           0 565  27  
##           1  42  12
```

```
sensitivity
```

```
## [1] 0.3076923
```

```
specificity
```

```
## [1] 0.9308072
```

```
# Model 2  
qda.fit <- qda(class ~ genergy + gpuls + nbumps + nbumps2 + nbumps4, data=seismic.train)  
qda.class=predict(qda.fit,seismic.test)$class  
  
confusion <- table(qda.class ,seismic.test$class)  
  
sensitivity <- confusion[2,2]/sum(confusion[,2])  
specificity <- confusion[1,1]/sum(confusion[,1])  
  
confusion
```

```
##  
## qda.class    0    1  
##           0 527  23  
##           1  80  16
```

```
sensitivity
```

```
## [1] 0.4102564
```

```
specificity
```

```
## [1] 0.8682043
```

## Regularized Discriminant Analysis after variable selection

```
rda.class  0  1  
          0 595 35  
          1  12  4
```

```
[1] 0.1025641
```

```
[1] 0.9802306
```

```
rda.class  0  1  
          0 572 33  
          1  35  6
```

```
[1] 0.1538462
```

```
[1] 0.9423394
```